

CHAPTER 3

**The Boltzmann Equation and
Its Hydrodynamic Limits**

François Golse

*Institut Universitaire de France, and Laboratoire Jacques-Louis Lions, Université Paris 7,
Boîte Courrier 187, F75252 Paris Cedex 05, France
E-mail: golse@math.jussieu.fr*

Contents

19	1. Introduction	3	19
20	2. Fluid dynamics: A presentation of models	5	20
21	2.1. The compressible Euler system	7	21
22	2.2. The compressible Navier–Stokes system	9	22
23	2.3. The acoustic system	10	23
24	2.4. The incompressible Euler equations	11	24
25	2.5. The incompressible Navier–Stokes equations	13	25
26	2.6. The temperature equation for incompressible flows	14	26
27	2.7. Coupling of the velocity and temperature fields by conservative forces	14	27
28	3. The Boltzmann equation and its formal properties	16	28
29	3.1. Conservation laws	19	29
30	3.2. Boltzmann’s <i>H</i> -theorem	23	30
31	3.3. <i>H</i> -theorem and a priori estimates	26	31
32	3.4. Further remarks on the <i>H</i> -theorem	34	32
33	3.5. The collision kernel	37	33
34	3.6. The linearized collision integral	44	34
35	4. Hydrodynamic scalings for the Boltzmann equation	51	35
36	4.1. Notion of a rarefied gas	51	36
37	4.2. The dimensionless Boltzmann equation	53	37
38	5. Compressible limits of the Boltzmann equation: Formal results	55	38
39	5.1. The compressible Euler limit: The Hilbert expansion	55	39
40	5.2. The compressible Navier–Stokes limit: The Chapman–Enskog expansion	59	40
41	5.3. The compressible Euler limit: The moment method	63	41
42	5.4. The acoustic limit	66	42
43	6. Incompressible limits of the Boltzmann equation: Formal results	68	43
44	6.1. The incompressible Navier–Stokes limit	68	44

HANDBOOK OF DIFFERENTIAL EQUATIONS
Evolutionary Equations, volume 2
Edited by C.M. Dafermos and E. Feireisl
© 2005 Elsevier B.V. All rights reserved

1	6.2. The incompressible Stokes and Euler limits	75	1
2	6.3. Other incompressible models	78	2
3	7. Mathematical theory of the Cauchy problem for hydrodynamic models	80	3
4	7.1. The Stokes and acoustic systems	80	4
5	7.2. The incompressible Navier–Stokes equations	82	5
6	7.3. The incompressible Navier–Stokes–Fourier system	84	6
7	7.4. The compressible Euler system	85	7
8	7.5. The incompressible Euler equations	87	8
9	8. Mathematical theory of the Cauchy problem for the Boltzmann equation	90	9
10	8.1. Global classical solutions for “small” data	90	10
11	8.2. The DiPerna–Lions theory	91	11
12	8.3. Variants of the DiPerna–Lions theory	100	12
13	9. The Hilbert expansion method: Application to the compressible Euler limit	103	13
14	10. The relative entropy method: Application to the incompressible Euler limit	107	14
15	11. Applications of the moment method	111	15
16	11.1. The acoustic limit	111	16
17	11.2. The Stokes–Fourier limit	113	17
18	11.3. The Navier–Stokes–Fourier limit	115	18
19	11.4. Sketch of the proof of the Navier–Stokes–Fourier limit by the moment method	116	19
20	11.5. The nonlinear compactness estimate	131	20
21	12. Conclusions and open problems	137	21
22	References	139	22
23			23
24			24
25			25
26			26
27			27
28			28
29			29
30			30
31			31
32			32
33			33
34			34
35			35
36			36
37			37
38			38
39			39
40			40
41			41
42			42
43			43
44			44
45			45

1. Introduction


The classical models of fluid dynamics, such as the Euler or Navier–Stokes equations, were first established by applying Newton’s second law of motion to each infinitesimal volume element of the fluid considered, see, for instance, Chapter 1 of [75]. While this method has the advantage of being universal – indeed, all hydrodynamic models can be obtained in this way – it has one major drawback: equations of state and transport coefficients (such as the viscosity or heat conductivity) are given as phenomenological or experimental data, and are not related to microscopic data (essentially, to the laws governing molecular interactions). As a matter of fact, a microscopic theory of liquids is most likely too complex to be of any use in deriving the macroscopic models of fluid mechanics. In the case of gases or plasmas, however, molecular interactions are on principle much more elementary, so that one can hope to express thermodynamic functions and transport coefficients in terms of purely mechanical data concerning collisions between gas molecules.

In fact, the subject of hydrodynamic limits goes back to the work of the founders J. Clerk Maxwell and L. Boltzmann, of the kinetic theory of gases. Both checked the consistency of their new – and, at the time, controversial – theory with the well-established laws of fluid mechanics. Interestingly, while the very existence of atoms was subject to heated debates, kinetic theory would provide estimates on the size of a gas molecule from macroscopic data such as the viscosity of the gas.

Much later, D. Hilbert formulated the question of hydrodynamic limits as a mathematical problem, as an example in his 6th problem on the axiomatization of physics [68]. In Hilbert’s own words “[...] Boltzmann’s work on the principles of mechanics suggests the problem of developing mathematically the limiting processes [...] which lead from the atomistic view to the laws of motion of continua”. Some years later, Hilbert himself attacked the problem in [69], as an application of his own fundamental work on integral equations.

There is an ambiguity in Hilbert’s formulation. Indeed, what is meant by “the atomistic view” could designate two very different theories. One is molecular dynamics (i.e., the N -body problem of classical mechanics with elastic collisions, assuming for simplicity all bodies to be spherical and of equal mass). The other possibility is to start from the kinetic theory of gases, and more precisely from the Boltzmann equation, which is what Hilbert himself did in [69]. However, one should be aware that the Boltzmann equation is not itself a “first principle” of physics, but a low density limit of molecular dynamics. In the days of Maxwell and Boltzmann, and maybe even at the time of Hilbert’s own papers on the subject, this may not have been so clear to everyone. In particular, much of the controversy on irreversibility could perhaps have been avoided with a clear understanding of the relations between molecular gas dynamics and the kinetic theory of gases.

In any case, the problem of hydrodynamic limits is to obtain rigorous derivations of macroscopic models such as the fundamental partial differential equations (PDEs) of fluid mechanics from a microscopic description of matter, be it molecular dynamics or the kinetic theory of gases. The situation can be illustrated by the following diagram.



Throughout the present chapter, we are concerned with only the vertical arrow in the diagram above. As a matter of fact, this is perhaps the part of the subject that is best understood so far, at least according to the mathematical standards of rigor.

The other arrows in this diagram correspond with situations that are only partially understood, and where certain issues are still clouded with mystery. Before starting our discussion of the hydrodynamic limits of the kinetic theory of gases, let us say a few words on these other limits and direct the interested reader at the related literature.

Although beyond the scope of this chapter, the horizontal arrow is of considerable interest to our discussion, being a justification of the kinetic theory of gases on the basis of the molecular gas dynamics (viewed as a first principle of classical, nonrelativistic physics). A rigorous derivation of the Boltzmann equation from molecular dynamics on short time intervals was obtained by Lanford [77]; see also the very nice rendition of Lanford's work in the book [28]. Hence, although not a first principle itself, the Boltzmann equation is rigorously derived from first principles and therefore has more physical legitimacy than phenomenological models (such as lattice gases or stochastic Hamiltonian models). Besides, the Boltzmann equation is currently used by engineers in aerospace industry, in vacuum technology, in nuclear engineering, as well as several other applied fields, a more complete list of those being available in the Proceedings of the Rarefied Gas Dynamics Symposia.

On the other hand, "formal" derivations of the Euler system for compressible fluids from molecular dynamics were proposed by Morrey [98]. Later on, S.R.S. Varadhan and his collaborators studied the same limit, however with a different method. Instead of taking molecular dynamics as their starting point, they modified slightly the N -body Hamiltonian by adding an arbitrarily small noise term to the kinetic energy; they also cut off high velocities at a threshold compatible with the maximum speed observed on the macroscopic system. Starting from this stochastic variant of molecular gas dynamics, they derived the Euler system of compressible fluids for short times (before the onset of singularities such as shock waves); see for instance [120] and the references therein, notably [104], see also [33], and the more recent reference [44]. The role of the extra noise term in their derivation is to guarantee some form of the ergodic principle, i.e., that the only invariant measure for the Hamiltonian in the limit of infinitely many particles is a local Gibbs state (parametrized by macroscopic quantities). At the time of this writing, deriving the Euler system of compressible fluids from molecular gas dynamics without additional noise terms as in [104] and for all positive times seems beyond reach.

For these reasons, we have limited our discussion to only the derivation of hydrodynamic models from the kinetic theory of gases, i.e., from the Boltzmann equation. For a

1 more general view of the subject of hydrodynamic limits, the reader is advised to read the 1
 2 excellent survey article by Esposito and Pulvirenti [41], whose selection of topics is quite 2
 3 different from ours. 3

4 This chapter is organized as follows: in Section 2 we review the classical models of fluid 4
 5 mechanics. Section 3 introduces the Boltzmann equation and discusses its structure and 5
 6 main formal properties. In Section 4 we discuss the dimensionless form of the Boltzmann 6
 7 equation and introduce its main scaling parameters. Sections 5 and 6 explain in detail 7
 8 the formal derivation of the most classical PDEs of fluid mechanics from the Boltzmann 8
 9 equation by several different methods. Section 7 recalls the known mathematical results 9
 10 on the Cauchy problem for the PDEs of fluid mechanics. In Section 8 we review the state 10
 11 of the art on the existence theory for the Boltzmann equation. Sections 9–11 sketch the 11
 12 mathematical proofs of the formal derivations described in Sections 6 and 7; here again, 12
 13 we present three different methods for establishing these hydrodynamic limits and discuss 13
 14 their respective merits. 14

15 We have chosen to emphasize compactness methods, leading to global results, and espe- 15
 16 cially the derivation of global weak solutions of the incompressible Navier–Stokes equa- 16
 17 tions from renormalized solutions of the Boltzmann equation. There is more than a simple 17
 18 matter of taste in this choice. Indeed, it is a nontrivial question to decide whether these 18
 19 hydrodynamic limits are intrinsic properties of the microscopic versus macroscopic mod- 19
 20 els governing the dynamics of gases, or simply an illustration of more or less standard 20
 21 techniques in asymptotic analysis. The second viewpoint leads to derivations of hydrody- 21
 22 namic models that fall short of describing any singular behavior beyond isolated shock 22
 23 waves in compressible gas dynamics. The first viewpoint uses the specific structure of the 23
 24 Boltzmann equation to design convergence proofs that are based on only the a priori esti- 24
 25 mates on this equation that have an intrinsic physical meaning; these convergence proofs 25
 26 are insensitive to whether singularities appear in finite time on the limiting hydrodynamic 26
 27 model. 27
 28
 29

30 2. Fluid dynamics: A presentation of models 30

31
 32 Usually, one thinks of a fluid – more generally, a continuous medium – as a set of material 32
 33 points which, at any given time t , fill a smooth domain in the Euclidean space \mathbb{R}^N , where 33
 34 $N = 1, 2, 3$ are the dimensions of physical interest. 34
 35

36 The purpose of fluid dynamics is to describe the state of the fluid at any instant of time 36
 37 with a *small* number of fields – such as the velocity or temperature fields – defined on the 37
 38 domain filled by the fluid. 38

39 These fields are governed by several partial differential equations that share a common 39
 40 structure which we briefly recall below. 40

41 Consider the motion of a continuous medium, and denote by 41

$$42 X(t, s; a) \in \mathbb{R}^N \quad 42$$

43
 44
 45 the position at time t of the material point which occupied position a at time s . 45

The kinematics of such a medium is based on the *parallel transport* along the family of curves $t \mapsto X(t, s; a)$ indexed by a (s , being the origin of times, is kept fixed). The infinitesimal description of this parallel transport involves the first-order differential operator

$$\frac{D}{Dt} = \partial_t + u(t, x) \cdot \nabla_x = \partial_t + \sum_{j=1}^N u_j(t, x) \partial_{x_j},$$

where the velocity field $u(t, x)$ is defined in terms of the particle paths $X(t, s; a)$ by the formula

$$\frac{d}{dt} X(t, s; a) = u(t, X(t, s; a)).$$

The operator $\frac{D}{Dt}$ is usually called *the material derivative*, and using it allows one to eliminate the trajectories $X(t, s; a)$. In other words, instead of following the motion of each material point, one looks at any fixed point in the Euclidean space \mathbb{R}^N , say x , and observes, at any given time t , the velocity $u(t, x)$ of the material point that is located at the position x at time t . This is called the *Eulerian description* of a continuous medium, whereas the description in terms of $X(t, s; a)$ is called the *Lagrangian description*. Interestingly, the connections between the kinetic theory of gases (or plasmas) and fluid dynamics are always formulated in terms of the Eulerian, instead of the Lagrangian description, although the latter may seem more natural when dealing with the motion of a gas at the atomic or molecular level.

Fluid dynamics rests on three fundamental laws – or equations:

- the continuity equation,
- the motion equation, and
- the energy balance equation.

The continuity equation states that the density ρ of the fluid is transported by the flow, i.e., that the measure $\rho(t, x) dx$ is the image of the measure $\rho(s, a) da$ under the map $a \mapsto X(t, s; a)$. The infinitesimal formulation of this fact is

$$\frac{D\rho}{Dt} = -\rho \operatorname{div}_x u. \quad (2.1)$$

The motion equation states that each portion of the fluid obeys Newton's second law of motion (i.e., $\frac{d}{dt}(\text{momentum}) = \text{force}$). The acceleration is computed in terms of the material derivative, and the infinitesimal formulation of the motion equation is

$$\rho \frac{Du}{Dt} = \operatorname{div}_x S + \rho \mathbf{f}, \quad (2.2)$$

where \mathbf{f} is the external force field (e.g., gravity, Lorentz force in the case of a plasma...) and S is the stress tensor. The meaning of S is as follows: at any given time t , isolate a smooth domain Ω in the fluid, denote by $\partial\Omega$ its boundary, by n_x the unit normal field

on $\partial\Omega$ pointing toward the outside of Ω , and by $d\sigma(x)$ the surface element on $\partial\Omega$. Then, the force exerted by the fluid outside Ω on the fluid inside Ω is

$$\int_{\partial\Omega} S(t, x) n_x d\sigma(x).$$

Finally, the energy balance equation involves the internal energy of the fluid per unit of mass E ; the total energy per unit of mass is $\frac{1}{2}|u|^2 + E$ (the sum of the kinetic energy and the internal energy). It states that the material derivative of the total energy of any portion of fluid is the sum of the works of the stresses and of the external force \mathbf{f} , minus the heat flux lost by that portion of fluid. Its infinitesimal formulation is

$$\rho \frac{D}{Dt} \left(\frac{1}{2}|u|^2 + E \right) = -\operatorname{div}_x Q + \operatorname{div}_x (Su) + \rho \mathbf{f} \cdot u, \quad (2.3)$$

where Q is the heat flux.

In the motion and energy balance equations, \mathbf{f} is a given vector field, while the density ρ , the velocity field u , the internal energy E , the stress tensor S and the heat flux Q are unknown. However, these quantities are usually not independent, but are related by *equations of state* that depend on the fluid considered.

Equations (2.1)–(2.3) are *Galilean invariant*. Specifically, let $v \in \mathbb{R}^3$; define the Galilean transformation

$$x' = x + vt, \quad u'(t, x') = u(t, x) + v, \quad \phi'(t, x') = \phi(t, x)$$

for $\phi = \rho, S, \mathbf{f}, E, Q$. Then, setting

$$\frac{D'}{Dt} = \partial_t + u' \cdot \nabla_{x'}$$

one deduces from (2.1)–(2.3) that

$$\frac{D'}{Dt} \rho' = -\rho' \operatorname{div}_{x'} u',$$

$$\rho' \frac{D'}{Dt} u' = \operatorname{div}_{x'} S' + \rho' \mathbf{f}',$$

$$\rho' \frac{D'}{Dt} \left(\frac{1}{2}|u'|^2 + E' \right) = -\operatorname{div}_{x'} Q' + \operatorname{div}_{x'} (S' u') + \rho' \mathbf{f}' \cdot u'.$$

2.1. The compressible Euler system

An *ideal fluid* is one where the effects of viscosity and thermal conductivity can be neglected. In this case, $Q = 0$ and the stress tensor is of the form

$$S = -pI,$$

1 where the scalar p is the pressure. Hence the system consisting of the continuity equation 1
2 the motion equation and the energy balance equation becomes 2

$$\begin{aligned}
 3 \quad & \partial_t \rho + \operatorname{div}_x(\rho u) = 0, \\
 4 \quad & \rho(\partial_t u + (u \cdot \nabla_x)u) = -\nabla_x p + \rho \mathbf{f}, \\
 5 \quad & \rho(\partial_t E + (u \cdot \nabla_x)E) = -p \operatorname{div}_x u + \rho \mathbf{f} \cdot u.
 \end{aligned}
 \tag{2.4}$$

9 Thus, the unknowns are the density ρ , the velocity field u , the pressure p and the internal 9
10 energy E . However, the quantities ρ , p and E are not independent, but are related by 10
11 *equations of state*. 11

12 Choosing the density ρ and the temperature θ as independent thermodynamic variables, 12
13 these equations of state are relations that express the pressure p and the internal energy E 13
14 in terms of ρ and θ 14

$$p \equiv p(\rho, \theta), \quad E \equiv E(\rho, \theta). \tag{2.5}$$

18 Hence (2.4) is a system of $N + 2$ partial differential equations for the unknowns ρ , u and θ ; 18
19 notice that there are in fact $N + 2$ scalar unknowns, ρ and θ , plus the N components of the 19
20 vector field u . 20

21 The case of a perfect gas is of particular importance for the rest of this chapter. In this 21
22 case, the equations of state are 22

$$p(\rho, \theta) = k\rho\theta, \quad e(\rho, \theta) = \frac{k\theta}{\gamma - 1}, \tag{2.6}$$

27 where k is the Boltzmann constant ($k = 1.38 \cdot 10^{-23} \text{ J K}^{-1}$) and $\gamma > 1$ is a constant called 27
28 the adiabatic exponent. For a perfect gas whose molecules have n degrees of freedom 28
29

$$\gamma = 1 + \frac{2}{n}.$$

33 For instance, in the case of a perfect monatomic gas, each molecule has 3 degrees of free- 33
34 dom (the coordinates of its center of mass); hence $\gamma = 5/3$. In the case of a diatomic gas, 34
35 each molecule has 5 degrees of freedom (the coordinates of its center of mass and the 35
36 direction of the line passing through the centers of both atoms); hence $\gamma = 7/5$. 36

37 From now on, we choose a temperature scale such that $k = 1$. 37

38 Adding the continuity equation to the motion equation and to the energy balance equa- 38
39 tion, one can recast (2.4) in the form 39

$$\begin{aligned}
 41 \quad & \partial_t \rho + \operatorname{div}_x(\rho u) = 0, \\
 42 \quad & \partial_t(\rho u) + \operatorname{div}_x(\rho u \otimes u) + \nabla_x(\rho\theta) = \rho \mathbf{f}, \\
 43 \quad & \partial_t \left(\rho \left(\frac{1}{2} |u|^2 + \frac{1}{\gamma - 1} \theta \right) \right) + \operatorname{div}_x \left(\rho u \left(\frac{1}{2} |u|^2 + \frac{\gamma}{\gamma - 1} \theta \right) \right) = \rho \mathbf{f} \cdot u.
 \end{aligned}
 \tag{2.7}$$

This is a degenerate parabolic system of partial differential equations in the unknowns ρ , u and θ . Observe that there is no diffusion term in the first equation, which is clear on physical grounds. Indeed, the meaning of the continuity equation is purely geometric – namely, the fact that the measure ρdx is transported by the fluid flow – and cannot be affected by physical assumptions on the fluid (such as whether the fluid is ideal or not).

2.3. The acoustic system

The acoustic waves in an ideal fluid are small amplitude disturbances of a constant equilibrium state. Therefore the propagation of acoustic waves is governed by the linearization at a constant state $(\bar{\rho}, \bar{u}, \bar{\theta})$ of the compressible Euler system. Without loss of generality, one can assume by Galilean invariance that $\bar{u} = 0$. The density, velocity and temperature fields are written as

$$\rho = \bar{\rho} + \tilde{\rho}, \quad u = \tilde{u}, \quad \theta = \bar{\theta} + \tilde{\theta},$$

where the letters adorned with tildes designate small disturbances of the background equilibrium state $(\bar{\rho}, 0, \bar{\theta})$. In the case of a perfect gas, and in the absence of external force (i.e., for $\mathbf{f} = 0$), the acoustic system takes the form

$$\begin{aligned} \partial_t \tilde{\rho} + \bar{\rho} \operatorname{div}_x \tilde{u} &= 0, \\ \partial_t \tilde{u} + \frac{\bar{\theta}}{\bar{\rho}} \nabla_x \tilde{\rho} + \nabla_x \tilde{\theta} &= 0, \\ \frac{1}{\gamma - 1} \partial_t \tilde{\theta} + \bar{\theta} \operatorname{div}_x \tilde{u} &= 0. \end{aligned} \tag{2.9}$$

By combining the first and the last equation in the system above, one can put it in the form

$$\begin{aligned} \partial_t \left(\frac{\tilde{\rho}}{\bar{\rho}} + \frac{\tilde{\theta}}{\bar{\theta}} \right) + \gamma \operatorname{div}_x \tilde{u} &= 0, \\ \partial_t \tilde{u} + \bar{\theta} \nabla_x \left(\frac{\tilde{\rho}}{\bar{\rho}} + \frac{\tilde{\theta}}{\bar{\theta}} \right) &= 0. \end{aligned} \tag{2.10}$$

Splitting the fluctuation of velocity field \tilde{u} as the sum of a gradient field and of a *solenoidal* (i.e., divergence-free) field

$$\tilde{u} = -\nabla_x \varphi + \tilde{u}^s, \quad \operatorname{div}_x \tilde{u}^s = 0,$$

one deduces from the system (2.10) – together with boundary conditions, or conditions at infinity, or else conditions on the mean value of the fields, whose detailed description does

1 not belong here – that

$$\begin{aligned}
 2 & \\
 3 & (\partial_{tt} - \gamma \bar{\theta} \Delta_x) \left(\frac{\tilde{\rho}}{\bar{\rho}} + \frac{\tilde{\theta}}{\bar{\theta}} \right) = 0, \\
 4 & \\
 5 & (\partial_{tt} - \gamma \bar{\theta} \Delta_x) \varphi = 0, \\
 6 & \\
 7 & \partial_t u^s = 0. \\
 8 &
 \end{aligned} \tag{2.11}$$

9 In other words, the acoustic system can be reduced to two independent wave equations
10 for $(\tilde{\rho}/\bar{\rho} + \tilde{\theta}/\bar{\theta})$ (the relative pressure fluctuation) and φ (the fluctuating stream function),
11 while the solenoidal part of the velocity fluctuation u^s is a constant of motion.

14 2.4. The incompressible Euler equations

15
16 Consider next the case of an incompressible, homogeneous ideal fluid. The evolution of
17 such a fluid is governed by the system (2.4) with $\rho = \text{const}$. The continuity and motion
18 equations in (2.4) reduce to

$$\begin{aligned}
 19 & \\
 20 & \text{div}_x u = 0, \\
 21 & \\
 22 & \partial_t u + (u \cdot \nabla_x) u = -\nabla_x \pi + \mathbf{f}, \\
 23 &
 \end{aligned} \tag{2.12}$$

24 where $\pi = p/\rho$. At variance with the compressible Euler system, there is no need of an
25 equation of state to determine π . Indeed, taking the divergence of both sides of the motion
26 equation leads to

$$27 \quad -\Delta_x \pi = \text{div}_x (u \cdot \nabla_x u) - \text{div}_x \mathbf{f} = \text{trace}((\nabla_x u)^2) - \text{div}_x \mathbf{f},$$

28
29 so that π can be expressed in terms of u by solving the Laplace equation. In other words,
30 π must be thought of as the Lagrange multiplier associated to the constraint $\text{div}_x u = 0$.

31 The incompressible Euler equations arise in a different context, namely in the description
32 of *incompressible flows* of compressible fluids (such as perfect gases, for instance).

33 The dimensionless number that monitors the compressibility is the *Mach number*, i.e.,
34 the ratio of the length of the velocity field to the speed of sound. In the case of a perfect
35 gas with adiabatic exponent γ , our discussion of the acoustic system above shows that the
36 speed of sound in the gas at a temperature θ is $c = \sqrt{\gamma \theta}$, so that the Mach number in that
37 case is

$$38 \quad Ma = \frac{|u|}{\sqrt{\gamma \theta}}. \tag{2.13}$$

39
40
41
42 With this definition, the Mach number is a local quantity, since u and θ are in general
43 functions of x and t . But one can replace $|u|$ and θ in the definition above by constant
44 quantities of the same order of magnitude, for instance by averages of $|u|$ and θ over large
45 spatial and temporal domains.

1 in the limit as $\varepsilon \rightarrow 0$. This discussion suggests that, in the small Mach number limit, flows 1
 2 of a compressible fluid such as a perfect gas are well described by the incompressible Euler 2
 3 equations. 3
 4

5 2.5. The incompressible Navier–Stokes equations 6 7

8
 9 Next, we start from the compressible Navier–Stokes system (2.8), and assume that the 9
 10 density ρ is a constant. As above, the continuity equation in (2.8) reduces to the incom- 10
 11 pressibility condition $\operatorname{div}_x u = 0$. Moreover, assuming that the viscosity λ is a constant, we 11
 12 find that the momentum equation reduces to 12
 13

$$14 \quad \rho(\partial_t u + (u \cdot \nabla_x)u) + \nabla_x p = \rho \mathbf{f} + \lambda \Delta_x u. \quad 14$$

15
 16 Defining the kinematic viscosity to be 16
 17

$$18 \quad \nu = \frac{\lambda}{\rho} \quad 18$$

19
 20 and setting $\pi = p/\rho$, we arrive at the incompressible Navier–Stokes equations 20
 21

$$22 \quad \operatorname{div}_x u = 0, \quad 22$$

$$23 \quad \partial_t u + (u \cdot \nabla_x)u + \nabla_x \pi = \mathbf{f} + \nu \Delta_x u. \quad 23$$

24
 25 We leave it to the reader to verify that the incompressible Navier–Stokes equations can be 24
 26 viewed as the small Mach number limit of the compressible Navier–Stokes system, as was 25
 27 done in the case of the incompressible Euler system. The scaling law is slightly different 26
 28 from the Euler case: for $\varepsilon = \sqrt{Ma}$, set 27
 28

$$29 \quad \rho_\varepsilon(t, x) = \rho\left(\frac{t}{\varepsilon^2}, \frac{x}{\varepsilon}\right), \quad 29$$

$$30 \quad u_\varepsilon(t, x) = \frac{1}{\varepsilon} u\left(\frac{t}{\varepsilon^2}, \frac{x}{\varepsilon}\right), \quad 30$$

$$31 \quad \theta_\varepsilon(t, x) = \theta\left(\frac{t}{\varepsilon^2}, \frac{x}{\varepsilon}\right), \quad 31$$

32
 33 where (ρ, u, θ) is a solution to the Navier–Stokes system (2.8), with $\mathbf{f} \equiv 0$. Then, to leading 32
 34 order as $\varepsilon \rightarrow 0$, u_ε satisfies (2.16) with $\mathbf{f} \equiv 0$. 33
 35

36
 37 So far, we have said nothing about the temperature field in incompressible flows; this 34
 38 will be the subject matter of the next subsection. 35
 39
 40
 41
 42
 43
 44
 45

2.6. The temperature equation for incompressible flows

Going back to the Navier–Stokes system (2.8) for a perfect gas with adiabatic exponent γ , we see that, in the incompressible case where $\rho = \text{const}$, the third equation reduces to

$$\frac{1}{\gamma - 1} \rho (\partial_t \theta + u \cdot \nabla_x \theta) = \text{div}_x (\kappa(\theta) \nabla_x \theta) + \frac{1}{2} \lambda D(u) : D(u). \quad (2.18)$$

On the right-hand side of (2.18), the first term represents the divergence of the heat flux due to thermal conduction, as described by Fourier’s law, while the second term represents the production of heat by intermolecular friction and is called the *viscous heating* term.

In some models that can be found in the literature, the viscous heating term is absent from the temperature equation. Whether the viscous heating term should be taken into account or not depends in fact on the relative size of the fluctuations of velocity field about its average value, and of the fluctuations of temperature field about its average values.

If the fluctuations of velocity field are of a smaller order than the square-root of the temperature fluctuations, then a straightforward scaling argument shows that the viscous heating term can indeed be neglected in (2.18). If however, the fluctuations of velocity field are at least of the same order of magnitude as the square-root of the temperature fluctuations, then the viscous heating term cannot be neglected in (2.18). We shall discuss this alternative further in the description of the incompressible hydrodynamic limits of the Boltzmann equation.

2.7. Coupling of the velocity and temperature fields by conservative forces

In our discussion of the incompressible flows as low Mach number limits, we have neglected so far the external force \mathbf{f} . Split it as the sum of a gradient field (i.e., of a conservative force) and of a solenoidal field

$$\mathbf{f} = -\nabla_x \phi + \mathbf{f}^s, \quad \text{div}_x \mathbf{f}^s = 0.$$

Scale ϕ and \mathbf{f}^s as

$$\phi_\varepsilon(t, x) = \frac{1}{\varepsilon} \phi\left(\frac{t}{\varepsilon^2}, \frac{x}{\varepsilon}\right), \quad \mathbf{f}_\varepsilon^s(t, x) = \frac{1}{\varepsilon^3} \mathbf{f}^s\left(\frac{t}{\varepsilon^2}, \frac{x}{\varepsilon}\right), \quad (2.19)$$

and assume that ρ_ε and θ_ε have fluctuations of order ε about their constant average values

$$\rho_\varepsilon = \bar{\rho} + \varepsilon \tilde{\rho}_\varepsilon, \quad \theta_\varepsilon = \bar{\theta} + \varepsilon \tilde{\theta}_\varepsilon.$$

In that case, the leading order in ε of the momentum equation in the Navier–Stokes system (2.8) reduces to

$$\nabla_x (\bar{\rho} \tilde{\theta}_\varepsilon + \bar{\theta} \tilde{\rho}_\varepsilon) + \bar{\rho} \nabla_x \phi_\varepsilon \simeq 0,$$

1 or in other words, 1

$$2 \frac{\tilde{\rho}_\varepsilon}{\bar{\rho}} + \frac{\tilde{\theta}_\varepsilon}{\bar{\theta}} + \frac{\phi_\varepsilon}{\bar{\theta}} \simeq C(t), \quad (2.20) \quad 3$$

4 an equality known as *Boussinesq's relation*. In many cases, the boundary conditions (or 5
6 decay at infinity, or else periodicity conditions) entail that $C(t) = 0$. 6

7 The next order in ε of the momentum equation in the Navier–Stokes system is 7
8

$$9 \bar{\rho}(\partial_t u_\varepsilon + (u_\varepsilon \cdot \nabla_x)u_\varepsilon) + \tilde{\rho}_\varepsilon \nabla_x \phi_\varepsilon \simeq \lambda \Delta_x u_\varepsilon + \bar{\rho} \mathbf{f}_\varepsilon + \text{gradient field} \quad 9$$

10 and one expresses the action of the conservative force $\tilde{\rho}_\varepsilon \nabla_x \phi_\varepsilon$ as 10
11

$$12 \tilde{\rho}_\varepsilon \nabla_x \phi_\varepsilon \simeq -\frac{\bar{\rho}}{\bar{\theta}} \left(\tilde{\theta}_\varepsilon \nabla_x \phi_\varepsilon + \frac{1}{2} \nabla_x (\phi_\varepsilon^2) \right) \quad 12$$

13 so that the momentum equation reduces to 13
14

$$15 \partial_t u_\varepsilon + (u_\varepsilon \cdot \nabla_x)u_\varepsilon - \frac{\theta_\varepsilon}{\bar{\theta}} \nabla_x \phi_\varepsilon \simeq \nu \Delta_x u_\varepsilon + \mathbf{f}_\varepsilon + \text{gradient field}. \quad (2.21) \quad 15$$

16 As for the temperature equation, one should refrain from using directly (2.18). Indeed, this 16
17 equation has been derived from (2.8) in the purely incompressible case where $\rho = \text{const}$, 17
18 while in the present case $\rho = \text{const}$ modulo terms of order ε . 18

19 In the present case, we must go back to the Navier–Stokes system (2.8) and write the 19
20 continuity and energy equation in terms of the fluctuations of density and temperature 20
21

$$22 \varepsilon(\partial_t \tilde{\rho}_\varepsilon + u_\varepsilon \cdot \nabla_x \tilde{\rho}_\varepsilon) + \bar{\rho} \operatorname{div}_x u_\varepsilon = \mathfrak{o}(\varepsilon), \quad 22$$

$$23 \frac{1}{\gamma - 1} \varepsilon(\partial_t \tilde{\theta}_\varepsilon + u_\varepsilon \cdot \nabla_x \tilde{\theta}_\varepsilon) + \bar{\theta} \operatorname{div}_x u_\varepsilon = \varepsilon \operatorname{div}_x \left(\frac{\kappa(\bar{\theta})}{\bar{\rho}} \nabla_x \tilde{\theta}_\varepsilon \right) + \mathfrak{o}(\varepsilon). \quad 23$$

24 Next we must eliminate $\operatorname{div}_x u_\varepsilon$ between both equations above; indeed, we only know that 24
25 $\operatorname{div}_x u_\varepsilon \simeq 0$ to leading order in ε , and it may not be true that $\operatorname{div}_x u_\varepsilon = \mathfrak{o}(\varepsilon)$. Dividing the 25
26 first equation above by $\bar{\rho}$ and the second by $\bar{\theta}$, one arrives at 26
27

$$28 \varepsilon(\partial_t + u_\varepsilon \cdot \nabla_x) \left(\frac{1}{\gamma - 1} \frac{\tilde{\theta}_\varepsilon}{\bar{\theta}} - \frac{\tilde{\rho}_\varepsilon}{\bar{\rho}} \right) = \varepsilon \frac{\kappa(\bar{\theta})}{\bar{\rho}} \Delta_x \left(\frac{\tilde{\theta}_\varepsilon}{\bar{\theta}} \right) + \mathfrak{o}(\varepsilon). \quad 28$$

29 We further eliminate the fluctuation of density by Boussinesq's relation, and eventually 29
30 arrive at 30
31

$$32 \frac{\gamma}{\gamma - 1} \bar{\rho}(\partial_t \theta_\varepsilon + u_\varepsilon \cdot \nabla_x \theta_\varepsilon) + \bar{\rho} u_\varepsilon \cdot \nabla_x \phi_\varepsilon \simeq \kappa(\bar{\theta}) \Delta_x \theta_\varepsilon. \quad (2.22) \quad 32$$

1 Collecting both equations (2.21) and (2.22), we arrive at the coupled system in the small
2 ε limit

$$\begin{aligned} 3 & \partial_t u + u \cdot \nabla_x u + \frac{\theta}{\bar{\rho}} \nabla_x \phi + \nabla_x \pi = \nu \Delta_x u, & \operatorname{div}_x u &= 0, \\ 4 & \partial_t \theta + u \cdot \nabla_x \theta + \frac{\gamma - 1}{\gamma} u \cdot \nabla_x \phi = \bar{\kappa} \Delta_x \theta, \end{aligned} \quad (2.23)$$

5 where

$$6 \quad \bar{\kappa} = \frac{\gamma - 1}{\gamma} \frac{\kappa(\bar{\theta})}{\bar{\rho}}.$$

7 It is interesting to compare the temperature equation in (2.23) with (2.18); notice that the
8 heat conductivity in (2.23) is $1/\gamma$ that in (2.18). Besides there is no viscous heating term in
9 the temperature equation in (2.23), at variance with (2.18). This, however, is a consequence
10 of the scaling considered in the discussion above: indeed, the fluctuations of velocity and
11 temperature fields are of the same order of magnitude, so that viscous heating is a lower-
12 order effect. On the contrary, if one sets the temperature fluctuations to be of the order of
13 the squared fluctuations of velocity field, one recovers a viscous heating term in (2.23).

14 The material in this section is fairly classical and can be found in most textbooks on
15 fluid mechanics; more information can be gathered from the excellent introductory section
16 of [86]; see also the classical treatise [75].

17 3. The Boltzmann equation and its formal properties

18 The Boltzmann equation is the model that governs the evolution of perfect gases in kinetic
19 theory. While fluid dynamics describes the state of a fluid with a few scalar or vector fields
20 defined on the domain filled by the fluid, such as the temperature or velocity fields, kinetic
21 theory describes the state of a gas with the *number density* (also called the *distribution*
22 *function*) $F \equiv F(t, x, v) \geq 0$ that is the density of gas molecules which, at time $t \geq 0$,
23 are located at the position $x \in \mathbb{R}^3$ and have velocity $v \in \mathbb{R}^3$. Put in other words, in any
24 infinitesimal volume $dx dv$ centered at the point $(x, v) \in \mathbb{R}^3 \times \mathbb{R}^3$ of the single particle
25 phase space, one can find approximately $F(t, x, v) dx dv$ like particles at time t . In the
26 classical kinetic theory of gases, the molecular radius is neglected, except in the collision
27 cross-section: this has important consequences, as will be seen later.


28 The Boltzmann equation takes the form

$$29 \quad \partial_t F + v \cdot \nabla_x F = \mathcal{B}(F, F), \quad (3.1)$$

30 where $\mathcal{B}(F, F)$ is the collision integral. This collision integral $\mathcal{B}(F, F)$ is a quadratic inte-
31 gral operator acting only on the v -argument of the number density F , and takes the form

$$32 \quad \mathcal{B}(F, F)(t, x, v) = \iint_{\mathbb{R}^3 \times \mathbb{S}^2} (F' F'_* - F F_*) b(v - v_*, \omega) d\omega dv_*, \quad (3.2)$$

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

Fig. 1. The pre- and post-collision velocities in the reference frame of the center of mass of the particle pair.

where the notations F , F_* , F' and F'_* designate respectively the values $F(t, x, v)$, $F(t, x, v_*)$, $F(t, x, v')$ and $F(t, x, v'_*)$, with $v' \equiv v'(v, v_*, \omega)$ and $v'_* \equiv v'_*(v, v_*, \omega)$ given in terms of v , v_* and ω by the formulas

$$v' = v - (v - v_*) \cdot \omega \omega, \quad v'_* = v_* + (v - v_*) \cdot \omega \omega, \quad (3.3)$$

where $\omega \in \mathbb{S}^2$ is an arbitrary unit vector. These formulas represent all the solutions $(v', v'_*) \in \mathbb{R}^3 \times \mathbb{R}^3$ of the system of equations

$$v' + v'_* = v + v_*, \quad |v'|^2 + |v'_*|^2 = |v|^2 + |v_*|^2, \quad (3.4)$$

where $(v, v_*) \in \mathbb{R}^3 \times \mathbb{R}^3$ is given. If (v, v_*) are the velocities of a pair of like particles before collision, and (v', v'_*) are the velocities of the same pair of particles after collision – or vice versa, equalities (3.4) express the conservation of momentum and kinetic energy during the collision.

Only the binary collisions are accounted for in Boltzmann's equation. Indeed, since the molecular radius is neglected in the kinetic theory of gases, one can show that collisions involving more than two particles are events that occur with probability zero, and therefore can be neglected for all practical purposes.

Moreover, kinetic energy is the only form of energy conserved during collisions. In fact, the Boltzmann collision integral (3.2) applies only to monatomic gases. Polyatomic gases can also be treated by the methods of kinetic theory; however this require using complicated variants of Boltzmann's original collision integral that involve vibrational and rotational energies in addition to the kinetic energy of the center of mass of each molecule; besides, these additional energy variables are quantized in certain applications. While such considerations are important for understanding some real gas effects, they lead to heavy

1 technicalities which do not belong to an expository article such as the present one. For these 1
 2 reasons, we shall implicitly restrict our attention to monatomic gases and to the collision 2
 3 integral (3.2) in the sequel. 3

4 The function $b \equiv b(V, \omega)$ is the *collision kernel*, an a.e. positive function that is of the 4
 5 form 5

$$6 \quad b(V, \omega) = |V| \Sigma(|V|, |\cos(\widehat{V, \omega})|), \quad (3.5) \quad 6$$

7 where Σ is the *scattering cross-section* (see Section 3.5 for a precise definition of this 7
 8 notion). 8

9 We shall discuss later the physical meaning of the function Σ , together with the usual 9
 10 mathematical assumptions on the collision kernel b . For the moment, assume that b is 10
 11 locally integrable on $\mathbb{R}^3 \times \mathbb{S}^2$, and consider a number density $F \equiv F(t, x, v)$ which, at 11
 12 any arbitrary instant of time t and location x , is continuous with compact support in the 12
 13 velocity variable v . Then, the collision integral $\mathcal{B}(F, F)(t, x, v)$ can be split as 13
 14
 15
 16

$$17 \quad \mathcal{B}(F, F)(t, x, v) = \mathcal{B}_+(F, F)(t, x, v) - \mathcal{B}_-(F, F)(t, x, v), \quad (3.6) \quad 17$$

18 where 18
 19

$$20 \quad \begin{aligned} 21 \quad \mathcal{B}_+(F, F)(t, x, v) &= \iint_{\mathbb{R}^3 \times \mathbb{S}^2} F' F'_* b(v - v_*, \omega) \, d\omega \, dv_*, & 21 \\ 22 \quad \mathcal{B}_-(F, F)(t, x, v) &= \iint_{\mathbb{R}^3 \times \mathbb{S}^2} F F_* b(v - v_*, \omega) \, d\omega \, dv_*. & 22 \end{aligned} \quad (3.7) \quad 23$$

24 are called respectively the *gain term* and the *loss term* in the collision integral $\mathcal{B}(F, F)$. 24
 25 The physical meaning of both the gain and loss terms – and that of the collision integral 25
 26 itself – can be explained in the following manner: 26

- 27 • $\mathcal{B}_-(F, F)(t, x, v) \, dv$ is the number of particles located at x at time t that exit the vol- 27
 28 ume element dv centered at v in the velocity space by colliding with another particle 28
 29 with an arbitrary velocity v_* located at the same position x at the same time t , and 29
- 30 • $\mathcal{B}_+(F, F)(t, x, v) \, dv$ is the number of particles located at x at time t that enter the 30
 31 volume element dv centered at v in the velocity space as the result of a collision 31
 32 involving two particles with pre-collisional velocities v' and v'_* at the same time t and 32
 33 the same position x . 33
 34
 35
 36

37 Notice that, in this model, collisions are purely local and instantaneous, which is another 37
 38 consequence of having neglected the molecular radius. Moreover, it is assumed that the 38
 39 joint distribution of any pair of particles located at the same position x at the same time t 39
 40 with velocities v and v_* and that are about to collide is the product 40

$$41 \quad F(t, x, v) F(t, x, v_*). \quad 41$$

42 In other words, such particles are assumed to be statistically uncorrelated; however, this as- 42
 43 sumption, which is crucial in the physical derivation of the Boltzmann equation, is needed 43
 44
 45

1 only for particle pairs about to collide – and is obviously false for a pair of particles having
2 just collided.

3 Going back to the Boltzmann equation (3.1) in the form

$$4 \quad \partial_t F = -v \cdot \nabla_x F + \mathcal{B}(F, F),$$

7 it follows from the above discussion that the first term on the right-hand side represents
8 the net number of particles entering the infinitesimal phase-space volume $dx dv$ centered
9 at (x, v) as the result of inertial motion of particles between collisions, while the second
10 term represents the net number of particles entering that same volume as the result of
11 instantaneous and purely local collisions.

14 3.1. Conservation laws

16 Throughout this subsection, it is assumed that the collision kernel b is locally integrable,

$$18 \quad b \in L^1_{\text{loc}}(\mathbb{R}^3 \times \mathbb{S}^2). \quad (3.8)$$

20 The first major result about the Boltzmann collision integral is the following proposition.

22 PROPOSITION 3.1. *Let $F \equiv F(v) \in C_c(\mathbb{R}^3)$ and $\phi \in C(\mathbb{R}^3)$. Then*

$$24 \quad \int_{\mathbb{R}^3} \mathcal{B}(F, F)(v) \phi(v) dv$$

$$26 \quad = \frac{1}{4} \iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F' F'_* - F F_*) (\phi + \phi_* - \phi' - \phi'_*) b(v - v_*, \omega) dv dv_* d\omega.$$

30 This result is essential to understanding the Boltzmann equation and especially its rela-
31 tions to hydrodynamics. For this reason, we shall give a complete proof of it.

33 PROOF OF PROPOSITION 3.1. The second relation in (3.4) and the fact that F is com-
34 pactly supported shows that the support of $(v, v_*, \omega) \mapsto F' F'_* - F F_*$ is compact in
35 $\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2$. Hence both integrals

$$37 \quad \frac{1}{4} \iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F' F'_* - F F_*) (\phi + \phi_* - \phi' - \phi'_*) b(v - v_*, \omega) dv dv_* d\omega$$

40 and

$$42 \quad \int_{\mathbb{R}^3} \mathcal{B}(F, F)(v) \phi(v) dv$$

$$44 \quad = \iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F' F'_* - F F_*) \phi b(v - v_*, \omega) dv dv_* d\omega$$

are well defined since F and ϕ are continuous and b satisfies (3.8). In the latter integral, apply the change of variables $(v, v_*) \mapsto (v_*, v)$, while keeping ω fixed: (3.3) show that (v', v'_*) is changed into (v'_*, v') , so that the expression $(F'F'_* - FF_*)$ is invariant, while (3.5) shows that the collision kernel satisfies $b(v_* - v, \omega) = b(v - v_*, \omega)$. Hence

$$\begin{aligned} & \iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F'F'_* - FF_*) \phi b(v - v_*, \omega) \, dv \, dv_* \, d\omega \\ &= \iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F'F'_* - FF_*) \phi_* b(v - v_*, \omega) \, dv \, dv_* \, d\omega \\ &= \frac{1}{2} \iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F'F'_* - FF_*) (\phi + \phi_*) b(v - v_*, \omega) \, dv \, dv_* \, d\omega. \end{aligned}$$

Now in the latter integral, for a.e. fixed $\omega \in \mathbb{S}^2$, apply the change of variables $(v, v_*) \mapsto (v', v'_*)$ defined by (3.3). It is easily seen that this transformation is an involution of $\mathbb{R}^3 \times \mathbb{R}^3$, so that this change of variables maps (v', v'_*) onto (v, v_*) : hence $F'F'_* - FF_*$ is transformed into its opposite $FF_* - F'F'_*$. Formulas (3.3) also show that

$$|v' - v'_*| = v - v_* \quad \text{and} \quad (v' - v'_*) \cdot \omega = -(v - v_*) \cdot \omega$$

so that, by (3.5), one has $b(v' - v'_*, \omega) = b(v - v_*, \omega)$. Finally, this change of variables is an isometry of $\mathbb{R}^3 \times \mathbb{R}^3$ by the second relation of (3.4), and therefore preserves the Lebesgue measure. Eventually, we have proved that

$$\begin{aligned} & \iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F'F'_* - FF_*) (\phi + \phi_*) b(v - v_*, \omega) \, dv \, dv_* \, d\omega \\ &= - \iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F'F'_* - FF_*) (\phi' + \phi'_*) b(v - v_*, \omega) \, dv \, dv_* \, d\omega \\ &= \frac{1}{2} \iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F'F'_* - FF_*) (\phi + \phi_* - \phi' - \phi'_*) b(v - v_*, \omega) \, dv \, dv_* \, d\omega \end{aligned}$$

and this entails the announced formula. \square

Notice that it may not be necessary to assume that F has compact support in Proposition 3.1. For instance, the same result holds for all F and $\phi \in C(\mathbb{R}^3)$ if there exists $m > 0$ such that

$$\begin{aligned} & |\phi(v)| + \int_{\mathbb{S}^2} b(v, \omega) \, d\omega = O(|v|^m) \quad \text{while} \quad F(v) = O(|v|^{-n}) \\ & \text{as } |v| \rightarrow +\infty, \text{ with } n > 2m + 3. \end{aligned} \tag{3.9}$$

An important consequence of this proposition is the following corollary.

1 COROLLARY 3.2. Under the same assumptions as in Proposition 3.1 or (3.9), one has 1

$$2 \int_{\mathbb{R}^3} \mathcal{B}(F, F)(v) \, dv = 0 \quad (\text{conservation of mass}), \quad 3$$

$$4 \int_{\mathbb{R}^3} v_k \mathcal{B}(F, F)(v) \, dv = 0 \quad (\text{conservation of momentum}), \quad 5$$

$$6 \int_{\mathbb{R}^3} \frac{1}{2} |v|^2 \mathcal{B}(F, F)(v) \, dv = 0 \quad (\text{conservation of energy}), \quad 7$$

8 for $k = 1, 2, 3$. 9

10 When applied to a solution of the Boltzmann equation $F \equiv F(t, x, v)$, these five re- 11
12 lations are the net conservation of mass – equivalently, of the total number of particles – 13
14 momentum and energy in each phase-space cylinder $dx \times \mathbb{R}_v^3$, where dx is any infinitesimal 15
16 phase-space element in the space of positions \mathbb{R}_x^3 . 17

18 PROOF OF COROLLARY 3.2. Assuming that $\phi(v)$ is one of the functions 1, v_k for 19
20 $k = 1, 2, 3$, and $\frac{1}{2}|v|^2$, one has 21

$$22 \phi(v) + \phi(v_*) - \phi(v') - \phi(v'_*) = 0 \quad 23$$

24 for each $(v, v_*, \omega) \in \mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2$, because of (3.4). Applying Proposition 3.1 shows the 25
26 five relations stated in Corollary 3.2. \square 27

28 Let $F \equiv F(t, x, v)$ be a solution of the Boltzmann equation; assume that $F(t, x, \cdot)$ is 29
30 continuous with compact support on \mathbb{R}_v^3 a.e. in $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^3$, or satisfies (3.9). Then 31
32 Corollary 3.2 implies that 33

$$34 \begin{aligned} 35 \partial_t \int_{\mathbb{R}^3} F \, dv + \operatorname{div}_x \int_{\mathbb{R}^3} v F \, dv &= 0, \\ 36 \partial_t \int_{\mathbb{R}^3} v F \, dv + \operatorname{div}_x \int_{\mathbb{R}^3} v \otimes v F \, dv &= 0, \\ 37 \partial_t \int_{\mathbb{R}^3} \frac{1}{2} |v|^2 F \, dv + \operatorname{div}_x \int_{\mathbb{R}^3} v \frac{1}{2} |v|^2 F \, dv &= 0. \end{aligned} \quad (3.10) \quad 38$$

39 These equalities are the local conservation laws of mass, momentum and energy in space- 40
41 time divergence form. 42

43 Assume further, for simplicity, that for a.e. $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^3$, 44

$$45 \int_{\mathbb{R}^3} F(t, x, v) \, dv > 0; \quad 46$$

1 define then

$$\begin{aligned}
 2 \quad & \rho(t, x) = \int_{\mathbb{R}^3} F(t, x, v) \, dv \quad (\text{macroscopic density}), \\
 3 \quad & u(t, x) = \frac{1}{\rho(t, x)} \int_{\mathbb{R}^3} v F(t, x, v) \, dv \quad (\text{bulk velocity}), \\
 4 \quad & \theta(t, x) = \frac{1}{\rho(t, x)} \int_{\mathbb{R}^3} \frac{1}{3} |v - u(t, x)|^2 F(t, x, v) \, dv \quad (\text{temperature}).
 \end{aligned}
 \tag{3.11}$$

5 With these definitions, the local conservation laws (3.10) take the form

$$\begin{aligned}
 6 \quad & \partial_t \rho + \operatorname{div}_x(\rho u) = 0, \\
 7 \quad & \partial_t(\rho u) + \operatorname{div}_x(\rho u \otimes u) + \nabla_x(\rho \theta) = -\operatorname{div}_x \int_{\mathbb{R}^3} A(v - u) F \, dv, \\
 8 \quad & \partial_t \left(\rho \left(\frac{1}{2} |u|^2 + \frac{3}{2} \theta \right) \right) + \operatorname{div}_x \left(\rho u \left(\frac{1}{2} |u|^2 + \frac{5}{2} \theta \right) \right) \\
 9 \quad & = -\operatorname{div}_x \int_{\mathbb{R}^3} B(v - u) F \, dv - \operatorname{div}_x \int_{\mathbb{R}^3} A(v - u) \cdot u F \, dv,
 \end{aligned}
 \tag{3.12}$$

10 where

$$A(z) = z \otimes z - \frac{1}{3} |z|^2, \quad B(z) = \frac{1}{2} (|z|^2 - 5)z.$$

11 The left-hand side of the equalities above coincides with that of the compressible Euler system (2.7) with $\gamma = 5/3$ (the adiabatic exponent for point particles, i.e., for particles with 3 degrees of freedom). The right-hand side, on the contrary, depends on the solution of the Boltzmann equation F and is in general not determined by the macroscopic variables ρ , u and θ .

12 However, in some limit, it may be possible to approximate the right-hand side of (3.12) by appropriate functions of ρ , u and θ , thereby arriving at a system in closed form with unknown (ρ, u, θ) .

13 For instance, deriving the compressible Euler system (2.7) as some asymptotic limit of the Boltzmann equation would consist in proving that the right-hand side of the second and third equations in (3.12) vanishes in that limit. Deriving the Navier–Stokes system (2.8) from the Boltzmann equation would consist in finding some (other) asymptotic limit such that

$$\int_{\mathbb{R}^3} A(v - u) F \, dv \simeq -\lambda D(u) \quad \text{and} \quad \int_{\mathbb{R}^3} B(v - u) F \, dv = -\kappa \nabla_x \theta,$$

14 and so on.

15 The problem of finding such closure relations is the key to all the derivations of hydrodynamic models from the Boltzmann equation.

3.2. Boltzmann's H -theorem

We have seen in the last subsection how the symmetries of the Boltzmann collision integral entail the local conservation of mass, momentum and energy.

Another important feature of these symmetries is that they also entail a variant of the second principle of thermodynamics, as we shall now explain.

PROPOSITION 3.3 (Boltzmann's H -theorem). *Assume that the collision kernel b satisfies (3.8), that $F \in C(\mathbb{R}^3)$ is positive and rapidly decaying at infinity, and that, for some $m > 0$, one has*

$$\int_{\mathbb{S}^2} b(v, \omega) d\omega + |\ln F(v)| = O(|v|^m) \quad \text{as } |v| \rightarrow +\infty.$$

Then

$$\begin{aligned} & \int_{\mathbb{R}^3} \mathcal{B}(F, F) \ln F \, dv \\ &= -\frac{1}{4} \iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F' F'_* - F F_*) \ln \left(\frac{F' F'_*}{F F_*} \right) b(v - v_*, \omega) \, dv \, dv_* \, d\omega \leq 0. \end{aligned}$$

Moreover, the following conditions are equivalent

- (i) $\int_{\mathbb{R}^3} \mathcal{B}(F, F) \ln F \, dv = 0$,
- (ii) $\mathcal{B}(F, F)(v) = 0$ for all $v \in \mathbb{R}^3$,
- (iii) F is a Maxwellian distribution, i.e., there exists $\rho, \theta > 0$ and $u \in \mathbb{R}^3$ such that $F = \mathcal{M}_{(\rho, u, \theta)}$, where

$$\mathcal{M}_{(\rho, u, \theta)}(v) := \frac{\rho}{(2\pi\theta)^{3/2}} e^{-|v-u|^2/(2\theta)} \quad \text{for each } v \in \mathbb{R}^3. \quad (3.13)$$

As was already the case of Proposition 3.1, Boltzmann's H -theorem is so essential in deriving hydrodynamic equations from the Boltzmann equation that we give a complete proof of it.

PROOF OF PROPOSITION 3.3. The assumptions on F and b are such that F , b and $\phi = \ln F$ satisfy the assumption (3.9). Applying Proposition 3.1 implies that

$$\begin{aligned} & \int_{\mathbb{R}^3} \mathcal{B}(F, F) \ln F \, dv \\ &= -\frac{1}{4} \iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F' F'_* - F F_*) \ln \left(\frac{F' F'_*}{F F_*} \right) b(v - v_*, \omega) \, dv \, dv_* \, d\omega. \end{aligned} \quad (3.14)$$

1 Since the logarithm is an increasing function, one has

$$2 \quad 3 \quad 4 \quad 5 \quad (f - g) \ln\left(\frac{f}{g}\right) = (f - g)(\ln f - \ln g) \geq 0 \quad \text{for each } f, g > 0,$$

6 so that the expression on the right-hand side of (3.14) is nonpositive.

7 If that expression is equal to zero, the integrand must vanish a.e., meaning that

$$8 \quad 9 \quad F' F'_* = F F_* \quad \text{for a.e. } (v, v_*, \omega) \in \mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2,$$

10 since the collision kernel b is a.e. positive. This accounts for the equivalence between
11 conditions (i) and (ii). That (iii) implies (i) is proved by inspection; for instance, one can
12 observe that, if F is a Maxwellian distribution, $\ln F$ is a linear combination of 1, v_1 , v_2 ,
13 v_3 and $|v|^2$, so that

$$14 \quad 15 \quad 16 \quad \ln F' + \ln F'_* - \ln F - \ln F_* = 0 \quad \text{for all } (v, v_*, \omega) \in \mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2,$$

17 because of the microscopic conservation laws (3.4). Finally, (i) implies (iii), as shown by
18 the next lemma, and this concludes the proof of Boltzmann's H -theorem. \square

19 LEMMA 3.4. *Let $\phi \geq 0$ a.e. be such that $(1 + |v|^2)\phi \in L^1(\mathbb{R}^3)$. If*

$$20 \quad 21 \quad 22 \quad 23 \quad 24 \quad \phi' \phi'_* = \phi \phi_* \quad \text{for a.e. } (v, v_*, \omega) \in \mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2,$$

25 then ϕ is either a.e. 0 or a Maxwellian (i.e., is of the form (3.13)).

26 The following proof is due to Perthame [105]; Boltzmann's original argument can be
27 found in Section 18 of [16].

28 PROOF OF LEMMA 3.4. After translation and multiplication by a constant, one can always
29 assume that

$$30 \quad 31 \quad 32 \quad 33 \quad 34 \quad \int_{\mathbb{R}^3} \phi(v) dv = 1, \quad \int_{\mathbb{R}^3} v \phi(v) dv = 0 \quad (3.15)$$

35 unless $\phi = 0$ a.e. Denoting by $\hat{\phi}$ the Fourier transform of ϕ , our assumptions implies that,
36 for a.e. $\omega \in \mathbb{S}^2$, one has

$$37 \quad 38 \quad 39 \quad 40 \quad 41 \quad 42 \quad 43 \quad 44 \quad 45 \quad \begin{aligned} \hat{\phi}(\xi) \hat{\phi}(\xi_*) &= \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \phi(v') \phi(v'_*) e^{-i\xi \cdot v - i\xi_* \cdot v_*} dv dv_* \\ &= \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \phi(v) \phi(v_*) e^{-i\xi \cdot v' - i\xi_* \cdot v'_*} dv dv_* \\ &= \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \phi(v) \phi(v_*) e^{-i\xi \cdot v - i\xi_* \cdot v_*} e^{i((\xi - \xi_*) \cdot \omega)((v - v_*) \cdot \omega)} dv dv_*. \end{aligned}$$

(Notice that the second equality follows from the same change of variables $(v, v_*) \mapsto (v', v'_*)$ as in the proof of Proposition 3.1.) In fact, this relation holds for all $\omega \in \mathbb{S}^2$ since both sides of the equality above are continuous in ω .

Since the left-hand side of the equality above is independent of ω , one can differentiate in ω to obtain that

$$0 = \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \phi(v) \phi(v_*) e^{-i\xi \cdot v - i\xi_* \cdot v_*} (v - v_*) \cdot \omega_0 \, dv \, dv_*$$

for any $\xi \neq \xi_* \in \mathbb{R}^3$ and $\omega_0 \in \mathbb{S}^2$ such that $\omega_0 \perp (\xi - \xi_*)$. In other words,

$$\omega_0 \perp (\xi - \xi_*) \implies (\nabla_\xi - \nabla_{\xi_*}) \hat{\phi}(\xi) \hat{\phi}(\xi_*) \perp \omega_0.$$

This implies that, for all $\xi \neq \xi_* \in \mathbb{R}^3$, one has

$$(\nabla_\xi - \nabla_{\xi_*}) \hat{\phi}(\xi) \hat{\phi}(\xi_*) \parallel (\xi - \xi_*). \quad (3.16)$$

Applying this with $\xi_* = 0$ leads to

$$\nabla_\xi \hat{\phi}(\xi) \parallel \xi,$$

on account of the normalization condition (3.15). Hence $\hat{\phi}$ is of the form

$$\hat{\phi}(\xi) = \psi(|\xi|^2), \quad \xi \in \mathbb{R}^3.$$

Writing (3.16) with this form of $\hat{\phi}$, one finds that

$$\xi \psi'(|\xi|^2) \psi(|\xi_*|^2) - \xi_* \psi(|\xi|^2) \psi'(|\xi_*|^2) \parallel (\xi - \xi_*).$$

Whenever ξ and ξ_* are not colinear, i.e., for a dense subset of all $\xi, \xi_* \in \mathbb{R}^3$, this implies that

$$\psi'(|\xi|^2) \psi(|\xi_*|^2) = \psi(|\xi|^2) \psi'(|\xi_*|^2).$$

Since $\phi \in L^1((1 + |v|^2) \, dv)$, $\hat{\phi} \in C^2(\mathbb{R})$ and the normalization conditions (3.15) imply that $\hat{\phi}(0) = 1$ and $\hat{\phi}'(0) = 0$; hence $\psi \in C^1(\mathbb{R}^2)$ and the relation above holds for each $(\xi, \xi_*) \in \mathbb{R}^3 \times \mathbb{R}^3$. This relation implies in turn that ψ is of the form

$$\psi(r) = e^{-\theta r/2}.$$

Hence $\hat{\phi}$ is of the form

$$\hat{\phi}(\xi) = e^{-\theta|\xi|^2/2},$$

so that ϕ is of the form $\phi = \mathcal{M}_{(1,0,\theta)}$. □

At this point, it is natural to introduce the notion of *collision invariant*.

1 DEFINITION 3.5. A collision invariant is a measurable function ϕ defined a.e. on \mathbb{R}^3 that
2 satisfies

$$3 \quad \phi(v) + \phi(v_*) - \phi(v') - \phi(v'_*) = 0 \quad \text{a.e. in } (v, v_*, \omega),$$

4 where $v' \equiv v'(v, v_*, \omega)$ and $v'_* \equiv v'_*(v, v_*, \omega)$ are defined by (3.3).
5

6 For instance, in the proof of Boltzmann's H -theorem, Maxwellian densities are charac-
7 terized as the densities whose logarithms are collision invariants.

8 A variant of Lemma 3.4 characterizes collision invariants.

9 PROPOSITION 3.6. A function ϕ is a collision invariant if and only if there exists five
10 constants $a_0, a_1, a_2, a_3, a_4 \in \mathbb{R}$ such that

$$11 \quad \phi(v) = a_0 + a_1 v_1 + a_2 v_2 + a_3 v_3 + a_4 |v|^2 \quad \text{a.e. in } \mathbb{R}^3.$$

12 See Section 3.1 in [28] for a proof of the proposition above.
13
14

15 3.3. H -theorem and a priori estimates

16 We conclude with the main application of Boltzmann's H -theorem, i.e., getting a priori
17 estimates on the solution of the Boltzmann equation. We shall discuss four different cases.
18

19 *Case 1: The periodic box.* Consider the Cauchy problem

$$20 \quad \begin{aligned} 21 \quad \partial_t F + v \cdot \nabla_x F &= \mathcal{B}(F, F), \quad (t, x, v) \in \mathbb{R}_+^* \times \mathbb{T}^3 \times \mathbb{R}^3, \\ 22 \quad F|_{t=0} &= F^{\text{in}}. \end{aligned}$$

23 Let F be a solution of the Boltzmann equation such that, for a.e. $(t, x) \in \mathbb{R}_+ \times \mathbb{T}^3$, $F(t, x, \cdot)$
24 satisfies the assumptions of Proposition 3.3. Then the number density F satisfies the lo-
25 cal entropy inequality (3.26). Integrating this differential inequality on $[0, t] \times \mathbb{T}^3$, one
26 arrives at

$$27 \quad \begin{aligned} 28 \quad & \iint_{\mathbb{T}^3 \times \mathbb{R}^3} F \ln F(t, x, v) \, dx \, dv \\ 29 \quad & + \frac{1}{4} \int_0^t \int_{\mathbb{T}^3} \iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F' F'_* - F F_*) \ln \left(\frac{F' F'_*}{F F_*} \right) b \, dv \, dv_* \, d\omega \, dx \, ds \\ 30 \quad & = \iint_{\mathbb{T}^3 \times \mathbb{R}^3} F^{\text{in}} \ln F^{\text{in}}(x, v) \, dx \, dv \end{aligned} \quad (3.17)$$

31 for each $t \geq 0$.

32 The following definition explains the name " H -theorem".
33
34
35
36
37
38
39
40
41
42
43
44
45

1 DEFINITION 3.7. Let $F \geq 0$ a.e. be an element of $L^1(\mathbb{T}^3 \times \mathbb{R}^3)$ such that

$$2 \int \int_{\mathbb{T}^3 \times \mathbb{R}^3} |F \ln F(x, v)| \, dx \, dv < +\infty. \quad 3$$

4 One denotes by $H(F)$ the quantity

$$5 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \quad H(F) = \int \int_{\mathbb{T}^3 \times \mathbb{R}^3} F \ln F(x, v) \, dx \, dv.$$

11 Whenever there is no risk of ambiguity, we use the notation $H(t)$ to designate
12 $H(F(t, \cdot, \cdot))$, when F is a solution of the Boltzmann equation. Equality (3.17) implies
13 that $H(F)$ is a nonincreasing function of time; it was this property that Boltzmann called
14 “the H -theorem”. Moreover, $H(F)$ is stationary only if F is a Maxwellian (see Sec-
15 tion 3.4.2). Hence, from the physical viewpoint, it is natural to think of $H(F(t, \cdot, \cdot))$ as
16 *minus the entropy* of the system of particles distributed under $F(t, \cdot, \cdot)$.

17 In order to obtain a bound on the entropy production, it is convenient to introduce another
18 (closely related) concept of entropy.

19
20 DEFINITION 3.8. Let $F \geq 0$ a.e. and $G > 0$ be two measurable functions on $\mathbb{T}^3 \times \mathbb{R}^3$; the
21 relative entropy of F with respect to G is

$$22 \quad 23 \quad 24 \quad 25 \quad 26 \quad 27 \quad 28 \quad 29 \quad 30 \quad 31 \quad 32 \quad 33 \quad 34 \quad 35 \quad H(F|G) = \int \int_{\mathbb{T}^3 \times \mathbb{R}^3} \left(F \ln \left(\frac{F}{G} \right) - F + G \right) \, dx \, dv.$$

Notice that the integrand in the definition of $H(F|G)$ is an a.e. nonnegative measurable
function, so that the relative entropy $H(F|G)$ is well defined as an element of $[0, +\infty]$.

Let $\rho, \theta > 0$ and $u \in \mathbb{R}^3$, then

$$36 \quad 37 \quad 38 \quad 39 \quad 40 \quad 41 \quad 42 \quad 43 \quad 44 \quad 45 \quad H(F|\mathcal{M}_{(\rho, u, \theta)}) = H(t) + \int \int_{\mathbb{T}^3 \times \mathbb{R}^3} \frac{|v - u|^2}{2\theta} F \, dx \, dv \\ - \left(1 + \ln \left(\frac{\rho}{(2\pi\theta)^{3/2}} \right) \right) \int \int_{\mathbb{T}^3 \times \mathbb{R}^3} F \, dx \, dv + \rho. \quad (3.18)$$

Hence, if $F \in L^1(\mathbb{T}^3 \times \mathbb{R}^3; (1 + |v|^2) \, dv \, dx)$ and if $H(0)$ is finite, then $H(t)$ is finite for
each $t \geq 0$, and

$$- \left(\left| \ln \left(\frac{\rho}{(2\pi\theta)^{3/2}} \right) \right| + \frac{|u|^2 + 1}{\theta} \right) \int \int_{\mathbb{T}^3 \times \mathbb{R}^3} (1 + |v|^2) F \, dv \, dx - \rho \\ \leq H(t) \leq H(0).$$

On the other hand, F also satisfies the local conservation of mass, momentum, and en-
ergy (3.10), so that, integrating these local conservation laws on $[0, t] \times \mathbb{T}^3$, one arrives at

the *global* variant of these conservation laws

$$\begin{aligned}
\iint_{\mathbb{T}^3 \times \mathbb{R}^3} F(t, x, v) \, dv \, dx &= \iint_{\mathbb{T}^3 \times \mathbb{R}^3} F(0, x, v) \, dv \, dx, \\
\iint_{\mathbb{T}^3 \times \mathbb{R}^3} v F(t, x, v) \, dv \, dx &= \iint_{\mathbb{T}^3 \times \mathbb{R}^3} v F(0, x, v) \, dv \, dx, \\
\iint_{\mathbb{T}^3 \times \mathbb{R}^3} \frac{1}{2} |v|^2 F(t, x, v) \, dv \, dx &= \iint_{\mathbb{T}^3 \times \mathbb{R}^3} \frac{1}{2} |v|^2 F(0, x, v) \, dv \, dx.
\end{aligned} \tag{3.19}$$

Since

$$\begin{aligned}
&\iint_{\mathbb{T}^3 \times \mathbb{R}^3} \frac{|v - u|^2}{2\theta} F \, dx \, dv \\
&= \frac{1}{2\theta} \iint_{\mathbb{T}^3 \times \mathbb{R}^3} (|v|^2 + |u|^2) F \, dx \, dv - \frac{1}{\theta} \iint_{\mathbb{T}^3 \times \mathbb{R}^3} u \cdot v F \, dx \, dv,
\end{aligned}$$

one has

$$H(F(t)|\mathcal{M}_{(\rho, u, \theta)}) = H(t) + \text{globally conserved quantities}$$

so that

$$H(F(t)|\mathcal{M}_{(\rho, u, \theta)}) - H(F(0)|\mathcal{M}_{(\rho, u, \theta)}) = H(t) - H(0).$$

Hence, the global entropy relation (3.17) is recast in terms of the relative entropy as

$$\begin{aligned}
&\frac{1}{4} \int_0^t \int_{\mathbb{T}^3} \iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F' F'_* - F F_*') \ln \left(\frac{F' F'_*}{F F_*'} \right) b \, dv \, dv_* \, d\omega \, dx \, ds \\
&= H(F(0)|\mathcal{M}_{(\rho, u, \theta)}) - H(F(t)|\mathcal{M}_{(\rho, u, \theta)})
\end{aligned} \tag{3.20}$$

for each $\rho, \theta > 0$ and each $u \in \mathbb{R}^3$. This implies, in particular,

- the relative entropy bound

$$0 \leq H(F(t)|\mathcal{M}_{(\rho, u, \theta)}) \leq H(F(0)|\mathcal{M}_{(\rho, u, \theta)}), \quad t \geq 0;$$

- the following entropy control

$$\begin{aligned}
&-\left(\left| \ln \left(\frac{\rho}{(2\pi\theta)^{3/2}} \right) \right| + \frac{|u|^2 + 1}{\theta} \right) \iint_{\mathbb{T}^3 \times \mathbb{R}^3} (1 + |v|^2) F^{\text{in}} \, dv \, dx - \rho \\
&\leq H(F)(t) \leq H(F)(0);
\end{aligned}$$

- the entropy production estimate

$$\begin{aligned} & \frac{1}{4} \int_0^{+\infty} \int_{\mathbb{T}^3} \iint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F' F'_* - F F_*') \ln \left(\frac{F' F'_*}{F F_*'} \right) b \, dv \, dv_* \, d\omega \, dx \, ds \\ & \leq H(F(0) | \mathcal{M}_{(\rho, u, \theta)}). \end{aligned}$$

Case 2: A bounded domain with specular reflection on the boundary. The periodic box is a somewhat academic choice of a spatial domain for studying the Boltzmann equation. The next case that we consider now is very similar but more realistic. Let Ω be a smooth, bounded domain of \mathbb{R}^3 . Starting from a given number density F^{in} at time $t = 0$, we consider the initial boundary value problem

$$\begin{aligned} \partial_t F + v \cdot \nabla_x F &= \mathcal{B}(F, F), & (x, v) \in \Omega \times \mathbb{R}^3, \\ F(t, x, v) &= F(t, x, \mathcal{R}_x v), & (x, v) \in \partial\Omega \times \mathbb{R}^3, \\ F|_{t=0} &= F^{\text{in}}, \end{aligned}$$

where \mathcal{R}_x designates the specular reflection defined by the outward unit normal n_x at $x \in \partial\Omega$

$$\mathcal{R}_x v = v - 2(v \cdot n_x) n_x.$$

Assume that the initial boundary value problem above has a solution F satisfying the assumptions of Proposition 3.3. One multiplies the Boltzmann equation above by $\ln F + 1$ and integrates first in v to obtain the identity (3.26), and then integrates in x , which leads to

$$\begin{aligned} & \frac{d}{dt} \iint_{\Omega \times \mathbb{R}^3} F \ln F \, dx \, dv + \iint_{\partial\Omega \times \mathbb{R}^3} F \ln F v \cdot n_x \, d\sigma(x) \, dv \\ & = -\frac{1}{4} \iint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F' F'_* - F F_*') \ln \left(\frac{F' F'_*}{F F_*'} \right) b(v - v_*, \omega) \, dv \, dv_* \, d\omega. \end{aligned}$$

Changing the v variable in the boundary term by $v \mapsto w = \mathcal{R}_x v$, one sees that $w \cdot n_x = -v \cdot n_x$, while the specular reflection condition satisfied by F on $\partial\Omega$ implies that $F(t, x, v) = F(t, x, w)$; besides this change of variables preserves the Lebesgue measure dv since \mathcal{R}_x is an isometry. Hence

$$\iint_{\partial\Omega \times \mathbb{R}^3} F \ln F v \cdot n_x \, d\sigma(x) \, dv = -\iint_{\partial\Omega \times \mathbb{R}^3} F \ln F w \cdot n_x \, d\sigma(x) \, dw = 0$$

and therefore

$$\begin{aligned} & \frac{d}{dt} H(F)(t) \\ & = -\frac{1}{4} \iint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F' F'_* - F F_*') \ln \left(\frac{F' F'_*}{F F_*'} \right) b(v - v_*, \omega) \, dv \, dv_* \, d\omega. \end{aligned} \tag{3.21}$$

Proceeding similarly with the local conservation laws (3.10) shows that

$$\frac{d}{dt} \iint_{\Omega \times \mathbb{R}^3} F \, dx \, dv = \frac{d}{dt} \iint_{\Omega \times \mathbb{R}^3} \frac{1}{2} |v|^2 F \, dx \, dv = 0. \quad (3.22)$$

At this point, we apply the formula (3.18) in the case where $u = 0$ and deduce from (3.22) that

$$H(F | \mathcal{M}_{(\rho, 0, \theta)}) = H(F) + \text{globally conserved quantities.}$$

Notice that, at variance with the case of the periodic box, the total momentum is not conserved, so that the formula above only holds with centered Maxwellians (i.e., Maxwellians with zero bulk velocity).

Therefore, as in the case of the periodic box, one has, for each $\rho, \theta > 0$,

$$\begin{aligned} & \frac{1}{4} \int_0^t \iiint_{\Omega \times \mathbb{R}^3 \times \mathbb{S}^2} (F' F'_* - F F_*') \ln \left(\frac{F' F'_*}{F F_*'} \right) b \, dv \, dv_* \, d\omega \, dx \, ds \\ & = H(F(0) | \mathcal{M}_{(\rho, 0, \theta)}) - H(F(t) | \mathcal{M}_{(\rho, 0, \theta)}) \end{aligned} \quad (3.23)$$

for each $t \geq 0$. Again we obtain

- the relative entropy bound

$$0 \leq H(F(t) | \mathcal{M}_{(\rho, u, \theta)}) \leq H(F(0) | \mathcal{M}_{(\rho, u, \theta)}), \quad t \geq 0;$$

- the following entropy control

$$\begin{aligned} & - \left(\left| \ln \left(\frac{\rho}{(2\pi\theta)^{3/2}} \right) \right| + \frac{1}{\theta} \right) \iint_{\mathbb{T}^3 \times \mathbb{R}^3} (1 + |v|^2) F^{\text{in}} \, dv \, dx - \rho \\ & \leq H(F)(t) \leq H(F)(0); \end{aligned}$$

- the entropy production estimate

$$\begin{aligned} & \frac{1}{4} \int_0^{+\infty} \iiint_{\Omega \times \mathbb{R}^3 \times \mathbb{S}^2} (F' F'_* - F F_*') \ln \left(\frac{F' F'_*}{F F_*'} \right) b \, dv \, dv_* \, d\omega \, dx \, ds \\ & \leq H(F(0) | \mathcal{M}_{(\rho, u, \theta)}). \end{aligned}$$

Case 3: The Euclidean space with Maxwellian equilibrium at infinity. Next we study cases where the spatial domain is unbounded. The simplest of such cases is that of a cloud of gas which is in Maxwellian equilibrium at infinity. Therefore, we consider the Cauchy problem

$$\partial_t F + v \cdot \nabla_x F = \mathcal{B}(F, F), \quad (t, x, v) \in \mathbb{R}_+^* \times \mathbb{R}^3 \times \mathbb{R}^3,$$

$$F(t, x, v) \rightarrow \mathcal{M}_{(\rho, u, \theta)}, \quad |x| \rightarrow +\infty,$$

$$F|_{t=0} = F^{\text{in}}.$$

We shall assume that F converges to the Maxwellian state $\mathcal{M}_{(\rho,u,\theta)}$ rapidly enough so that the relative entropy

$$\begin{aligned} H(F(t)|\mathcal{M}_{(\rho,u,\theta)}) \\ = \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \left(F \ln \left(\frac{F}{\mathcal{M}_{(\rho,u,\theta)}} \right) - F + \mathcal{M}_{(\rho,u,\theta)} \right) dx dv < +\infty \end{aligned}$$

for each $t \geq 0$. We claim that the same entropy relation as in the case of the three-torus also holds in the present situation

$$\begin{aligned} \frac{1}{4} \int_0^t \iint_{\mathbb{R}^3} \iint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F' F'_* - F F_*) \ln \left(\frac{F' F'_*}{F F_*} \right) b dv dv_* d\omega dx ds \\ = H(F(0)|\mathcal{M}_{(\rho,u,\theta)}) - H(F(t)|\mathcal{M}_{(\rho,u,\theta)}) \end{aligned} \quad (3.24)$$

for each $t \geq 0$. However, this equality is not obtained in the same way, since neither the globally conserved quantities nor the H -function itself are well-defined objects in this case (these quantities involve divergent integrals because of the Maxwellian condition at infinity).

Observe instead that, by the same argument as in the case of the three-torus, one has

$$\begin{aligned} \int_{\mathbb{R}^3} \left(F \ln \left(\frac{F}{\mathcal{M}_{(\rho,u,\theta)}} \right) - F + \mathcal{M}_{(\rho,u,\theta)} \right) dv \\ = \int_{\mathbb{R}^3} F \ln F dv + \frac{1}{2\theta} \int_{\mathbb{R}^3} (|v|^2 + |u|^2) F dv \\ - \frac{1}{\theta} \int_{\mathbb{R}^3} u \cdot v F dv + \left(1 + \ln \left(\frac{\rho}{(2\pi\theta)^{3/2}} \right) \right) \int_{\mathbb{R}^3} F dv + \rho \end{aligned}$$

while

$$\begin{aligned} \int_{\mathbb{R}^3} v \left(F \ln \left(\frac{F}{\mathcal{M}_{(\rho,u,\theta)}} \right) - F + \mathcal{M}_{(\rho,u,\theta)} \right) dv \\ = \int_{\mathbb{R}^3} v F \ln F dv + \frac{1}{2\theta} \int_{\mathbb{R}^3} v (|v|^2 + |u|^2) F dv \\ - \frac{1}{\theta} \int_{\mathbb{R}^3} v u \cdot v F dv + \left(1 + \ln \left(\frac{\rho}{(2\pi\theta)^{3/2}} \right) \right) \int_{\mathbb{R}^3} v F dv. \end{aligned}$$

In other words,

$$\begin{aligned} \int_{\mathbb{R}^3} \left(F \ln \left(\frac{F}{\mathcal{M}_{(\rho,u,\theta)}} \right) - F + \mathcal{M}_{(\rho,u,\theta)} \right) dv \\ = \int_{\mathbb{R}^3} F \ln F dv + \text{locally conserved quantity} \end{aligned}$$

1 while

$$\begin{aligned} & \int_{\mathbb{R}^3} v \left(F \ln \left(\frac{F}{\mathcal{M}(\rho, u, \theta)} \right) - F + \mathcal{M}(\rho, u, \theta) \right) dv \\ &= \int_{\mathbb{R}^3} v F \ln F dv + \text{flux of that locally conserved quantity} \end{aligned}$$

9 so that

$$\begin{aligned} & \partial_t \int_{\mathbb{R}^3} \left(F \ln \left(\frac{F}{\mathcal{M}(\rho, u, \theta)} \right) - F + \mathcal{M}(\rho, u, \theta) \right) dv \\ &+ \operatorname{div}_x \int_{\mathbb{R}^3} v \left(F \ln \left(\frac{F}{\mathcal{M}(\rho, u, \theta)} \right) - F + \mathcal{M}(\rho, u, \theta) \right) dv \\ &= \partial_t \int_{\mathbb{R}^3} F \ln F dv + \operatorname{div}_x \int_{\mathbb{R}^3} v F \ln F dv \end{aligned}$$

19 so that

$$\begin{aligned} & \partial_t \int_{\mathbb{R}^3} \left(F \ln \left(\frac{F}{\mathcal{M}(\rho, u, \theta)} \right) - F + \mathcal{M}(\rho, u, \theta) \right) dv \\ &+ \operatorname{div}_x \int_{\mathbb{R}^3} v \left(F \ln \left(\frac{F}{\mathcal{M}(\rho, u, \theta)} \right) - F + \mathcal{M}(\rho, u, \theta) \right) dv \\ &= -\frac{1}{4} \iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F' F'_* - F F_*) \ln \left(\frac{F' F'_*}{F F_*} \right) b dv dv_* d\omega. \end{aligned}$$

30 Integrating further on $[0, t] \times \mathbb{R}^3$, one arrives at (3.24).

31 To summarize, we deduce from (3.24) that

- 32 • the relative entropy bound

$$34 \quad 0 \leq H(F(t) | \mathcal{M}(\rho, u, \theta)) \leq H(F^{\text{in}} | \mathcal{M}(\rho, u, \theta)) \quad \text{for each } t \geq 0;$$

- 36 • the entropy production estimate

$$\begin{aligned} & \frac{1}{4} \int_0^{+\infty} \iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F' F'_* - F F_*) \ln \left(\frac{F' F'_*}{F F_*} \right) b dv dv_* d\omega dx ds \\ & \leq H(F^{\text{in}} | \mathcal{M}(\rho, u, \theta)). \end{aligned}$$

44 *Case 4: The Euclidean space with vacuum at infinity.* Finally, we consider the case of a
45 cloud of gas expanding in the vacuum. As we shall see, this case is slightly different from

1 the previous ones. Consider the Cauchy problem

$$\begin{aligned}
 2 \quad & \partial_t F + v \cdot \nabla_x F = \mathcal{B}(F, F), \quad (t, x, v) \in \mathbb{R}_+^* \times \mathbb{R}^3 \times \mathbb{R}^3, \\
 3 \quad & F(t, x, v) \rightarrow 0, \quad |x|, |v| \rightarrow +\infty, \\
 4 \quad & F|_{t=0} = F^{\text{in}}.
 \end{aligned}$$

5 We shall assume that F vanishes rapidly enough at infinity so that the relative entropy

$$6 \quad H(F(t)|\mathcal{G}) < +\infty \quad \text{for each } t \geq 0,$$

7 where \mathcal{G} is the centered reduced Gaussian

$$8 \quad \mathcal{G}(x, v) = \frac{1}{(2\pi)^3} e^{-(|x|^2 + |v|^2)/2}.$$

9 Assume that

$$10 \quad \iint_{\mathbb{R}^3 \times \mathbb{R}^3} F^{\text{in}} (|\ln F^{\text{in}}| + |x|^2 + |v|^2 + 1) dx dv < +\infty.$$

11 We claim that, for each ≥ 0 ,

$$12 \quad \iint_{\mathbb{R}^3 \times \mathbb{R}^3} |x - tv|^2 F(t, x, v) dx dv = \iint_{\mathbb{R}^3 \times \mathbb{R}^3} |x|^2 F(0, x, v) dx dv. \quad (3.25)$$

13 Indeed

$$\begin{aligned}
 14 \quad & \frac{d}{dt} \iint_{\mathbb{R}^3 \times \mathbb{R}^3} |x - tv|^2 F(t, x, v) dx dv \\
 15 \quad & = \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \partial_t (|x - tv|^2 F(t, x, v)) dx dv \\
 16 \quad & = \iint_{\mathbb{R}^3 \times \mathbb{R}^3} (\partial_t + v \cdot \nabla_x) (|x - tv|^2 F(t, x, v)) dx dv \\
 17 \quad & = \iint_{\mathbb{R}^3 \times \mathbb{R}^3} |x - tv|^2 (\partial_t + v \cdot \nabla_x) F(t, x, v) dx dv \\
 18 \quad & = \int_{\mathbb{R}^3} \left(\int_{\mathbb{R}^3} |x - tv|^2 \mathcal{B}(F, F)(t, x, v) dv \right) dx = 0.
 \end{aligned}$$

19 Observe that

$$\begin{aligned}
 20 \quad & H(F|\mathcal{G}) = H(F) + \frac{1}{2} \iint_{\mathbb{R}^3 \times \mathbb{R}^3} (|x|^2 + |v|^2) F dx dv \\
 21 \quad & + (3 \ln(2\pi) - 1) \iint_{\mathbb{R}^3 \times \mathbb{R}^3} F dx dv + 1.
 \end{aligned}$$

1 Because of (3.25), one has

$$\begin{aligned}
 & \iint_{\mathbb{R}^3 \times \mathbb{R}^3} |x|^2 F(t) \, dx \, dv \\
 & \leq 2 \iint_{\mathbb{R}^3 \times \mathbb{R}^3} |x - tv|^2 F(t) \, dx \, dv + 2t^2 \iint_{\mathbb{R}^3 \times \mathbb{R}^3} |v|^2 F(t) \, dx \, dv \\
 & = 2 \iint_{\mathbb{R}^3 \times \mathbb{R}^3} |x|^2 F(0) \, dx \, dv + 2t^2 \iint_{\mathbb{R}^3 \times \mathbb{R}^3} |v|^2 F(0) \, dx \, dv
 \end{aligned}$$

11 so that

$$\begin{aligned}
 & - \iint_{\mathbb{R}^3 \times \mathbb{R}^3} |x|^2 F^{\text{in}} \, dx \, dv - \left(\frac{1}{2} + t^2 \right) \iint_{\mathbb{R}^3 \times \mathbb{R}^3} |v|^2 F^{\text{in}} \, dx \, dv \\
 & - (3 \ln(2\pi) - 1) \iint_{\mathbb{R}^3 \times \mathbb{R}^3} F^{\text{in}} \, dx \, dv - 1 \leq H(t) \leq H(0).
 \end{aligned}$$

19 Integrating on $[0, t] \times \mathbb{R}^3$ the local entropy equality (3.26), one arrives at the equality

$$\begin{aligned}
 & H(0) - H(t) \\
 & = \frac{1}{4} \int_0^t \iint_{\mathbb{R}^3} \iint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F' F'_* - F F_*') \ln \left(\frac{F' F'_*}{F F_*'} \right) b \, dv \, dv_* \, d\omega \, dx \, ds \\
 & \leq C(1 + t^2) \iint_{\mathbb{R}^3 \times \mathbb{R}^3} (1 + |x|^2 + |v|^2 + |\ln F^{\text{in}}|) F^{\text{in}} \, dv \, dx.
 \end{aligned}$$

28 As we shall see below, Cases 1–3 are the most useful in the context of hydrodynamic limits. Case 4 is also interesting, although not for hydrodynamic limits: it provides one of the important estimates in the construction of global weak solutions to the Boltzmann equation by R. DiPerna and P.-L. Lions (see further).

32 Another case, which we did not discuss in spite of its obvious interest for applications, is that of a spatial domain that is the complement in \mathbb{R}^3 of a regular compact set, assuming specular reflection of the particles at the boundary of the domain. This case is handled by a straightforward adaptation of the arguments in Cases 2 and 3.

36 Let us now briefly discuss some of the main consequences of Boltzmann's H -theorem.

40 3.4. Further remarks on the H -theorem

42 **3.4.1. H -theorem and the second principle of thermodynamics.** To begin with, let F be a solution of the Boltzmann equation such that, for a.e. $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^3$, $F(t, x, \cdot)$ satisfies the assumptions of Proposition 3.3. Multiplying both sides of the Boltzmann equation by $\ln F + 1$ and applying Proposition 3.3 and Corollary 3.2 with $\phi \equiv 1$, one arrives at the

1 identity

$$\begin{aligned}
 & \partial_t \int_{\mathbb{R}^3} F \ln F \, dv + \operatorname{div}_x \int_{\mathbb{R}^3} v F \ln F \, dv \\
 &= -\frac{1}{4} \iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F' F'_* - F F_*) \ln \left(\frac{F' F'_*}{F F_*} \right) b(v - v_*, \omega) \, dv \, dv_* \, d\omega \\
 &\leq 0.
 \end{aligned} \tag{3.26}$$

2
3
4
5
6
7
8
9
10 It is interesting to compare the equality (3.26) with the second principle of thermodynamics
11 applied to any portion of a fluid in a smooth domain Ω . Denoting by n_x the outward unit
12 normal field on $\partial\Omega$, and by s the entropy per unit of mass in the fluid, one has

$$\frac{d}{dt} \int_{\Omega} \rho s(t, x) \, dt \geq - \int_{\partial\Omega} \rho s u(t, x) \cdot n_x \, d\sigma(x) - \int_{\partial\Omega} \frac{q(t, x)}{\theta(t, x)} \cdot n_x \, d\sigma(x),$$

13
14
15
16
17 where ρ is the density of the fluid, u the velocity field, θ the temperature, q the heat flux
18 and $d\sigma(x)$ the surface element on $\partial\Omega$. The infinitesimal version of this inequality is

$$\partial_t(\rho s) + \operatorname{div}_x \left(\rho s u + \frac{q}{\theta} \right) \geq 0, \tag{3.27}$$

19
20
21
22 which is obviously analogous to (3.26). In particular,

$$\text{the quantity } - \int_{\mathbb{R}^3} F \ln F \, dv \text{ is analogous to } \rho s$$

23
24
25
26
27 and

$$\text{the quantity } - \int_{\mathbb{R}^3} v F \ln F \, dv \text{ is analogous to } \rho s u,$$

28
29
30
31
32 while the quantity

$$\frac{1}{4} \iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F' F'_* - F F_*) \ln \left(\frac{F' F'_*}{F F_*} \right) b(v - v_*, \omega) \, dv \, dv_* \, d\omega$$

33
34
35
36
37 is the local entropy production. Notice that fluid dynamics does not in general provide any
38 expression of the entropy production in terms of ρ , u , θ , s and q . On the contrary, in the
39 kinetic theory of gases, the entropy production is given in terms of the number density by
40 the integral above.

41
42 **3.4.2. Relaxation towards equilibrium.** One application of the second principle of ther-
43 modynamics is the relaxation towards equilibrium for closed systems. Assume that a gas
44 described by the Boltzmann equation is enclosed in some container Ω that is a smooth,
45 bounded domain of \mathbb{R}^3 . At the microscopic level, we assume that the gas molecules are

1 reflected on the surface of the container without exchanging heat. One model for this is 1
 2 the ideal situation where each molecule impinging on the boundary of the container is 2
 3 specularly reflected, this being Case 2 of Section 3.3. 3

4 Starting from a given number density F^{in} at time $t = 0$, we consider the initial boundary 4
 5 value problem 5

$$\begin{aligned} 6 & \partial_t F + v \cdot \nabla_x F = \mathcal{B}(F, F), \quad (x, v) \in \Omega \times \mathbb{R}^3, \\ 7 & F(t, x, v) = F(t, x, \mathcal{R}_x v), \quad (x, v) \in \partial\Omega \times \mathbb{R}^3, \\ 8 & F|_{t=0} = F^{\text{in}}, \end{aligned}$$

9 where \mathcal{R}_x designates the specular reflection defined by the outward unit normal n_x at 9
 10 $x \in \partial\Omega$ 10
 11 11

$$12 \quad \mathcal{R}_x v = v - 2v \cdot n_x n_x. \quad 12$$

13 Now pick any sequence $t_n \rightarrow +\infty$ such that 13
 14 14

$$15 \quad F_n(t, x, v) := F(t + t_n, x, v) \rightarrow E(t, x, v) \quad 15$$

16 in a weak topology that is compatible with the conservation laws (3.22). Then, by weak 16
 17 convergence and convexity, the bound on entropy production obtained in Case 2 of 17
 18 Section 3.3 implies that $E(t, x, v)$ is a local Maxwellian – meaning that the function 18
 19 $v \mapsto E(t, x, v)$ is a.e. a Maxwellian with parameters ρ, u, θ that are functions of t, x – 19
 20 which satisfies 20

$$\begin{aligned} 21 & (\partial_t E + v \cdot \nabla_x E)(t, x, v) = 0, \quad (x, v) \in \Omega \times \mathbb{R}^3, \\ 22 & E(t, x, v) = E(t, x, \mathcal{R}_x v), \quad (x, v) \in \partial\Omega \times \mathbb{R}^3. \end{aligned} \quad 21$$

23 Whenever Ω is not rotationally invariant with respect to some axis of symmetry, the only 23
 24 local Maxwellians that solve the system of equations (3.29) are the global Maxwellians of 24
 25 the form 25

$$26 \quad E(t, x, v) = \mathcal{M}_{(\rho, 0, \theta)}(v) \quad \text{for some constant } \rho, \theta > 0. \quad 26$$

27 Since we assumed that the conservation laws (3.22) are compatible with the topology in 27
 28 which the long time limit holds, the constants ρ and θ are given by 28
 29 29

$$30 \quad \iint_{\Omega \times \mathbb{R}^3} F^{\text{in}} dx dv = \rho |\Omega|, \quad \iint_{\Omega \times \mathbb{R}^3} \frac{1}{2} F^{\text{in}} dx dv = \frac{3}{2} \rho \theta |\Omega|. \quad 30$$

31 However, if Ω is rotationally invariant around some axis of symmetry, the system (3.29) 31
 32 has other solutions than the global Maxwellians (3.30), namely all functions of the form 32
 33 33

$$34 \quad E(t, x, v) = \mathcal{M}_{(\rho, 0, \theta)}(v) e^{\lambda(k \times (x - x_0)) \cdot v}, \quad 34$$

1 where $\lambda \neq 0$ is a constant, while the axis of rotational symmetry for Ω is the line of 1
2 direction $k \in \mathbb{S}^2$ passing through x_0 . 2

3 In fact, Lions proved in Section V of [84] that the convergence (3.28) is locally uniform 3
4 in t with values in the strong topology of $L^1(\Omega \times \mathbb{R}^3)$. 4

5 Let us mention that, in spite of his apparent simplicity, the problem of relaxation to- 5
6 ward equilibrium for the Boltzmann equation is still open, in spite of recent progress by 6
7 Desvillettes and Villani [35]. The main issue is the lack of a tightness estimate in the 7
8 v variable for $|v|^2 F(t, x, v)$ as $t \rightarrow +\infty$ that would apply to all initial data of finite mass, 8
9 energy and entropy. Short of such an estimate, the part of the argument above identify- 9
10 ing the temperature in terms of the initial data fails. The only cases where such estimates 10
11 have been obtained correspond to initial data that are already close enough to some global 11
12 Maxwellian, or that are independent of the space variable (i.e., the space homogeneous 12
13 case). 13

14 However incomplete, this discussion shows the importance of Boltzmann's H -theorem 14
15 whenever one seeks to estimate how close to the class of local Maxwellians a given solution 15
16 of the Boltzmann equation may be. This particular point is of paramount importance for 16
17 hydrodynamic limits. 17

18

19

20

20 3.5. The collision kernel 20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

22 So far, our discussion of the Boltzmann equation – in fact, of the Boltzmann collision 22
23 integral – did not use much of the properties of the collision kernel b . Indeed, we only 23
24 took advantage of the symmetries of b in (3.5) and some additional bounds such as (3.8) 24
25 or (3.9). 25

26 However, the derivation of hydrodynamic limits requires further properties of the colli- 26
27 sion integral, for which a more extensive discussion of the collision kernel becomes neces- 27
28 sary. 28

29 First we recall some elementary facts concerning the two-body problem. Consider two 29
30 points of unit mass subject to a repulsive interaction potential $U \equiv U(r)$, where r is the 30
31 distance between these two points. In other words, assume that U satisfies the properties 31

$$32 \quad U \in C^\infty(\mathbb{R}_+^*) \text{ is decreasing,} \quad \lim_{r \rightarrow 0^+} U(r) = +\infty, \quad \lim_{r \rightarrow +\infty} U(r) = 0^+. \quad 33$$

35 It is well known that both points stay in the same plane for all times. Pick a Galilean 35
36 frame where one of the points is at rest; then, the trajectory of the moving point is easily 36
37 expressed in polar coordinates (r, θ) with the fixed point as origin. Choosing the origin of 37
38 polar angles to be the line asymptotic to the trajectory of the moving particle in the past, 38
39 the trajectory is determined as follows. Let v be the speed of the moving particle at infinity, 39
40 and let h be the impact parameter defined by 40

$$41 \quad vh = r^2(t)\dot{\theta}(t), \quad t \in \mathbb{R}, \quad 42$$

44 this quantity being a well-known first integral of the motion (see Figure 2). In other words, 44
45 h is the distance between the line asymptotic to the trajectory of the moving particle in the 45




Fig. 2. Deflection of a particle subject to a radial repulsion potential from a particle at rest.

past and the parallel line going through the particle at rest (see Figure 2). Let $z_* > 0$ be the unique solution to

$$1 - z_*^2 - \frac{4}{v^2} U\left(\frac{h}{z_*}\right) = 0,$$

and set

$$r_* = \frac{h}{z_*} \quad \text{and} \quad \theta_* = \int_0^{z_*} \frac{dz}{\sqrt{1 - z^2 - 4/v^2 U(h/z)}}. \tag{3.31}$$

The point of polar coordinates (r_*, θ_*) is the apse of the trajectory, i.e., the closest to the particle at rest. Then the trajectory of the moving particles is given in polar coordinates by the equation

$$\theta = \begin{cases} \int_0^{h/r} \frac{dz}{\sqrt{1 - z^2 - 4/v^2 U(h/z)}} & \text{for } \theta \in (0, \theta_*] \text{ and } r > r_*, \\ 2\theta_* - \int_0^{h/r} \frac{dz}{\sqrt{1 - z^2 - 4/v^2 U(h/z)}} & \text{for } \theta \in [\theta_*, 2\theta_*) \text{ and } r > r_*. \end{cases}$$

Notice in particular that the moving particle is deflected of an angle $2\theta_*$.

Next we recall the notion of *scattering cross-section*. Pick an arbitrary relative speed v at infinity, and consider the deflection angle $\chi_* = \pi - 2\theta_*$ as a function of the impact parameter h . It is easily seen that

$$\chi_* \text{ is decreasing, } \quad \lim_{h \rightarrow 0^+} \chi_*(h) = \pi^- \quad \text{and} \quad \lim_{h \rightarrow +\infty} \chi_*(h) = 0^+.$$

Because the two-body problem is invariant by any rotation around the line D passing through the particle at rest that is parallel to the asymptote in the past to the trajectory




Fig. 3. The scattering cross-section corresponding to the relative speed v and in the direction χ corresponding to the impact parameter h .

of the moving particle, consider the map

$$\begin{aligned} \mathbb{R}^2 \setminus \{0\} &\simeq \mathbb{R}_+^* \times [0, 2\pi) \rightarrow (0, \pi) \times [0, 2\pi) \simeq \mathbb{S}^2 \setminus \{N, S\}, \\ (h, \phi) &\mapsto (\chi_*(h), \phi), \end{aligned} \tag{3.32}$$

where the first identification is through polar coordinates in the plane orthogonal to D with origin $h = 0$ (the intersection of D with that plane), while the second identification is through spherical coordinates, with D as the polar axis and $\mathbb{S}^2 \cap D = \{N, S\}$, see Figure 3. The image of the Lebesgue measure under this map is a surface measure on \mathbb{S}^2 of the form

$$S(v, \chi) \sin \chi \, d\chi \, d\phi,$$

and $S(v, \chi)$ is the scattering cross-section in the direction χ corresponding to the relative speed v . Because S is the density with respect to the Euclidean surface element on the unit sphere (which is dimensionless) of the image of the two-dimensional Lebesgue measure (which has the dimension of a surface) under the map (3.32), it has the dimension of a surface, which justifies the name ‘‘cross-section’’.

The scattering cross-section clearly depends upon the computation of the deflection angle in (3.31)

$$S(v, \chi) = \frac{h}{\sin \chi |\chi'_*(h)|} \Big|_{\chi_*(h)=\chi}.$$

Here are the scattering cross-sections for a few typical interactions:

- in the case of a hard sphere interaction,

$$U(r) = 0 \quad \text{if } r \geq d_0 \quad \text{and} \quad U(r) = +\infty \quad \text{if } 0 < r < d_0.$$

In this example, U is not decreasing but only nonincreasing and has finite range. Therefore, the definition of the scattering cross-section must be modified as follows. The map (3.32) is replaced with

$$\begin{aligned} B(0, d_0) &\simeq (0, d_0) \times [0, 2\pi) \rightarrow [0, \pi) \times [0, 2\pi) \simeq \mathbb{S}^2 \setminus \{N, S\}, \\ (h, \phi) &\mapsto (\chi_*(h), \phi), \end{aligned} \tag{3.33}$$

and $S(v, \chi) \sin \chi \, d\chi \, d\phi$ is the image under the above map of the restriction to the disk $B(0, d_0) \subset \mathbb{R}^2$ of the Lebesgue measure. With this slightly modified definition, it is found that

$$S(v, \chi) = \frac{1}{4} d_0^2;$$

- if $U(r) = kr^{-s}$ with $s > 0$, set

$$\vartheta(l) = \int_0^{\zeta(l)} \frac{dz}{\sqrt{1 - z^2 - 2(z/l)^s}},$$

where $l = (v^2/2k)^{1/s}h$, and where $\zeta(l)$ is the only positive root of the denominator of the integrand above; set $\vartheta \mapsto \lambda(\vartheta)$ to be the inverse of $l \mapsto \vartheta(l)$ so defined. Then

$$S(v, \chi) = (2k)^{2/s} v^{-4/s} \frac{\beta((\pi - \chi)/2)}{\sin \chi},$$

where $\beta(\vartheta) = \lambda(\vartheta)\lambda'(\vartheta)$. One finds that β is singular near $\theta = \pi/2$, which corresponds to $\chi = 0$ and $l \rightarrow +\infty$, i.e., to collisions with small deflection angles, or equivalently to the case of *grazing collisions*

$$\beta(\theta) \simeq C \left(\frac{\pi}{2} - \theta \right)^{-1-2/s} \quad \text{as } \theta \rightarrow \frac{\pi}{2}^- ,$$

while $\beta(\theta) = O(\theta)$ for $\theta \rightarrow 0$.

Although the usual definition of the scattering cross-section involves the deflection angle χ , one might find it easier to use instead the angle $\theta = (\pi - \chi)/2$ (see Figure 2). It is easily seen that the scattering cross-section S as above can be expressed in terms of a function $\Sigma \equiv \Sigma(v, \mu)$ defined on $\mathbb{R}_+ \times [0, 1)$ by the formula

$$\Sigma(v, |\cos \theta|) = \frac{1}{2} S(v, \pi - 2\theta) |\cos \theta|.$$

This function Σ has the following geometric interpretation: the mapping

$$\mathbb{S}^2 \setminus \{N, S\} \simeq (0, \pi) \times [0, 2\pi) \rightarrow \mathbb{S}^2 \setminus \{N\},$$

$$\omega \simeq (\theta, \phi) \mapsto (-\cos(2\theta), \sin(2\theta) \cos \phi, \sin(2\theta) \sin \phi)$$




Fig. 4. The double cover $\theta \mapsto \chi$.

is a double cover (see Figure 4) and the image of the surface measure $\Sigma(v, |\cos \theta|) \times \sin \theta \, d\theta \, d\phi$ under the above mapping is $2S(v, \chi) \sin \chi \, d\chi \, d\phi$.

Then, the collision kernel b is given by

$$b(V, \omega) = |V| \Sigma(|V|, |\cos(\widehat{V}, \omega)|).$$

With the formulas for the scattering cross-section given above, one sees that

- in the case of hard-spheres with radius r_0 ,

$$b(V, \omega) = \frac{1}{2} r_0^2 |V \cdot \omega|; \tag{3.34}$$

- in the case of an interaction potential $U(r) = kr^{-s}$ for $s > 0$,

$$b(V, \omega) = \frac{1}{4} (2k)^{2/s} |V|^{1-4/s} \frac{\beta(\theta)}{\sin \theta} \quad \text{with } \theta = (\widehat{V}, \omega) \tag{3.35}$$

with $\beta(\theta) = O(\theta)$ as $\theta \rightarrow 0^+$ while $\beta(\theta) = O((\pi/2 - \theta)^{-1-2/s})$ as $\theta \rightarrow \pi^-/2$;

- whenever $s = 4$, the collision kernel b is independent of $|V|$; such potentials are usually referred to as *Maxwellian* potentials, and considerably facilitate the analysis of the collision integral;
- for $s = 1$, which is the case of a repulsive Coulomb potential, one has

$$b(V, \omega) = k^2 |V|^{-3} \frac{\beta(\theta)}{\sin \theta} \quad \text{with } \beta(\theta) = O\left(\left(\frac{\pi}{2} - \theta\right)^{-3}\right) \text{ as } \theta \rightarrow \frac{\pi}{2}. \tag{3.36}$$

Observe that, for any inverse power-law potential $U(r) = kr^{-s}$,

$$\int_{\mathbb{S}^2} b(V, \omega) \, d\omega = +\infty \tag{3.37}$$

1 because of the singularity at $\theta = \pi/2$, so that our earlier assumption (3.8) is violated 1
 2 by such potentials. In particular, whenever the particle interaction is given by an inverse 2
 3 power-law potential, one cannot split the collision integral as 3
 4

$$5 \quad \mathcal{B}(F, F) = \mathcal{B}_+(F, F) - \mathcal{B}_-(F, F) \quad 5$$

6
 7 as was done earlier in this section. One way around this is to define $\mathcal{B}(F, F)$ in the sense 7
 8 of distributions, as follows 8
 9

$$10 \quad \langle \mathcal{B}(F, F), \phi \rangle \quad 10$$

$$11 \quad = \frac{1}{4} \iint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F' F'_* - F F_*') (\phi + \phi_* - \phi' - \phi_*') b(v - v_*, \omega) dv dv_* d\omega \quad 11$$

12
 13 for each $F \in C_c^1(\mathbb{R}^3)$ and $\phi \in C_c^1(\mathbb{R}^3)$. Then 13
 14

$$15 \quad (F' F'_* - F F_*') (\phi + \phi_* - \phi' - \phi_*') = O(|(v - v_*) \cdot \omega|^2) \quad 15$$

16
 17 so that the integrand in the right-hand side above is $O((\pi/2 - \theta)^{1-2/s})$. This procedure 16
 18 can handle all inverse power-law potentials $U(r) = kr^{-s}$ for $s > 1$; however, the Coulomb 17
 19 case $s = 1$ remains excluded. 18
 20

21 Observe that, in addition to the singularity in the deflection angle at $\theta = \pi/2$, the col- 21
 22 lision kernel of the Coulomb potential also has a $|v - v_*|^{-3}$ singularity in the velocity 22
 23 variable. Physicists deal with the latter singularity by introducing a further truncation near 23
 24 $V = 0$; the dependence upon this truncation parameter of the collision integral is only loga- 24
 25 rithmic, so that the result does not depend “too much” on the truncation parameter. In other 25
 26 words, the collision integral is computed modulo a scaling factor known as the *Coulomb* 26
 27 *logarithm* (see [76], Section 41). 27
 28

29 Another way of avoiding the singularity of the collision kernel at $\theta = \pi/2$ consists in 29
 30 assuming that the interaction potential is truncated at large distance, in other words, that 30
 31

$$32 \quad U(r) = U(r_C) \quad \text{whenever } r \geq r_C. \quad (3.38) \quad 32$$

33
 34 In this case (as in the hard sphere case), U is not decreasing but only nonincreasing, and the 34
 35 definition of the scattering cross-section must be modified as in (3.33) with r_C in the place 35
 36 of d_0 . Call b_C the collision kernel corresponding to the potential truncated as in (3.38), 36
 37 and S_C, Σ_C the associated scattering cross-section. Then 37
 38

$$39 \quad \int_{\mathbb{S}^2} b_C(V, \omega) d\omega = |V| \int_0^{2\pi} \int_0^\pi \Sigma_C(|V|, |\cos \theta|) \sin \theta d\theta d\phi \quad 39$$

$$40 \quad = |V| \int_0^{2\pi} \int_0^\pi S_C(|V|, \chi) \sin \chi d\chi d\phi \quad 40$$

$$41 \quad = |V| \int_0^{r_C} \int_0^{2\pi} h dh d\phi = |V| \pi r_C^2 \quad 41$$

1 since $S_C(v, \chi) \sin \chi \, d\chi \, d\phi$ is the image under the map (3.33) of the Lebesgue measure on 1
 2 the two-dimensional disk of radius r_C . Hence the truncation (3.38) leads to a collision ker- 2
 3 nel whose integral over the angle variables is finite, thereby avoiding the divergence (3.37) 3
 4 that occurs for any infinite range, inverse power law potential. 4

5 Yet another way of avoiding the singularity of the collision kernel at $\theta = \pi/2$ was pro- 5
 6 posed by Grad [64]. He considered molecular force laws for which the collision kernel 6
 7 satisfies the condition 7

$$8 \quad \frac{b(V, \omega)}{|\cos(V, \omega)|} \leq C(|V| + |V|^{1-\varepsilon}), \quad (3.39) \quad 9$$

10 where $C > 0$ and $\varepsilon \in (0, 1)$. Comparing the case of a power law potential $U(r) = kr^{-s}$ 10
 11 with Grad's assumption above, we see that the collision kernel in (3.35) satisfies (3.39) 11
 12 provided that one modifies the function β near $\theta = \pi/2$ so that $\beta(\theta) = O(\pi/2 - \theta)$. One 12
 13 possibility is to replace β with 13
 14 15

$$16 \quad \tilde{\beta}(\theta) = \beta(\theta) \mathbb{1}_{\theta \leq \theta_0}, \quad (3.40) \quad 17$$

18 where $\theta_0 \in (0, \frac{\pi}{2})$ is some arbitrary value. Then, the associated truncated collision kernel 18
 19 \tilde{b} satisfies the bounds 19
 20 21

$$22 \quad \tilde{b}(V, \omega) \leq C_b(1 + |V|)^{1-4/s} \quad \text{and} \quad \int_{\mathbb{S}^2} b(V, \omega) \, d\omega \geq \frac{1}{C_b}(1 + |V|)^{1-4/s} \quad 23$$

24 25 26 for some positive constant C_b . The potential U is called a *hard cut-off potential* if $s \geq 4$, 27
 28 and a *soft cut-off potential* if $s < 4$.

29 Grad defined more general classes of hard and soft cut-off potentials; specifically, a gen- 29
 30 eral hard cut-off potential corresponds to the condition 30
 31

$$32 \quad \int_{\mathbb{S}^2} b(V, \omega) \, d\omega \geq \frac{c|V|}{1 + |V|} \quad 32$$

33 34 while a soft cut-off potential is defined by the condition 35
 36

$$37 \quad \int_{\mathbb{S}^2} b(V, \omega) \, d\omega \leq c(1 + |V|^{\varepsilon-1}) \quad 37$$

38 39 for some $c > 0$ and $\varepsilon \in (0, 1)$, in addition to the bound (3.39). In the sequel, we shall mostly 40
 41 restrict our attention to those hard cut-off potentials that satisfy the same bound (3.41) as 41
 42 in the inverse power law case. 42

43 The terminology "cut-off potential" attached to Grad's cut-off prescription, is somewhat 43
 44 unfelicitous. Indeed, it is not equivalent to truncating the potential at large intermolecu- 44
 45 lar distances as in (3.38). Indeed, the angular truncation (3.40) prohibits grazing collisions 45

with deflection angle less than a threshold $\pi - 2\theta_0$ that is independent of the relative velocity V . On the contrary, with a potential truncated as in (3.38), there exist grazing collisions with deflection angle arbitrarily small for large enough relative velocity $|V|$. Hence, it would be more appropriate to refer to Grad's procedure as leading to a "cut-off scattering cross-section" rather than a "cut-off potential". Yet the latter terminology is commonly used in the literature, so that changing it would only cause confusion.

Let us conclude this subsection with a few words on the physical relevance of Grad's cut-off assumption. Grad observed that, in gases of neutral particles with short range interactions, grazing collisions are not statistically dominant as in the case of plasmas, where the long range effect of the Coulomb interaction must be accounted for. The latter case requires using a mean-field description of the long-range interaction potential, in addition to the Landau collision integral, an approximation of Boltzmann's collision integral in the regime of essentially grazing collisions.

That the Coulomb potential, probably the best known interaction in physics, is a singular point in the theory of the Boltzmann collision integral may seem highly regrettable. However, in view of the remark above, the reader should bear in mind that the Boltzmann equation is essentially meant to model collisional processes in neutral gases with short-range molecular interactions, and not in plasmas, so that the Coulomb potential is not really relevant in this context.

3.6. The linearized collision integral

Let ρ and $\theta > 0$, and pick $u \in \mathbb{R}^3$; the linearization at $\mathcal{M}_{(\rho,u,\theta)}$ of Boltzmann's collision integral is defined as follows

$$\mathcal{L}\mathcal{M}_{(\rho,u,\theta)}f = -2\mathcal{M}_{(\rho,u,\theta)}^{-1}\mathcal{B}(\mathcal{M}_{(\rho,u,\theta)}, \mathcal{M}_{(\rho,u,\theta)}f), \quad (3.42)$$

where \mathcal{B} is the bilinear operator obtained by polarization from the Boltzmann collision integral. In other words,

$$\mathcal{L}\mathcal{M}_{(\rho,u,\theta)}f = \iint_{\mathbb{R}^3 \times \mathbb{S}^2} (f + f_* - f' - f'_*)b(v - v_*, \omega)\mathcal{M}_{(\rho,u,\theta)}(v_*) dv_* d\omega, \quad (3.43)$$

where f_* , f' and f'_* are the values of f at v_* , v' and v'_* , respectively, and where v' and v'_* are determined in terms of v , v_* and ω by the usual collision relations (3.3).

The dependence on the parameters ρ , u and θ of the linearized collision integral is handled most easily by using the translation and scaling invariance of \mathcal{L} .

3.6.1. Translation and scaling invariance of $\mathcal{L}\mathcal{M}_{(\rho,u,\theta)}$. We introduce the following notation for the actions of translation and scaling transformations on functions defined on \mathbb{R}^3 :

$$\tau_u f(v) = f(v - u), \quad m_\lambda f(v) = \lambda^{-3} f\left(\frac{v}{\lambda}\right). \quad (3.44)$$

1 For instance, with these notations

$$2 \mathcal{M}_{(\rho,u,\theta)} = \rho m_{\sqrt{\theta}} \tau_u \mathcal{M}_{(1,0,1)}. \quad (3.45) \quad 3$$

4
5 Given a collision kernel $b \equiv b(z, \omega)$, we denote by \mathcal{B}^b Boltzmann's collision integral de-
6 fined by this collision kernel. With the notation so defined, a straightforward change of
7 variables in the collision integral leads to the following relation

$$8 \tau_u \mathcal{B}^b(\Phi, \Phi) = \mathcal{B}^b(\tau_u \Phi, \tau_u \Phi), \quad 9$$

$$10 m_\lambda \mathcal{B}^b(\Phi, \Phi) = \lambda^3 \mathcal{B}^{m_\lambda b}(m_\lambda \Phi, m_\lambda \Phi) \quad (3.46) \quad 11$$

12
13 for each continuous, rapidly decaying $\Phi \equiv \Phi(v)$, where, in the expression $m_\lambda b$, it is un-
14 derstood that the scaling transformation acts on the first argument of b , i.e., on the relative
15 velocity. The analogous formula for the linearized collision operator is

$$16 m_{\sqrt{\theta}} \tau_u \mathcal{L}_{\mathcal{M}_{(1,0,1)}}^b \phi = \rho \theta^{3/2} \mathcal{L}_{\mathcal{M}_{(\rho,u,\theta)}}^{m_{\sqrt{\theta}} b} (m_{\sqrt{\theta}} \tau_u \phi). \quad (3.47) \quad 17$$

18
19 This relation shows that it is enough to study the linearization of the collision integral at
20 the centered reduced Gaussian

$$21 M = \mathcal{M}_{(1,0,1)} \quad 22$$

23 with an arbitrary collision kernel b . 24

25 **3.6.2. Rotational invariance of $\mathcal{L}_{\mathcal{M}_{(1,0,1)}}$.** The orthogonal group $O_3(\mathbb{R})$ acts on functions
26 on \mathbb{R}^3 by the formula 27

$$28 f_R(v) = f(R^T v), \quad R \in O_3(\mathbb{R}), v \in \mathbb{R}^3; \quad (3.48) \quad 29$$

30 likewise its action on vector fields is defined by 31

$$32 U_R(v) = RU(R^T v), \quad R \in O_3(\mathbb{R}), v \in \mathbb{R}^3, \quad (3.49) \quad 33$$

34 while its action on symmetric matrix fields is given by 35

$$36 S_R(v) = RS(R^T v)R^T, \quad R \in O_3(\mathbb{R}), v \in \mathbb{R}^3. \quad (3.50) \quad 37$$

38 The Boltzmann collision integral is obviously invariant under the action of $O_3(\mathbb{R})$ – indeed,
39 the microscopic collision process is isotropic. In fact, an elementary change of variables in
40 the collision integral shows that 41

$$42 \mathcal{B}(\Phi_R, \Phi_R) = \mathcal{B}(\Phi, \Phi)_R \quad (3.51) \quad 43$$

44

45

1 for each continuous, rapidly decaying Φ . Since the centered unit Gaussian $M = \mathcal{M}_{(1,0,1)}$ 1
 2 is a radial function, this rotation invariance property goes over to \mathcal{L}_M 2

$$3 \quad \mathcal{L}_M(\phi_R) = (\mathcal{L}_M\phi)_R. \quad (3.52) \quad 4$$

5
 6 Extending \mathcal{L}_M to act componentwise on vector or matrix fields on \mathbb{R}^3 , one finds that 6
 7

$$8 \quad \mathcal{L}_M(U_R) = (\mathcal{L}_MU)_R \quad (3.53) \quad 9$$

10 for continuous, rapidly decaying vector fields U and 11

$$12 \quad \mathcal{L}_M(S_R) = (\mathcal{L}_MS)_R \quad (3.54) \quad 13$$

14
 15 for continuous, rapidly decaying symmetric matrix fields S , where the notations U_R and S_R 15
 16 are as in (3.48)–(3.50). 16

17 As we shall see below, this $O_3(\mathbb{R})$ -invariance of \mathcal{L}_M has important consequences: it 17
 18 implies in particular that the viscosity and heat conductivity are scalar quantities (and not 18
 19 matrices). 19

20
 21 **3.6.3. The Fredholm property.** Henceforth, we assume that the collision kernel b satisfies 21
 22 a hard cut-off assumption in the sense of Grad [64], i.e., there exists $\alpha \in [0, 1]$ and $C_b > 0$ 22
 23 such that, for a.e. $z \in \mathbb{R}^3$ and $\omega \in \mathbb{S}^2$, one has 23
 24

$$25 \quad 0 < b(z, \omega) \leq C_b(1 + |z|)^\alpha \quad \text{and} \quad (3.55) \quad 25$$

$$26 \quad \int_{\mathbb{S}^2} b(z, \omega) d\omega \geq \frac{1}{C_b}(1 + |z|)^\alpha. \quad 26$$

27
 28 Consider \mathcal{L} , the linearization of the Boltzmann collision integral at the centered, reduced 27
 29 Gaussian state M above. (Notice that we have discarded the dependence of \mathcal{L} on b and M 28
 30 for notational simplicity.) From (3.43), we infer that \mathcal{L} can be split as the sum of a local 29
 31 (multiplication) operator and of an integral operator, as follows 30
 32

$$33 \quad \mathcal{L}\phi(v) = a(|v|)\phi(v) - \mathcal{K}\phi(v), \quad (3.56) \quad 34$$

35 where a is the collision frequency 35
 36

$$37 \quad a(|v|) = \iint_{\mathbb{R}^3 \times \mathbb{S}^2} b(v - v_*, \omega) M_* dv_* d\omega. \quad (3.57) \quad 38$$

39
 40 The nonlocal operator \mathcal{K} is further split into two parts 40
 41

$$42 \quad \mathcal{K}\phi = \mathcal{K}_1\phi - \mathcal{K}_2\phi, \quad (3.58) \quad 43$$

1 where

$$\begin{aligned}
 \mathcal{K}_1\phi &= \iint_{\mathbb{R}^3 \times \mathbb{S}^2} \phi_* b(v - v_*, \omega) M_* dv_* d\omega \\
 \mathcal{K}_2\phi &= \iint_{\mathbb{R}^3 \times \mathbb{S}^2} (\phi' + \phi'_*) b(v - v_*, \omega) M_* dv_* d\omega \\
 &= 2 \iint_{\mathbb{R}^3 \times \mathbb{S}^2} \phi' b(v - v_*, \omega) M_* dv_* d\omega.
 \end{aligned}
 \tag{3.59}$$

2
3
4
5
6
7
8
9
10
11 (This last formula is not entirely obvious; it rests on a change of variables that will be
12 explained later.) It is clear that \mathcal{K}_1 is a compact operator on $L^2(M dv)$; that \mathcal{K}_2 shares the
13 same property is much less obvious, and was proved by Hilbert in the hard sphere case.
14 Fifty years later, Grad introduced the cut-off assumption which now bears his name and
15 used it in particular to extend Hilbert's result to all cut-off potentials.

16
17 LEMMA 3.9 (Hilbert [69], Grad [64]). *Assume that b is a collision kernel that satisfies the*
18 *hard cut-off assumption (3.55). Then the operator \mathcal{K}_2 is compact on $L^2(M dv)$.*

19
20
21 We shall not give the proof of the Hilbert–Grad lemma here, since it is rather long and
22 technical; the interested reader is referred to the lucid exposition by Glassey [50], or to the
23 treatise by Cercignani, Illner and Pulvirenti [28] for the hard sphere case.

24 Instead, we shall digress from our discussion of the linearized collision integral and
25 mention a new result by Lions which can be viewed as a nonlinear analogue of the Hilbert–
26 Grad result.

27
28 LEMMA 3.10 (Lions [84]). *Assume that b is the collision kernel of a hard sphere gas:*
29 *$b(z, \omega) = |z \cdot \omega|$, and consider the gain term in the Boltzmann collision integral*

$$\mathcal{B}_+(F, F) = \iint_{\mathbb{R}^3 \times \mathbb{S}^2} F' F'_* |(v - v_*) \cdot \omega| dv_* d\omega.$$

30
31
32
33
34 Then \mathcal{B}_+ maps $L^2_{\text{comp}}(\mathbb{R}^3)$ continuously into $H^1_{\text{loc}}(\mathbb{R}^3)$.

35
36
37 Actually, this is not exactly Lions' result; in the reference [84], only compactly supported
38 collision kernels are considered. The proof in [84] is based upon the L^2 -continuity of
39 Fourier integral operators of order 0. The statement above was later proved by Bouchut
40 and Desvillettes [17] by a very simple and elegant argument.

41 Deriving the Hilbert–Grad lemma from the Lions lemma would require additional es-
42 timates in order to handle contributions from large $|v|$. Since obtaining these estimates
43 requires essentially as much work as does the proof of the Hilbert–Grad lemma, we do not
44 claim that the Lions lemma leads to a simpler proof of the compactness of \mathcal{K} ; it is however
45 of great independent interest.

The main properties of \mathcal{L} are now summarized in the following theorem.

THEOREM 3.11. *Assume that the collision kernel b satisfies the hard cut-off assumption (3.55). Then \mathcal{L} is an unbounded self-adjoint nonnegative Fredholm operator, with domain $D(\mathcal{L}) = L^2(a^2 M dv)$ (a being the collision frequency defined in (3.57)). Its nullspace is the space of collision invariants*

$$\text{Ker } \mathcal{L} = \text{span}\{1, v_1, v_2, v_3, |v|^2\}.$$

Finally, \mathcal{L} satisfies the following relative coercivity property on $(\text{Ker } \mathcal{L})^\perp$: there exists $C_0 > 0$ such that, for each $\phi \in L^2(a M dv)$ – the form domain of \mathcal{L} – one has

$$\int_{\mathbb{R}^3} \phi \mathcal{L} \phi M dv \geq C_0 \int_{\mathbb{R}^3} (\phi - \Pi \phi)^2 a M dv, \tag{3.60}$$

where Π is the $L^2(M dv)$ -orthogonal projection on $\text{Ker } \mathcal{L}$.

SKETCH OF THE PROOF. Let us briefly explain how these various facts are established. It follows from the Hilbert–Grad lemma that \mathcal{K} is a compact operator on $L^2(M dv)$. Hence $\mathcal{L} = a - \mathcal{K}$ is an unbounded Fredholm operator with domain $D(\mathcal{L}) = L^2(a^2 M dv)$. That \mathcal{L} is self-adjoint comes from the symmetries of Boltzmann’s collision integral, especially from the computation of

$$\int_{\mathbb{R}^3} \mathcal{B}(F, F) \phi dv$$

as explained in Proposition 3.1. By the same argument, one sees that

$$\int_{\mathbb{R}^3} \phi \mathcal{L} \phi M dv = \frac{1}{4} \iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (\phi + \phi_* - \phi' - \phi'_*)^2 b M dv M_* dv_* d\omega \geq 0 \tag{3.61}$$

so that \mathcal{L} is a nonnegative operator; moreover, if $\mathcal{L} \phi = 0$, the integral on the right-hand side of the above equality vanishes, so that

$$\phi + \phi' - \phi' - \phi'_* = 0 \quad \text{a.e. in } v, v_*, \omega.$$

Hence ϕ is a collision invariant (see Definition 3.5) and this characterizes $\text{Ker } \mathcal{L}$, see Proposition 3.6. As for the last inequality, the spectral theory of compact operators implies that \mathcal{L} has a spectral gap in $L^2(M dv)$

$$\int_{\mathbb{R}^3} \phi \mathcal{L} \phi M dv \geq C^* \int_{\mathbb{R}^3} (\phi - \Pi \phi)^2 M dv. \tag{3.62}$$

1 On the other hand, by continuity of \mathcal{K} on $L^2(M dv)$, one has

$$\begin{aligned} 2 \int_{\mathbb{R}^3} \phi \mathcal{L} \phi M dv &= \int_{\mathbb{R}^3} (\phi - \Pi \phi) \mathcal{L} (\phi - \Pi \phi) M dv \\ 3 &\geq \int_{\mathbb{R}^3} (\phi - \Pi \phi)^2 a M dv - \|\mathcal{K}\|^2 \int_{\mathbb{R}^3} (\phi - \Pi \phi)^2 M dv. \end{aligned}$$

4
5
6
7
8 Combining both inequalities leads to the announced estimate. \square

9
10 The improved spectral gap estimate above is due to Bardos, Caflisch and Nicolaenko [6].
11 Also, an elegant explicit estimate for the $L^2(M dv)$ spectral gap can be found in [5].

12 In fact, it is interesting to notice that the weighted spectral gap estimate (3.60) is in some
13 sense more intrinsic than the unweighted one. Indeed, in the case of soft cut-off potentials,
14 (3.62) is false, but (3.60) (which is of course a weaker statement since the collision fre-
15 quency vanishes for large velocities in the soft potential case) holds true: this was proved
16 in [58], based on decay estimates on the gain term in the linearized collision integral due to
17 Grad and Caflisch [23]. In other words, while the spectral gap estimate (??) holds for hard
18 potentials only, the weighted spectral gap estimate holds for hard as well as soft potentials,
19 in the cut-off case.

20 An important consequence of Theorem 3.11 is that the integral equation

$$21 \mathcal{L} \phi = \psi, \quad \psi \in L^2(M dv), \tag{3.63}$$

22 satisfies the *Fredholm alternative*:

- 23 • either $\psi \perp \text{Ker } \mathcal{L}$, in which case (3.63) has a unique solution

$$24 \phi_0 \in L^2(a^2 M dv) \cap (\text{Ker } \mathcal{L})^\perp;$$

25 then any solution of (3.63) is of the form

$$26 \phi = \phi_0 + \phi_1, \quad \text{where } \phi_1 \text{ is an arbitrary element of } \text{Ker } \mathcal{L};$$

- 27 • or $\psi \notin (\text{Ker } \mathcal{L})^\perp$, in which case (3.63) has no solution.

28
29
30
31
32
33
34 EXAMPLE. Consider the vector field

$$35 B(v) = \frac{1}{2}(|v|^2 - 5)v$$

36 and the matrix field

$$37 A(v) = v \otimes v - \frac{1}{3}|v|^2 I.$$

38
39
40
41
42
43 Clearly

$$44 A_{jk} \perp \text{Ker } \mathcal{L}, \quad B_l \perp \text{Ker } \mathcal{L}, \quad A_{jk} \perp B_l, \quad j, k, l = 1, 2, 3. \tag{3.64}$$

1 In fact, more is true

2
3
$$\int_{\mathbb{R}^3} A(v) f(|v|^2) M dv = 0,$$

4
5
$$\int_{\mathbb{R}^3} A(v) v f(|v|^2) M dv = 0,$$

6
7
$$\int_{\mathbb{R}^3} B(v) f(|v|^2) M dv = 0,$$

8
9
$$\int_{\mathbb{R}^3} B(v) \cdot v M dv = 0.$$

10
11 The second and third formulas are obvious since A is even and B odd. As for the first
12 formula, observe that A is an isotropic matrix, in the sense that $A(Rv) = RA(v)R^T$
13 for each $R \in O_3(\mathbb{R})$ – with the notation (3.50) for the action of $O_3(\mathbb{R})$ on symmetric matrices,
14 $A_R = A$ for each $R \in O_3(\mathbb{R})$. Hence the matrix

15
16
$$\int_{\mathbb{R}^3} A(v) f(|v|^2) M dv$$

17
18 commutes with any $R \in O_3(\mathbb{R})$ – as can be seen by changing v into Rv in the above
19 integral – and is therefore a scalar multiple of the identity matrix. But

20
21
$$\text{trace} \int_{\mathbb{R}^3} A(v) f(|v|^2) M dv = \int_{\mathbb{R}^3} \text{trace} A(v) f(|v|^2) M dv = 0$$

22
23 and hence this scalar multiple of the identity matrix is null. The fourth and last formula is
24 based on the following elementary recursion formula for Gaussian integrals

25
26
$$\int_{\mathbb{R}^3} |v|^n M dv = (n + 1) \int_{\mathbb{R}^3} |v|^{n-2} M dv, \quad n \geq 2 \tag{3.65}$$

27
28 (to see this, use spherical coordinates and integrate by parts).

29
30 In particular, the Fredholm alternative implies the existence of a matrix field \widehat{A} and of a
31 vector field \widehat{B} such that

32
33
$$\begin{aligned} \mathcal{L}\widehat{A} &= A & \text{and} & & \widehat{A} &\perp \text{Ker } \mathcal{L}, \\ \mathcal{L}\widehat{B} &= B & \text{and} & & \widehat{B} &\perp \text{Ker } \mathcal{L}. \end{aligned} \tag{3.66}$$

34
35 Observe that

36
37
$$\begin{aligned} \mathcal{L}(\widehat{A}_R) &= A_R = A, & \widehat{A}_R &\perp \text{Ker } \mathcal{L} & \text{for all } R \in O_3(\mathbb{R}), \\ \mathcal{L}(\widehat{B}_R) &= B_R = B, & \widehat{B}_R &\perp \text{Ker } \mathcal{L} & \text{for all } R \in O_3(\mathbb{R}), \end{aligned}$$

so that, by the uniqueness part in the Fredholm alternative

$$\widehat{A}_R = \widehat{A} \quad \text{and} \quad \widehat{B}_R = \widehat{B} \quad \text{for all } R \in O_3(\mathbb{R}).$$

An elementary geometric argument (see [34]) shows the existence of two scalar functions

$$\mathbf{a}: \mathbb{R}_+ \rightarrow \mathbb{R}, \quad \mathbf{b}: \mathbb{R}_+ \rightarrow \mathbb{R}$$

such that

$$\widehat{A}(v) = \mathbf{a}(|v|^2)A(v) \quad \text{and} \quad \widehat{B}(v) = \mathbf{b}(|v|^2)B(v). \quad (3.67)$$

As we shall see further, the viscosity and heat conductivity of a gas are expressed as Gaussian integrals of the scalar functions \mathbf{a} and \mathbf{b} , and therefore are scalar quantities themselves.

4. Hydrodynamic scalings for the Boltzmann equation

This short section introduces in particular the Knudsen number, Kn , a dimensionless parameter of considerable importance for the derivation of hydrodynamic models from the kinetic theory of gases.

4.1. Notion of a rarefied gas

Think of a monatomic gas as a cloud of \mathcal{N} hard spheres of radius r confined in a container of volume V ; we shall call *excluded volume* the volume V_e that the \mathcal{N} gas molecules would occupy if tightly packed somewhere in the container (as oranges in a grocery store). Clearly

$$\frac{4}{3}\mathcal{N}\pi r^3 \leq V_e \leq \mathcal{N}(2r)^3,$$

and we shall call a *perfect gas* one for which $V_e \ll V$. It is well known that the equation of state for a perfect gas is given by the Boyle–Mariotte law

$$p = k\rho\theta,$$

where p , ρ and θ are the pressure, the density and the temperature in the gas, while k designates the Boltzmann constant ($k = 1.38 \cdot 10^{-23} \text{ J K}^{-1}$).

EXAMPLE. For a monatomic gas at room temperature and atmospheric pressure, about $\mathcal{N} = 10^{20}$ molecules with radius 10^{-8} cm are to be found in a volume V of 1 cm^3 . Hence, the excluded volume satisfies $V_e \leq 10^{20} \cdot (2 \cdot 10^{-8})^3 = 8 \cdot 10^{-4} \text{ cm}^3$, so that $V_e \ll V$.

1 Another important notion is that of *mean free path*: it is the average distance between 1
 2 two successive collisions for one gas molecule picked at random. There is more than one 2
 3 precise definition of that notion, in particular, because there are several choices of proba- 3
 4 bility measures for computing the mean. One choice could be to use the empirical notion 4
 5 of mean: Pick a typical particle, wait until after this particle has collided $n \gg 1$ times with 5
 6 other particles; the ratio of the distance traveled by the particle between the initial time and 6
 7 the last collision time to the number n of collisions should converge, in the large n limit, 7
 8 to one notion of mean free path. This definition, however, does not provide us with an easy 8
 9 way of estimating the mean free path, because it is fairly difficult to compute the trajectory 9
 10 of one particle in the cloud of the $\mathcal{N} - 1$ other particles. 10

11 Intuitively, one expects that the larger the number of particles in the container, the 11
 12 smaller the mean free path; likewise, the bigger the particles, the smaller the mean free 12
 13 path. This suggests using the following formula to estimate the order of magnitude of the 13
 14 mean free path, 14

$$15 \text{ mean free path} \simeq \frac{1}{(\mathcal{N}/(V - V_e)) \times \mathcal{A}}, \quad 15$$

16 where \mathcal{A} is the surface area of the section of the particles. A mathematical justification of 16
 17 the above formula for the mean free path can be found in [111]; see also [39]. 17

18 EXAMPLE. For the same monatomic gas as above, $\mathcal{N} = 10^{20}$ molecules, $V - V_e \simeq$ 18
 19 $V = 1 \text{ cm}^3$, while $\mathcal{A} = \pi \cdot (10^{-8})^2 \text{ cm}^2$. This gives a mean free path of the order 19
 20 of $3 \cdot 10^{-5} \text{ cm} \ll 1 \text{ cm}$ (the size of the container). Hence, a gas molecule will bump 20
 21 into roughly 10^4 other particles when traveling a distance comparable to the size of the 21
 22 container. 22
 23
 24
 25
 26

27 The example above shows that even in the case of a perfect gas (i.e., for an excluded 27
 28 volume negligible when compared to the size of the container) any given particle can col- 28
 29 lide with a large number of other particles. See, for instance, [2] for more information on 29
 30 the importance of the excluded volume in this context. 30
 31

32 EXAMPLE. For the same monatomic gas as above, lower the pressure from $p = 1 \text{ atm}$ to 32
 33 $p = 10^{-4} \text{ atm}$. Then, only $\mathcal{N} = 10^{16}$ molecules are to be found in the container, which 33
 34 gives a mean free path of the order of 0.3 cm , comparable to the size of the container. 34
 35

36 These examples suggest that the natural way of measuring the degree of rarefaction in 36
 37 the gas is by using the Knudsen number, Kn , defined as the ratio of the mean free path to 37
 38 the size of the container, or more generally, as 38
 39

$$40 Kn = \frac{\text{mean free path}}{\text{macroscopic length scale}}. \quad 40$$

41 In other words, 41
 42

- 43 • a rarefied gas is a gas for which $Kn \gtrsim 1$, while 43
 - 44 • a gas in hydrodynamic regime satisfies $Kn \ll 1$. 44
- 45 45

4.2. The dimensionless Boltzmann equation

In the discussion above, the introduction of the Knudsen number was based on physical arguments. We shall see below that it also appears naturally when writing the Boltzmann equation in dimensionless variables.

Choose a macroscopic length scale L and time scale T , and a reference temperature Θ . This defines *two* velocity scales:

- one is the speed at which some macroscopic portion of the gas is transported over a distance L in time T , i.e.,

$$U = \frac{L}{T};$$

- the other one is the *thermal speed* of the molecules with energy $\frac{3}{2}k\Theta$; in fact, it is more natural to define this velocity scale as

$$c = \sqrt{\frac{5}{3} \frac{k\Theta}{m}},$$

m being the molecular mass, which is the *speed of sound* in a monatomic gas at the temperature Θ .

Define next the dimensionless variables involved in the Boltzmann equation, i.e., the dimensionless time, space and velocity variables as

$$\hat{t} = \frac{t}{T}, \quad \hat{x} = \frac{x}{L} \quad \text{and} \quad \hat{v} = \frac{v}{c}.$$

Define also the dimensionless number density

$$\hat{F}(\hat{t}, \hat{x}, \hat{v}) = \frac{L^3 c^3}{\mathcal{N}^3} F(t, x, v),$$

where \mathcal{N} is the total number of gas molecules in a volume L^3 . Finally, we must rescale the collision kernel b . As mentioned earlier, $b(z, \omega)$ is the relative velocity multiplied by the scattering cross-section of the gas molecules; define

$$\hat{b}(\hat{z}, \omega) = \frac{1}{c \times \pi r^2} b(z, \omega) \quad \text{with} \quad \hat{z} = \frac{z}{c},$$

where r is the molecular radius.

If f satisfies the Boltzmann equation

$$\partial_t F + v \cdot \nabla_x F = \iint_{\mathbb{R}^3 \times \mathbb{S}^2} (F' F'_* - F F_*') b(v - v_*, \omega) dv_* d\omega,$$

1 then

$$2 \quad \frac{L}{cT} \partial_t \widehat{F} + \widehat{v} \cdot \nabla_{\widehat{x}} \widehat{F} = \frac{\mathcal{N} \pi r^2}{L^2} \iint_{\mathbb{R}^3 \times \mathbb{S}^2} (\widehat{F}' \widehat{F}'_* - \widehat{F} \widehat{F}_*) \widehat{b}(\widehat{v} - \widehat{v}_*, \omega) d\widehat{v}_* d\omega. \quad 3$$

4 The factor multiplying the collision integral is

$$5 \quad L \times \frac{\mathcal{N} \times \pi r^2}{L^3} = \frac{L}{\text{mean free path}} = \frac{1}{Kn}, \quad 6$$

7 where Kn is the Knudsen number defined above. The factor multiplying the time derivative

$$8 \quad \frac{(1/T) \times L}{c} =: St \quad 9$$

10 is called the *kinetic Strouhal number* (by analogy with the notion of Strouhal number used
11 in the dynamics of vortices). Hence the dimensionless form of the Boltzmann equation (see
12 Section 2.9 in [115]) is

$$13 \quad St \partial_t \widehat{F} + \widehat{v} \cdot \nabla_{\widehat{x}} \widehat{F} = \frac{1}{Kn} \iint_{\mathbb{R}^3 \times \mathbb{S}^2} (\widehat{F}' \widehat{F}'_* - \widehat{F} \widehat{F}_*) \widehat{b}(\widehat{v} - \widehat{v}_*, \omega) d\widehat{v}_* d\omega. \quad (4.1) \quad 14$$

15 There is some arbitrariness in the way the length, time and temperature scales L , T , Θ are
16 chosen. For instance, if $F^{\text{in}} \equiv F^{\text{in}}(x, v)$ is the initial number density (at time $t = 0$), one
17 can choose

$$18 \quad \begin{aligned} 19 \quad \frac{1}{L} &= \frac{\iint_{\mathbb{R}^3 \times \mathbb{R}^3} |\nabla_x F^{\text{in}}| dx dv}{\iint_{\mathbb{R}^3 \times \mathbb{R}^3} F^{\text{in}} dx dv}, \\ 20 \quad U &= \frac{\int_{\mathbb{R}^3} |\int_{\mathbb{R}^3} v F^{\text{in}} dv| dx}{\iint_{\mathbb{R}^3 \times \mathbb{R}^3} F^{\text{in}} dx dv}, \\ 21 \quad \Theta &= \frac{1/(3k) \iint_{\mathbb{R}^3 \times \mathbb{R}^3} m |v|^2 F^{\text{in}} dx dv}{\iint_{\mathbb{R}^3 \times \mathbb{R}^3} F^{\text{in}} dx dv}, \end{aligned} \quad 22$$

23 and define $T = L/U$. In addition to Sone's book [115], we also refer to the Introduction
24 of [10] for a presentation of the Boltzmann equation in dimensionless variables.

25 All hydrodynamic limits of the Boltzmann equation correspond to situations where the
26 Knudsen number, Kn , satisfies

$$27 \quad Kn \ll 1. \quad 28$$

29 In other words, the Knudsen number governs the transition from the kinetic theory of
30 gases to hydrodynamic models, just as the Reynolds number in fluid mechanics governs
31 the transition from laminar to turbulent flows, except that the hydrodynamic limit is much
32 better understood than the latter situation. 33

But there is no universal prescription for the Strouhal number in the context of the hydrodynamic limit; as we shall see below, various hydrodynamic regimes can be derived from the Boltzmann equation by appropriately tuning the Strouhal number.

5. Compressible limits of the Boltzmann equation: Formal results

In this section, we study the dimensionless Boltzmann equation (4.1) in the case where

$$Kn = \varepsilon \ll 1, \quad St = 1.$$

The dimensionless Boltzmann equation takes the form

$$\partial_t F_\varepsilon + v \cdot \nabla_x F_\varepsilon = \frac{1}{\varepsilon} \mathcal{B}(F_\varepsilon, F_\varepsilon), \quad (5.1)$$

where

$$\mathcal{B}(F, F) = \iint_{\mathbb{R}^3 \times \mathbb{S}^2} (F' F'_* - F F_*') b(v - v_*, \omega) dv_* d\omega.$$

The collision kernel b is assumed to satisfy Grad's hard cut-off assumption

$$0 < b(z, \omega) \leq C_b (1 + |z|)^\alpha \quad \text{and} \quad \int_{\mathbb{S}^2} b(z, \omega) d\omega \geq \frac{1}{C_b} (1 + |z|)^\alpha \quad (5.2)$$

for a.e. $z \in \mathbb{R}^3$ and $\omega \in \mathbb{S}^2$, where $\alpha \in [0, 1]$ and $C_b > 0$.

5.1. The compressible Euler limit: The Hilbert expansion

In [69], Hilbert proposed to seek the solution of (5.1) as a formal power series in ε ,

$$F_\varepsilon(t, x, v) = \sum_{k \geq 0} \varepsilon^k F_k(t, x, v) \quad (5.3)$$

with $F_k \in C^\infty(\mathbb{R}_+ \times \mathbb{R}^3; \mathcal{S}(\mathbb{R}^3))$ for each $k \geq 0$.

The equations governing the coefficients F_k are obtained by inserting the right-hand side of (5.3) in (5.1) and equating the coefficients multiplying the successive powers of ε .

Order ε^{-1} . One finds

$$0 = \mathcal{B}(F_0, F_0),$$

hence F_0 is a local Maxwellian, meaning that there exists $\rho \equiv \rho(t, x) > 0$, $\theta \equiv \theta(t, x) > 0$ and $u \equiv u(t, x) \in \mathbb{R}^3$ such that

$$F_0(t, x, v) = \mathcal{M}_{(\rho(t,x), u(t,x), \theta(t,x))}(v) \quad \text{a.e.}$$

1 *Order* ε^0 . One finds

$$2 \quad (\partial_t + v \cdot \nabla_x) F_0 = \mathcal{B}(F_1, F_0) + \mathcal{B}(F_0, F_1) \quad 3$$

4 which can be recast in terms of the linearization at the local Maxwellian F_0 of the collision
5 integral (see (3.42)) as

$$6 \quad \mathcal{L}_{F_0} \left(\frac{F_1}{F_0} \right) = -(\partial_t + v \cdot \nabla_x) \ln F_0. \quad 7 \quad (5.4) \quad 8$$

9 This is precisely an integral equation for F_1/F_0 of the form (3.63) studied above.

10 Let us compute the right-hand side of the above equality, i.e., the expression

$$11 \quad (\partial_t + v \cdot \nabla_x) \ln F_0. \quad 12$$

13 For convenience, we shall denote by V the vector

$$14 \quad V = \frac{1}{\sqrt{\theta}}(v - u). \quad 15$$

16 Then

$$17 \quad (\partial_t + v \cdot \nabla_x) \ln \mathcal{M}_{(\rho, u, \theta)} \quad 18$$

$$19 \quad = \frac{1}{\rho} (\partial_t + v \cdot \nabla_x) \rho - \frac{3}{2} \frac{1}{\theta} (\partial_t + v \cdot \nabla_x) \theta \quad 20$$

$$21 \quad + \frac{v - u}{\theta} (\partial_t + v \cdot \nabla_x) u + \frac{|v - u|^2}{2\theta^2} (\partial_t + v \cdot \nabla_x) \theta. \quad 22$$

23 We shall rearrange the right-hand side of the above equality and express it as a linear
24 combination of the functions

$$25 \quad 1, \quad V_j, \quad \frac{1}{2}(|V|^2 - 3), \quad A(V)_{kl} \quad \text{and} \quad B(V)_j, \quad j, k, l = 1, 2, 3. \quad 26$$

27 One finds that

$$28 \quad (\partial_t + v \cdot \nabla_x) \ln \mathcal{M}_{(\rho, u, \theta)} \quad 29$$

$$30 \quad = \frac{1}{\rho} (\partial_t \rho + u \cdot \nabla_x \rho + \rho \operatorname{div}_x u) \quad 31$$

$$32 \quad + \frac{1}{\sqrt{\theta}} V \cdot \left(\partial_t u + u \cdot \nabla_x u + \nabla_x \theta + \frac{\theta}{\rho} \nabla_x \rho \right) \quad 32$$

$$33 \quad + \frac{1}{2} (|V|^2 - 3) \frac{1}{\theta} \left(\partial_t \theta + u \cdot \nabla_x \theta + \frac{2}{3} \theta \operatorname{div}_x u \right) \quad 33$$

$$34 \quad + A(V) : \nabla_x u + 2B(V) \cdot \nabla_x \sqrt{\theta}. \quad 34 \quad (5.5) \quad 35$$

1 Because of the orthogonality relations (3.64), the last two terms on the right-hand side 1
 2 of (5.5) belong to $(\text{Ker } \mathcal{L}_{F_0})^\perp$, while the first three terms there are in $\text{Ker } \mathcal{L}_{F_0}$. Therefore, 2
 3 the solvability condition for the Fredholm integral equation (5.4) consists in setting to 0 3
 4 the coefficients of the functions 1, V_j and $\frac{1}{2}(|V|^2 - 3)$, i.e., 4
 5

$$\begin{aligned} 6 \quad & \partial_t \rho + u \cdot \nabla_x \rho + \rho \operatorname{div}_x u = 0, \\ 7 \quad & \partial_t u + u \cdot \nabla_x u + \nabla_x \theta + \frac{\theta}{\rho} \nabla_x \rho = 0, \\ 8 \quad & \partial_t \theta + u \cdot \nabla_x \theta + \frac{2}{3} \theta \operatorname{div}_x u = 0, \end{aligned}$$

9
 10
 11 or in other words, 11
 12

$$\begin{aligned} 13 \quad & \partial_t \rho + \operatorname{div}_x(\rho u) = 0, \\ 14 \quad & \partial_t u + u \cdot \nabla_x u + \frac{1}{\rho} \nabla_x(\rho \theta) = 0, \\ 15 \quad & \partial_t \theta + u \cdot \nabla_x \theta + \frac{2}{3} \theta \operatorname{div}_x u = 0. \end{aligned} \tag{5.6}$$

16
 17
 18
 19
 20
 21 We recognize the system of Euler equations for a compressible fluid (2.7), in the case of a 21
 22 perfect monatomic gas; i.e., for $\gamma = 0$, so that the pressure law and the internal energy are 22
 23 given respectively by 23
 24

$$25 \quad \text{pressure} = \rho \theta \quad \text{and} \quad \text{internal energy} = \frac{3}{2} \theta. \tag{5.6}$$

26
 27
 28 Assuming that ρ , u and θ satisfy these Euler equations, we solve for F_1 the Fredholm 28
 29 integral equation (5.4) to find 29
 30

$$\begin{aligned} 31 \quad & F_1 = -\frac{1}{\rho} F_0(\mathbf{a}(\theta, |V|)A(V) \cdot \nabla_x u + 2\mathbf{b}(\theta, |V|)B(V) \cdot \nabla_x \sqrt{\theta}) \\ 32 \quad & + F_0\left(\frac{\rho_1}{\rho} + \frac{1}{\sqrt{\theta}} V \cdot u_1 + \frac{1}{2}(|V|^2 - 3)\frac{\theta_1}{\theta}\right). \end{aligned}$$

33
 34
 35
 36
 37 The second term on the right-hand side of the formula giving F_1 represents the arbitrary 37
 38 element of $\text{Ker } \mathcal{L}_{F_0}$ that appears in the general solution of the integral equation (3.63), 38
 39 while the scalar quantities $\mathbf{a}(\theta, V)$ and $\mathbf{b}(\theta, V)$ satisfy (see (3.67)) 39
 40

$$41 \quad \mathcal{L}_{F_0}(\mathbf{a}(\theta, |V|)A(V)) = A(V) \quad \text{and} \quad \mathcal{L}_{F_0}(\mathbf{b}(\theta, |V|)B(V)) = B(V). \tag{3.67}$$

42
 43
 44 More precisely, let b be the collision kernel satisfying (5.2) considered in this chapter. 43
 45 Applying the notation in Section 3.6.1, and especially (3.44), we define scalar functions 44
 46 $\mathbf{a}(\theta, \cdot)$ and $\mathbf{b}(\theta, \cdot)$ as in (3.67) with a collision kernel $m_{1/\sqrt{\theta}} b$. 45

1 *Order ε^1 .* One finds that

$$2 \quad \partial_t F_1 + v \cdot \nabla_x F_1 - \mathcal{B}(F_1, F_1) = 2\mathcal{B}(F_0, F_2)$$

4 which can be put in the form

$$5 \quad \mathcal{L}_{F_0} \left(\frac{F_2}{F_0} \right) = \mathcal{Q}_{F_0} \left(\frac{F_1}{F_0}, \frac{F_1}{F_0} \right) - \frac{1}{F_0} (\partial_t + v \cdot \nabla_x) F_1$$

7 and this is therefore a Fredholm integral equation of the type (3.63). Here \mathcal{Q}_{F_0} is the Boltzmann collision integral intertwined with the multiplication by the local Maxwellian F_0 , i.e.,

$$10 \quad \mathcal{Q}_{F_0}(\phi, \phi) = F_0^{-1} \mathcal{B}(F_0 \phi, F_0 \phi).$$

12 For this equation to have a solution, one must verify the compatibility conditions

$$14 \quad \partial_t \int_{\mathbb{R}^3} \begin{pmatrix} 1 \\ v \\ \frac{1}{2}|v|^2 \end{pmatrix} F_1 \, dv + \operatorname{div}_x \int_{\mathbb{R}^3} v \otimes \begin{pmatrix} 1 \\ v \\ \frac{1}{2}|v|^2 \end{pmatrix} F_1 \, dv = 0.$$

16 These five compatibility conditions are five PDEs for the five unknown functions ρ_1 , u_1 and θ_1 .

18 *Order ε^n .* One finds

$$20 \quad \partial_t F_n + v \cdot \nabla_x F_n - \sum_{k+l=n, 1 \leq k, l \leq n} \mathcal{B}(F_k, F_l) = 2\mathcal{B}(F_0, F_{n+1})$$

22 which is a Fredholm equation of the same type as above.

23 Here again, the compatibility condition reduces to

$$25 \quad \partial_t \int_{\mathbb{R}^3} \begin{pmatrix} 1 \\ v \\ \frac{1}{2}|v|^2 \end{pmatrix} F_n \, dv + \operatorname{div}_x \int_{\mathbb{R}^3} v \otimes \begin{pmatrix} 1 \\ v \\ \frac{1}{2}|v|^2 \end{pmatrix} F_n \, dv = 0.$$

27 More generally, the compatibility condition at order $n + 1$ (to guarantee the existence of F_{n+1}) provides the system of five equations satisfied by that part of F_n which belongs to the nullspace of \mathcal{L}_{F_0} . See [115] or Chapter V, Section 2 in [27] for more on Hilbert's expansion.

29 *Conclusion.* The Hilbert expansion method shows that the leading order in ε of Hilbert's formal solution (5.3) of the scaled Boltzmann equation (5.1) with Strouhal number $St = 1$ and Knudsen number $Kn = \varepsilon \ll 1$ is a local Maxwellian state whose parameters are governed by the Euler equations of gas dynamics (for a perfect monatomic gas).

In general, one cannot hope that Hilbert's formal power series (5.3) has a positive radius of convergence. Yet, Hilbert's expansion method can be used to obtain a rigorous derivation of the Euler equations of gas dynamics from the Boltzmann equation, as we shall see in Section 9.

5.2. The compressible Navier–Stokes limit: The Chapman–Enskog expansion

In this subsection, we shall seek higher-order (in ε) corrections to the compressible Euler system. The Hilbert expansion presented above is not well suited for this purpose, because linear combinations of collision invariants (i.e., hydrodynamic modes) appear at each order in ε instead of being all concentrated in the leading order term.

For that reason, we shall use a slightly different expansion method, the Chapman–Enskog expansion. Thus, we seek a solution of the scaled Boltzmann equation (5.1) as a Chapman–Enskog formal power series,

$$F_\varepsilon(t, x, v) = \sum_{n \geq 0} \varepsilon^n F^{(n)}[\vec{P}(t, x)](v), \quad (5.7)$$

parametrized by the vector \vec{P} of conserved densities of F_ε .

NOTATION. $F^n[\vec{P}(t, x)](v)$ designates a quantity that depends smoothly on \vec{P} and any finite number of its derivatives with respect to the x -variable at the same point (t, x) , and on the v -variable.

In particular, $F^n[\vec{P}(t, x)](v)$ does not contain time-derivatives of \vec{P} : the Chapman–Enskog method is based on eliminating $\partial_t \vec{P}$ in favor of x -derivatives via conservation laws satisfied by \vec{P} .

That \vec{P} is the vector of conserved densities of F_ε means that

$$\int_{\mathbb{R}^3} F^{(0)}[\vec{P}](v) \begin{pmatrix} 1 \\ v \\ \frac{1}{2}|v|^2 \end{pmatrix} dv = \vec{P}, \quad (5.8)$$

$$\int_{\mathbb{R}^3} F^{(n)}[\vec{P}](v) \begin{pmatrix} 1 \\ v \\ \frac{1}{2}|v|^2 \end{pmatrix} dv = \vec{0}, \quad n \geq 1.$$

These conserved densities satisfy a formal system of conservation laws of the form

$$\partial_t \vec{P} = \sum_{n \geq 0} \varepsilon^n \operatorname{div}_x \Phi^{(n)}[\vec{P}], \quad (5.9)$$

where the formal fluxes $\Phi^{(n)}$ are obtained from the local conservation laws associated to the Boltzmann equation by the formulas

$$\Phi^{(n)}[\vec{P}] = - \int_{\mathbb{R}^3} v \otimes \begin{pmatrix} 1 \\ v \\ \frac{1}{2}|v|^2 \end{pmatrix} F^{(n)}[\vec{P}](v) dv \quad (5.10)$$

for $n \geq 0$.

Let us analyze the first orders in ε of the Chapman–Enskog expansion.

Order 0. One has

$$\mathcal{B}(F^{(0)}[\vec{P}], F^{(0)}[\vec{P}]) = 0 \quad \text{and thus} \quad F^{(0)}[\vec{P}] = \mathcal{M}_{(\rho, u, \theta)},$$

here

$$\vec{P} = \begin{pmatrix} \rho \\ \rho u \\ \rho(\frac{1}{2}|u|^2 + \frac{3}{2}\theta) \end{pmatrix}, \quad \Phi^{(0)}[\vec{P}] = - \begin{pmatrix} \rho u \\ \rho u^{\otimes 2} + \rho \theta I \\ \rho u(\frac{1}{2}|u|^2 + \frac{5}{2}\theta) \end{pmatrix}.$$

Hence the formal conservation law at order 0 is

$$\partial_t \vec{P} = \operatorname{div}_x \Phi^{(0)}[\vec{P}] \quad \text{mod } \mathcal{O}(\varepsilon)$$

whose explicit form is

$$\begin{aligned} \partial_t \rho + u \cdot \nabla_x \rho + \rho \operatorname{div}_x u &= 0, \\ \partial_t u + (u \cdot \nabla_x) u + \frac{1}{\rho} \nabla_x(\rho \theta) &= 0 \quad \text{mod } \mathcal{O}(\varepsilon), \\ \partial_t \theta + u \cdot \nabla_x \theta + \frac{2}{3} \theta \operatorname{div}_x u &= 0. \end{aligned} \quad (5.11)$$

In other words, the 0th order of the Chapman–Enskog expansion gives the compressible Euler system, as does the Hilbert expansion.

The Hilbert and Chapman–Enskog methods differ at order 1 in ε , as we shall see below.

Order 1. One has

$$(\partial_t + v \cdot \nabla_x) F^{(0)}[\vec{P}] = 2\mathcal{B}(F^{(0)}[\vec{P}], F^{(1)}[\vec{P}]). \quad (5.12)$$

Using the formal conservation law at order 0, we eliminate $\partial_t F^{(0)}[\vec{P}]$ and replace it with x -derivatives of $F^{(0)}[\vec{P}]$, by using (5.5) and the conservation laws at order 0 (5.11) (i.e., Euler’s system) as follows

$$\begin{aligned} (\partial_t + v \cdot \nabla_x) \mathcal{M}_{(\rho, u, \theta)} \\ = \mathcal{M}_{(\rho, u, \theta)} (A(V) : D(u) + 2B(V) \cdot \nabla_x \sqrt{\theta}) + \mathcal{O}(\varepsilon) \end{aligned} \quad (5.13)$$

with the notation

$$V = \frac{v-u}{\sqrt{\theta}}, \quad A(V) = V \otimes V - \frac{1}{3}|v|^2 I, \quad B(V) = \frac{1}{2}V(|V|^2 - 5),$$

and where $D(u)$ is the traceless part of the deformation tensor of u

$$D(u) = \frac{1}{2} \left(\nabla_x u + (\nabla_x u)^T - \frac{2}{3} \operatorname{div}_x u I \right).$$

In view of (5.12) and (5.13), $F^{(1)}[\vec{P}]$ is determined by the conditions

$$A(V) : D(u) + 2B(V) \cdot \nabla_x \sqrt{\theta} = -\mathcal{L}_{\mathcal{M}(\rho, u, \theta)} \left(\frac{F^{(1)}[\vec{P}]}{\mathcal{M}(\rho, u, \theta)} \right),$$

$$\int F^{(1)}[\vec{P}](v) \begin{pmatrix} 1 \\ v \\ \frac{1}{2}|v|^2 \end{pmatrix} dv = 0.$$

By Hilbert's lemma, $\mathcal{L}_{\mathcal{M}(1, u, \theta)}$ is a Fredholm operator on $L^2(M dv)$; thus

$$F^{(1)}[\vec{P}](v) = -\mathcal{M}(1, u, \theta) \left(\mathbf{a}(\theta, |V|) A(V) : D(u) + 2\mathbf{b}(\theta, |V|) B(V) \cdot \nabla_x \sqrt{\theta} \right),$$

where we recall that the scalar functions \mathbf{a} and \mathbf{b} are defined by

$$\mathcal{L}_{\mathcal{M}(1, u, \theta)} (\mathbf{a}(\theta, |V|) A(V)) = A(V) \quad \text{and} \quad \mathbf{a}(\theta, |V|) A(V) \perp \operatorname{Ker} \mathcal{L}_{\mathcal{M}(1, u, \theta)}$$

while

$$\mathcal{L}_{\mathcal{M}(1, u, \theta)} (\mathbf{b}(\theta, |V|) B(V)) = B(V) \quad \text{and} \quad \mathbf{b}(\theta, |V|) B(V) \perp \operatorname{Ker} \mathcal{L}_{\mathcal{M}(1, u, \theta)}.$$

Hence the first-order correction to the fluxes in the formal conservation law is

$$\Phi^{(1)}[\vec{P}] = \begin{pmatrix} 0 \\ \mu(\theta) D(u) \\ \mu(\theta) D(u) \cdot u + \kappa(\theta) \nabla_x \theta \end{pmatrix}.$$

Therefore, the formal conservation law at first order is

$$\partial_t \vec{P} = \operatorname{div}_x \Phi^{(0)}[\vec{P}] + \varepsilon \operatorname{div}_x \Phi^{(1)}[\vec{P}] \quad \text{mod } \mathcal{O}(\varepsilon^2),$$

i.e., the compressible Navier–Stokes system with $O(\varepsilon)$ dissipation terms

$$\begin{aligned} \partial_t \rho + \operatorname{div}_x(\rho u) &= 0, \\ \partial_t(\rho u) + \operatorname{div}_x(\rho u \otimes u) + \nabla_x(\rho \theta) &= \varepsilon \operatorname{div}_x(\mu D(u)) \pmod{O(\varepsilon^2)}, \\ \partial_t \left(\rho \left(\frac{1}{2}|u|^2 + \frac{3}{2}\theta \right) \right) + \operatorname{div}_x \left(\rho u \left(\frac{1}{2}|u|^2 + \frac{5}{2}\theta \right) \right) & \\ &= \varepsilon \operatorname{div}_x(\kappa \nabla_x \theta) + \varepsilon \operatorname{div}_x(\mu D(u) \cdot u). \end{aligned} \quad (5.14)$$

The viscosity μ and heat conduction κ are computed as follows

$$\begin{aligned} \theta \int_{\mathbb{R}^3} \mathbf{a}(\theta, |V|) A_{ij}(V) A_{kl}(V) \mathcal{M}_{(1,u,\theta)} dv &= \mu(\theta) \left(\delta_{ik} \delta_{jl} + \delta_{il} \delta_{jk} - \frac{2}{3} \delta_{ij} \delta_{kl} \right), \\ \theta \int_{\mathbb{R}^3} \mathbf{b}(\theta, |V|) B_i(V) B_j(V) \mathcal{M}_{(1,u,\theta)} dv &= \kappa(\theta) \delta_{ij}, \end{aligned}$$

or in other words,

$$\begin{aligned} \mu(\theta) &= \frac{2}{15} \theta \int_0^{+\infty} \mathbf{a}(\theta, r) r^6 e^{-r^2/2} \frac{dr}{\sqrt{2\pi}}, \\ \kappa(\theta) &= \frac{1}{6} \theta \int_0^{+\infty} \mathbf{b}(\theta, r) r^4 (r^2 - 5)^2 e^{-r^2/2} \frac{dr}{\sqrt{2\pi}}. \end{aligned} \quad (5.15)$$

In the hard sphere case, one finds that the viscosity and heat conduction are of the form

$$\mu(\theta) = \mu_0 \sqrt{\theta}, \quad \kappa(\theta) = \kappa_0 \sqrt{\theta},$$

where μ_0 and ν_0 are positive constants.

Conclusion. The Chapman–Enskog expansion method shows that the first-order (in ε) correction to the compressible Euler system in the limit of (5.1) – with $St = 1$ and $Kn = \varepsilon \rightarrow 0$ – is the compressible Navier–Stokes system (5.14).

Notice that the class of Navier–Stokes systems obtained in this way is by no means the most general: only the equation of state of a perfect monatomic gas can be obtained in this way, as in the case of the compressible Euler limit of the Boltzmann equation. In addition, the viscous dissipation tensor obtained in this way involves *only one* viscosity coefficient instead of two.

Finally, the Chapman–Enskog expansion can be pushed further to obtain higher-order corrections to the compressible Euler equations. For instance, the second-order correction to the compressible Euler equations is a system known as *the Burnett equations*; further corrections have also been computed and are known as *the super-Burnett equations*. See Chapter 5, Section 3 in [27] and Section 25 in [63] for more material on the Chapman–Enskog expansion, as well as for a comparison with Hilbert’s expansion.

It should be noted however that these further corrections to the Euler system, beyond the Navier–Stokes system, i.e., the Burnett and super-Burnett equations, are in general not well posed, so that their practical interest in fluid mechanics is unclear. Therefore, we shall not pursue this line of investigation. However, Levermore recently proposed a subtle modification of the Chapman–Enskog expansion which leads to well-posed variants of the Burnett systems [82].

5.3. The compressible Euler limit: The moment method

We shall now present a method for deriving hydrodynamic equations from the Boltzmann equation that differs from either the Hilbert or Chapman–Enskog expansions. It consists of passing to the limit as the Knudsen number vanishes in the local conservation laws of mass, momentum and energy that are satisfied by “well-behaved” solutions of the Boltzmann equation. We describe this method on the derivation of the compressible Euler equations from the scaled Boltzmann equation (5.1).

Start from the Cauchy problem for the Boltzmann equation in the periodic box

$$\begin{aligned} \partial_t F_\varepsilon + v \cdot \nabla_x F_\varepsilon &= \frac{1}{\varepsilon} \mathcal{B}(F_\varepsilon, F_\varepsilon), \quad (t, x, v) \in \mathbb{R}_+^* \times \mathbb{T}^3 \times \mathbb{R}^3, \\ F_\varepsilon|_{t=0} &= \mathcal{M}_{(\rho^{\text{in}}, u^{\text{in}}, \theta^{\text{in}})}. \end{aligned} \tag{5.16}$$

As before, the collision kernel b in the Boltzmann collision integral satisfies the cut-off assumption (5.2).

THEOREM 5.1. *Let $\rho^{\text{in}} \geq 0$ a.e., u^{in} and $\theta^{\text{in}} > 0$ a.e. be such that*

$$\int_{\mathbb{T}^3} \rho^{\text{in}} (1 + |u^{\text{in}}|) (|u^{\text{in}}|^2 + \theta^{\text{in}} + |\ln \rho^{\text{in}}| + |\ln \theta^{\text{in}}|) dx < +\infty.$$

For each $\varepsilon > 0$, let F_ε be a solution of (5.16) that satisfies the local conservation laws of mass, momentum, and energy, as well as the local entropy relation. Assume that

$$F_\varepsilon \rightarrow F \quad \text{a.e.}$$

as well as

$$\begin{aligned} \int_{\mathbb{R}^3} F_\varepsilon dv &\rightarrow \int_{\mathbb{R}^3} F dv && \text{in } C(\mathbb{R}_+; \mathcal{D}'(\mathbb{T}^3)), \\ \int_{\mathbb{R}^3} v F_\varepsilon dv &\rightarrow \int_{\mathbb{R}^3} v F dv && \text{in } C(\mathbb{R}_+; \mathcal{D}'(\mathbb{T}^3)), \\ \int_{\mathbb{R}^3} |v|^2 F_\varepsilon dv &\rightarrow \int_{\mathbb{R}^3} |v|^2 F dv && \text{in } C(\mathbb{R}_+; \mathcal{D}'(\mathbb{T}^3)), \end{aligned}$$

1 while

$$2 \int_{\mathbb{R}^3} v \otimes v F_\varepsilon \, dv \rightarrow \int_{\mathbb{R}^3} v \otimes v F \, dv \quad \text{in } \mathcal{D}'(\mathbb{R}_+^* \times \mathbb{T}^3),$$

$$3 \int_{\mathbb{R}^3} v |v|^2 F_\varepsilon \, dv \rightarrow \int_{\mathbb{R}^3} v |v|^2 F \, dv \quad \text{in } \mathcal{D}'(\mathbb{R}_+^* \times \mathbb{T}^3),$$

$$4 \int_{\mathbb{R}^3} F_\varepsilon \ln F_\varepsilon \, dv \rightarrow \int_{\mathbb{R}^3} F \ln F \, dv \quad \text{in } \mathcal{D}'(\mathbb{R}_+^* \times \mathbb{T}^3),$$

$$5 \int_{\mathbb{R}^3} v F_\varepsilon \ln F_\varepsilon \, dv \rightarrow \int_{\mathbb{R}^3} v F \ln F \, dv \quad \text{in } \mathcal{D}'(\mathbb{R}_+^* \times \mathbb{T}^3),$$

6 as $\varepsilon \rightarrow 0$. Then

$$7 F = \mathcal{M}_{(\rho, u, \theta)},$$

8 where (ρ, u, θ) is an entropic solution of the compressible Euler system

$$9 \begin{aligned} 10 \partial_t \rho + \operatorname{div}_x(\rho u) &= 0, \\ 11 \partial_t(\rho u) + \operatorname{div}_x(\rho u \otimes u) + \nabla_x(\rho \theta) &= 0, \\ 12 \partial_t \left(\rho \left(\frac{1}{2} |u|^2 + \frac{3}{2} \theta \right) \right) + \operatorname{div}_x \left(\rho u \left(\frac{1}{2} |u|^2 + \frac{5}{2} \theta \right) \right) &= 0, \end{aligned} \quad (5.17)$$

13 that satisfies the initial condition

$$14 (\rho, u, \theta)|_{t=0} = (\rho^{\text{in}}, u^{\text{in}}, \theta^{\text{in}}). \quad (5.18)$$

15 PROOF. The moment method involves three steps.

16 *Step 1* (Entropy production bound implies convergence to local equilibrium). The entropy relation in the 3-torus implies the entropy production bound

$$17 \begin{aligned} 18 \frac{1}{4} \int_0^t \int_{\mathbb{T}^3} \int \int \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F'_\varepsilon F'_{\varepsilon*} - F_\varepsilon F_{\varepsilon*}) \ln \left(\frac{F'_\varepsilon F'_{\varepsilon*}}{F_\varepsilon F_{\varepsilon*}} \right) b \, dv \, dv_* \, d\omega \\ 19 \leq \varepsilon H(\mathcal{M}_{(\rho^{\text{in}}, u^{\text{in}}, \theta^{\text{in}})} | \bar{\mathcal{M}}), \end{aligned} \quad (5.19)$$

20 where $\bar{\mathcal{M}}$ is any global Maxwellian state, for instance, one could choose

$$21 \bar{\mathcal{M}} = \mathcal{M}_{(\bar{\rho}^{\text{in}}, \bar{u}^{\text{in}}, \bar{\theta}^{\text{in}})},$$

22 where

$$23 \bar{\rho}^{\text{in}} = \int_{\mathbb{T}^3} \rho^{\text{in}}(x) \, dx,$$

$$\begin{aligned} \overline{u^{\text{in}}} &= \frac{1}{\rho^{\text{in}}} \int_{\mathbb{T}^3} \rho^{\text{in}} u^{\text{in}}(x) \, dx, \\ \overline{\theta^{\text{in}}} &= \frac{1}{\rho^{\text{in}}} \int_{\mathbb{T}^3} \rho^{\text{in}} \left(\frac{1}{3} |u^{\text{in}} - \overline{u^{\text{in}}}|^2 + \theta^{\text{in}} \right) (x) \, dx. \end{aligned}$$

Next we apply Fatou's lemma: assuming that $F_\varepsilon \rightarrow F$ a.e. on $\mathbb{R}_+ \times \mathbb{T}^3 \times \mathbb{R}^3$, one has

$$\begin{aligned} 0 &\leq \int_0^t \int_{\mathbb{T}^3} \iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F' F'_* - F F_*) \ln \left(\frac{F' F'_*}{F F_*} \right) b \, dv_* \, d\omega \, dx \, ds \\ &\leq \liminf_{\varepsilon \rightarrow 0} \int_0^t \int_{\mathbb{T}^3} \iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F'_\varepsilon F'_{\varepsilon*} - F_\varepsilon F_{\varepsilon*}) \ln \left(\frac{F'_\varepsilon F'_{\varepsilon*}}{F_\varepsilon F_{\varepsilon*}} \right) b \, dv_* \, d\omega \, dx \, ds \\ &= 0. \end{aligned}$$

Hence F is a local Maxwellian, i.e.,

$$F(t, x, v) = \mathcal{M}_{(\rho(t,x), u(t,x), \theta(t,x))}(v)$$

for some $\rho(t, x) \geq 0$, $\theta(t, x) > 0$ and $u(t, x) \in \mathbb{R}^3$.

Step 2 (Passing to the limit in the local conservation laws). For each positive ε , the number density F_ε satisfies the local conservation laws recalled below

$$\begin{aligned} \partial_t \int_{\mathbb{R}^3} F_\varepsilon \, dv + \operatorname{div}_x \int_{\mathbb{R}^3} v F_\varepsilon \, dv &= 0, \\ \partial_t \int_{\mathbb{R}^3} v F_\varepsilon \, dv + \operatorname{div}_x \int_{\mathbb{R}^3} v \otimes v F_\varepsilon \, dv &= 0, \\ \partial_t \int_{\mathbb{R}^3} \frac{1}{2} |v|^2 F_\varepsilon \, dv + \operatorname{div}_x \int_{\mathbb{R}^3} v \frac{1}{2} |v|^2 F_\varepsilon \, dv &= 0. \end{aligned} \tag{5.20}$$

It follows from our assumptions and Step 1 that

$$\begin{aligned} \int_{\mathbb{R}^3} F_\varepsilon \, dv &\rightarrow \int_{\mathbb{R}^3} \mathcal{M}_{(\rho, u, \theta)} \, dv = \rho, \\ \int_{\mathbb{R}^3} v F_\varepsilon \, dv &\rightarrow \int_{\mathbb{R}^3} v \mathcal{M}_{(\rho, u, \theta)} \, dv = \rho u, \\ \int_{\mathbb{R}^3} v \otimes v F_\varepsilon \, dv &\rightarrow \int_{\mathbb{R}^3} v \otimes v \mathcal{M}_{(\rho, u, \theta)} \, dv = \rho(u \otimes u + \theta I), \\ \int_{\mathbb{R}^3} v |v|^2 F_\varepsilon \, dv &\rightarrow \int_{\mathbb{R}^3} v |v|^2 \mathcal{M}_{(\rho, u, \theta)} \, dv = \rho u \left(\frac{1}{2} |u|^2 + \frac{5}{2} \theta \right), \end{aligned}$$

1 in the limit as $\varepsilon \rightarrow 0$. Hence the functions ρ and θ , and the vector field u satisfy the system 1
 2 of PDEs (5.17). It also satisfies the initial condition (5.18) because the convergence of the 2
 3 conserved densities is locally uniform in t . 3
 4

5 *Step 3 (Passing to the limit in the local entropy relation).* Finally, we recall the local 5
 6 entropy relation 6
 7

$$8 \quad \partial_t \int_{\mathbb{R}^3} F_\varepsilon \ln F_\varepsilon \, dv + \operatorname{div}_x \int_{\mathbb{R}^3} v F_\varepsilon \ln F_\varepsilon \, dv = -\text{local entropy production rate} \leq 0. \quad 8$$

10 It follows from our assumptions that 10
 11

$$12 \quad \int_{\mathbb{R}^3} F_\varepsilon \ln F_\varepsilon \, dv \rightarrow \int_{\mathbb{R}^3} \mathcal{M}_{(\rho,u,\theta)} \ln \mathcal{M}_{(\rho,u,\theta)} \, dv \quad 12$$

$$13 \quad = \rho \ln \left(\frac{\rho}{\theta^{3/2}} \right) - \frac{3}{2} (1 + \ln(2\pi)) \rho, \quad 13$$

$$14 \quad \int_{\mathbb{R}^3} v F_\varepsilon \ln F_\varepsilon \, dv \rightarrow \int_{\mathbb{R}^3} v \mathcal{M}_{(\rho,u,\theta)} \ln \mathcal{M}_{(\rho,u,\theta)} \, dv \quad 14$$

$$15 \quad = \rho u \ln \left(\frac{\rho}{\theta^{3/2}} \right) - \frac{3}{2} (1 + \ln(2\pi)) \rho u, \quad 15$$

16 so that, by passing to the limit in the local entropy relation, on account of the continuity 16
 17 equation in (5.17), one arrives at the differential inequality 17
 18

$$19 \quad \partial_t \left(\rho \ln \left(\frac{\rho}{\theta^{3/2}} \right) \right) + \operatorname{div}_x \left(\rho u \ln \left(\frac{\rho}{\theta^{3/2}} \right) \right) \leq 0. \quad 19$$

20 In other words, (ρ, u, θ) is a solution of the compressible Euler equations that satisfies the 20
 21 Lax–Friedrichs entropy condition. \square 21
 22

23 Theorem 5.1 and its proof are taken from [7]. 23
 24

25 **5.4. The acoustic limit** 25

26 Start from the Boltzmann equation with the same scaling as before, but with initial data 26
 27 that are small perturbations of a uniform Maxwellian equilibrium. By Galilean invariance, 27
 28 one can assume without loss of generality that this uniform equilibrium is the centered 28
 29 reduced Gaussian 29
 30

$$31 \quad M(v) = \mathcal{M}_{(1,0,1)}(v) = \frac{1}{(2\pi)^{3/2}} e^{-|v|^2/2}. \quad 31$$

1 Thus, the problem (5.16) reduces to

$$\begin{aligned}
 2 & \\
 3 & \partial_t F_\varepsilon + v \cdot \nabla_x F_\varepsilon = \frac{1}{\varepsilon} \mathcal{B}(F_\varepsilon, F_\varepsilon), \quad (t, x, v) \in \mathbb{R}_+^* \times \mathbb{T}^3 \times \mathbb{R}^3, \\
 4 & \\
 5 & F_\varepsilon|_{t=0} = \mathcal{M}_{(1+\eta_\varepsilon \rho^{\text{in}}, \eta_\varepsilon u^{\text{in}}, 1+\eta_\varepsilon \theta^{\text{in}})}, \\
 6 &
 \end{aligned} \tag{5.22}$$

7 where $0 < \varepsilon \ll 1$ and $0 < \eta_\varepsilon \ll 1$. The same moment method as above shows that the
 8 limiting behavior of the solution to (5.22) under these assumptions is governed by the
 9 acoustic system.

10
 11
 12 **THEOREM 5.2.** *Assume that $\rho^{\text{in}}, u^{\text{in}}$ and θ^{in} belong to $L^2(\mathbb{T}^3)$ and that $\eta_\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0$.
 13 For each $\varepsilon > 0$, let F_ε be a solution of (5.22) that satisfies the local conservation laws of
 14 mass momentum and energy, as well as the local entropy relation.*

15 *Assume that*

$$16 \quad g_\varepsilon = \frac{F_\varepsilon - M}{\eta_\varepsilon M} \rightarrow g \quad \text{in the sense of distributions,}$$

17
 18
 19
 20 while

$$21 \quad \eta_\varepsilon \mathcal{B}(F_\varepsilon - M, F_\varepsilon - M) \rightarrow 0 \quad \text{in the sense of distributions,}$$

22
 23
 24 as well as

$$25 \quad \int_{\mathbb{R}^3} g_\varepsilon M \, dv \rightarrow \int_{\mathbb{R}^3} g M \, dv \quad \text{in } C(\mathbb{R}_+; \mathcal{D}'(\mathbb{T}^3)),$$

$$26 \quad \int_{\mathbb{R}^3} v g_\varepsilon M \, dv \rightarrow \int_{\mathbb{R}^3} v g M \, dv \quad \text{in } C(\mathbb{R}_+; \mathcal{D}'(\mathbb{T}^3)),$$

$$27 \quad \int_{\mathbb{R}^3} |v|^2 g_\varepsilon M \, dv \rightarrow \int_{\mathbb{R}^3} |v|^2 g M \, dv \quad \text{in } C(\mathbb{R}_+; \mathcal{D}'(\mathbb{T}^3)),$$

28
 29
 30
 31
 32
 33
 34 and

$$35 \quad \int_{\mathbb{R}^3} v \otimes v g_\varepsilon M \, dv \rightarrow \int_{\mathbb{R}^3} v \otimes v g M \, dv \quad \text{in } \mathcal{D}'(\mathbb{R}_+^* \times \mathbb{T}^3),$$

$$36 \quad \int_{\mathbb{R}^3} v |v|^2 g_\varepsilon M \, dv \rightarrow \int_{\mathbb{R}^3} v |v|^2 g M \, dv \quad \text{in } \mathcal{D}'(\mathbb{R}_+^* \times \mathbb{T}^3)$$

37
 38
 39
 40
 41
 42 as $\varepsilon \rightarrow 0$. Then

$$43 \quad g(t, x, v) = \rho(t, x) + u(t, x) \cdot v + \theta(t, x) \frac{1}{2} (|v|^2 - 3),$$

44

45

1 where (ρ, u, θ) is the solution of the acoustic system

$$\begin{aligned}
 2 & \partial_t \rho + \operatorname{div}_x u = 0, \\
 3 & \partial_t u + \nabla_x (\rho + \theta) = 0, \\
 4 & \frac{3}{2} \partial_t \theta + \operatorname{div}_x u = 0,
 \end{aligned}
 \tag{5.23}$$

8 that satisfies the initial condition

$$10 \quad (\rho, u, \theta)|_{t=0} = (\rho^{\text{in}}, u^{\text{in}}, \theta^{\text{in}}).
 \tag{5.24}$$

13 The proof of this theorem is an easy variant of the formal compressible Euler limit, and
 14 is left to the reader. See [53] for the missing details of this formal proof.

16 6. Incompressible limits of the Boltzmann equation: Formal results

18 So far, we have discussed various limits of the kinetic theory of gases leading to hydrodynamic models for compressible fluids that satisfy the equation of state of perfect gases. As
 19 we shall see in this section, incompressible hydrodynamic models describing *incompressible flows* of perfect gases can also be derived from the Boltzmann equation.

24 6.1. The incompressible Navier–Stokes limit

26 The scaling on the Boltzmann equation that leads to the incompressible Navier–Stokes
 27 equations in the hydrodynamic limit is defined by

$$29 \quad Kn = St = \varepsilon \ll 1.$$

31 However, this scaling is not sufficient by itself: as in all long time scalings, one should
 32 assume that the length and time scales L and T that enter the definition of St capture the
 33 speed of the fluid motion. In other words, situations where

$$35 \quad |u(t, x)| \gg \frac{L}{T} \quad \text{must be excluded,}$$

38 where u is the bulk velocity of the gas, i.e.,

$$40 \quad u(t, x) = \frac{\int_{\mathbb{R}^3} v F \, dv}{\int_{\mathbb{R}^3} F \, dv}.$$

43 In other words, since $St = (L/T)/\text{speed of sound}$, one must take

$$45 \quad Ma = O(St).$$

The case $Ma \sim St$ corresponds to the largest possible velocity field compatible with the above condition, and therefore leads to the Navier–Stokes equation, while the case $Ma = o(\varepsilon)$ leads to the linearized version of the Navier–Stokes equations, i.e., the Stokes equations.

Hence the complete Navier–Stokes scaling is

$$Kn = St = Ma = \varepsilon \ll 1. \quad (6.1)$$

We consider therefore the scaled Boltzmann equation posed on the spatial domain \mathbb{R}^3 with uniform Maxwellian equilibrium at infinity – without loss of generality, this Maxwellian equilibrium is assumed to be the centered reduced Gaussian distribution

$$M(v) = \mathcal{M}_{(1,0,1)}(v) = \frac{1}{(2\pi)^{3/2}} e^{-|v|^2/2}.$$

The problem to be studied is therefore

$$\begin{aligned} \varepsilon \partial_t F_\varepsilon + v \cdot \nabla_x F_\varepsilon &= \frac{1}{\varepsilon} \mathcal{B}(F_\varepsilon, F_\varepsilon), \quad (t, x, v) \in \mathbb{R}_+^* \times \mathbb{R}^3 \times \mathbb{R}^3, \\ F_\varepsilon(t, x, v) &\rightarrow M \quad \text{as } |x| \rightarrow +\infty. \end{aligned} \quad (6.2)$$

That $Ma = O(\varepsilon)$ is seen on the number density F_ε , and not on the Boltzmann equation itself. Here is an example of number density with $Ma = O(\varepsilon)$.

EXAMPLE 1. Take F_ε of the form

$$F_\varepsilon(t, x, v) = \mathcal{M}_{(1, \varepsilon u(t, x), 1)}(v).$$

Indeed,

$$\frac{\int_{\mathbb{R}^3} v F_\varepsilon \, dv}{\int_{\mathbb{R}^3} F_\varepsilon \, dv} := \varepsilon u.$$

The speed of sound for the state of the gas described by F_ε is $\sqrt{\frac{5}{3}}\theta$ where

$$\theta(t, x) := \frac{\int_{\mathbb{R}^3} (1/3) |v - \varepsilon u|^2 F_\varepsilon \, dv}{\int_{\mathbb{R}^3} F_\varepsilon \, dv} = 1.$$

Hence the Mach number for the state of the gas associated to F_ε is

$$\frac{\varepsilon |u(t, x)|}{\sqrt{(5/3)\theta(t, x)}} = O(\varepsilon).$$

Here is another example that also involves fluctuations of temperature.

1 EXAMPLE 2. Take F_ε of the form

$$2 \quad 3 \quad 4 \quad 5 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \quad 11 \quad 12 \quad 13 \quad 14 \quad 15 \quad 16 \quad 17 \quad 18 \quad 19 \quad 20 \quad 21 \quad 22 \quad 23 \quad 24 \quad 25 \quad 26 \quad 27 \quad 28 \quad 29 \quad 30 \quad 31 \quad 32 \quad 33 \quad 34 \quad 35 \quad 36 \quad 37 \quad 38 \quad 39 \quad 40 \quad 41 \quad 42 \quad 43 \quad 44 \quad 45$$

$$F_\varepsilon(t, x, v) = \mathcal{M}_{(1-\varepsilon\theta(t,x), \frac{\varepsilon u(t,x)}{1-\varepsilon\theta(t,x)}, \frac{1}{1-\varepsilon\theta(t,x)})}(v)$$

with $\theta \in L^\infty(\mathbb{R}_+ \times \mathbb{R}^3)$ and $\varepsilon\|\theta\|_{L^\infty} < 1$. One easily checks that

$$\frac{\int_{\mathbb{R}^3} v F_\varepsilon \, dv}{\int_{\mathbb{R}^3} F_\varepsilon \, dv} = \varepsilon u + O(\varepsilon^2),$$

while

$$\frac{\int_{\mathbb{R}^3} (1/3)|v - \varepsilon u(t, x)/(1 - \varepsilon\theta(t, x))|^2 F_\varepsilon \, dv}{\int_{\mathbb{R}^3} F_\varepsilon \, dv} = 1 + \varepsilon\theta + O(\varepsilon^2).$$

Hence the Mach number is

$$\frac{\varepsilon u + O(\varepsilon^2)}{\sqrt{(5/3)(1 + \varepsilon\theta + O(\varepsilon^2))}} = O(\varepsilon).$$

More generally, if F_ε is a number density of the form

$$F_\varepsilon = M(1 + \varepsilon g_\varepsilon) \quad \text{such that} \quad g_\varepsilon \geq -\frac{1}{\varepsilon} \quad \text{a.e.}, \quad (6.3)$$

one can check that, provided that $\|g_\varepsilon\|_{L^\infty} = O(1)$, the Mach number for the state of the gas defined by F_ε is $O(\varepsilon)$.

Hence, we shall supplement the scaled Boltzmann equation (6.2) with the initial condition

$$F_\varepsilon|_{t=0} \equiv \mathcal{M}_{(1-\varepsilon\theta^{\text{in}}(x), \frac{\varepsilon u^{\text{in}}(x)}{1-\varepsilon\theta^{\text{in}}(x)}, \frac{1}{1-\varepsilon\theta^{\text{in}}(x)})}(v), \quad (6.4)$$

where

$$\operatorname{div}_x u^{\text{in}} = 0.$$

For each $\varepsilon > 0$, define the number density fluctuation

$$g_\varepsilon = \frac{F_\varepsilon - M}{\varepsilon M}.$$

In terms of the number density fluctuation g_ε , the Boltzmann equation (6.2) takes the form

$$\varepsilon \partial_t g_\varepsilon + v \cdot \nabla_x g_\varepsilon + \frac{1}{\varepsilon} \mathcal{L}_M g_\varepsilon = \mathcal{Q}_M(g_\varepsilon, g_\varepsilon), \quad t > 0, x, v \in \mathbb{R}^3, \quad (6.5)$$

where \mathcal{Q}_M is the collision integral intertwined with the multiplication by M

$$\mathcal{Q}_M(\phi, \phi) = M^{-1} \mathcal{B}(M\phi, M\phi). \quad (6.6)$$

We shall also need notation for moments

$$\langle \phi \rangle = \int_{\mathbb{R}^3} \phi(v) M(v) dv.$$

Hence the local conservation laws of mass momentum and energy satisfied by F_ε take the form

$$\begin{aligned} \varepsilon \partial_t \langle g_\varepsilon \rangle + \operatorname{div}_x \langle v g_\varepsilon \rangle &= 0 && \text{(mass),} \\ \varepsilon \partial_t \langle v g_\varepsilon \rangle + \operatorname{div}_x \langle v \otimes v g_\varepsilon \rangle &= 0 && \text{(momentum),} \\ \varepsilon \partial_t \left\langle \frac{1}{2} |v|^2 g_\varepsilon \right\rangle + \operatorname{div}_x \left\langle v \frac{1}{2} |v|^2 g_\varepsilon \right\rangle &= 0 && \text{(energy).} \end{aligned} \quad (6.7)$$

THEOREM 6.1 (Bardos, Golse and Levermore [8,9]). *For each $\varepsilon > 0$, let F_ε be a solution of (6.2)–(6.4). Assume that*

$$\frac{F_\varepsilon - M}{\varepsilon M} \rightarrow g \quad \text{in the sense of distributions on } \mathbb{R}_+^* \times \mathbb{R}^3,$$

and that F_ε satisfies the local conservation laws of mass, momentum and energy, and that

$$\langle v g_\varepsilon \rangle \rightarrow \langle v g \rangle \quad \text{and} \quad \langle (|v|^2 - 5) g_\varepsilon \rangle \rightarrow \langle (|v|^2 - 5) g \rangle \quad \text{in } C(\mathbb{R}_+, \mathcal{D}'(\mathbb{R}^3))$$

while

$$\mathcal{L}_M g_\varepsilon \rightarrow \mathcal{L}_M g \quad \text{in } \mathcal{D}'(\mathbb{R}_+^* \times \mathbb{R}^3)$$

and all formally small terms vanish in the sense of distributions as $\varepsilon \rightarrow 0$. Assume further that

$$\begin{aligned} \langle v \otimes v g_\varepsilon \rangle &\rightarrow \langle v \otimes v g \rangle, & \langle \mathcal{B} g_\varepsilon \rangle &\rightarrow \langle \mathcal{B} g \rangle, \\ \langle \widehat{A} \mathcal{Q}_M(g_\varepsilon, g_\varepsilon) \rangle &\rightarrow \langle \widehat{A} \mathcal{Q}_M(g_\varepsilon, g_\varepsilon) \rangle & \text{and} & \langle \widehat{A} \otimes v g_\varepsilon \rangle \rightarrow \langle \widehat{A} \otimes v g \rangle, \\ \langle \widehat{B} \mathcal{Q}_M(g_\varepsilon, g_\varepsilon) \rangle &\rightarrow \langle \widehat{B} \mathcal{Q}_M(g_\varepsilon, g_\varepsilon) \rangle & \text{and} & \langle \widehat{B} \otimes v g_\varepsilon \rangle \rightarrow \langle \widehat{B} \otimes v g \rangle \end{aligned}$$

in the sense of distributions on $\mathbb{R}_+^* \times \mathbb{R}^3$.

Then g is of the form

$$g(t, x, v) = u(t, x) \cdot v + \theta(t, x) \frac{1}{2} (|v|^2 - 5),$$

where (u, θ) satisfy the incompressible Navier–Stokes–Fourier system

$$\begin{aligned} \partial_t u + \operatorname{div}_x(u \otimes u) + \nabla_x p &= \nu \Delta_x u, & \operatorname{div}_x u &= 0, \\ (\partial_t \theta + \operatorname{div}_x(u \theta)) &= \kappa \Delta_x \theta, \end{aligned} \quad (6.8)$$

where

$$\begin{aligned} \nu &= \frac{1}{10} \langle \widehat{A} : A \rangle = \frac{2}{15} \int_0^{+\infty} \mathbf{a}(1, r) r^6 e^{-r^2/2} \frac{dr}{\sqrt{2\pi}}, \\ \kappa &= \frac{2}{15} \langle \widehat{B} \cdot B \rangle = \frac{1}{15} \int_0^{+\infty} \mathbf{b}(1, r) r^2 (r^2 - 5) e^{-r^2/2} \frac{dr}{\sqrt{2\pi}}. \end{aligned} \quad (6.9)$$

We recall that

$$A(v) = v \otimes v - \frac{1}{3} |v|^2 I, \quad B(v) = \frac{1}{2} (|v|^2 - 5) v, \quad (6.10)$$

and that there exists \widehat{A} and $\widehat{B} \in L^2(\mathbb{R}^3, M \, dv)$ uniquely determined by

$$\begin{aligned} \mathcal{L}_M \widehat{A} &= A, & \widehat{A} &\perp \operatorname{Ker} \mathcal{L}_M, \\ \mathcal{L}_M \widehat{B} &= B, & \widehat{B} &\perp \operatorname{Ker} \mathcal{L}_M. \end{aligned} \quad (6.11)$$

Furthermore, there exists two scalar functions \mathbf{a} and \mathbf{b} such that

$$\widehat{A}(v) = \mathbf{a}(1, |v|) A(v) \quad \text{and} \quad \widehat{B}(v) = \mathbf{b}(1, |v|) B(v). \quad (6.12)$$

PROOF OF THEOREM 6.1. We shall use the moment method, although either the Hilbert or the Chapman–Enskog expansions would also lead to the incompressible Navier–Stokes limit. However, the moment method is the closest to a complete (instead of formal) convergence proof for global solutions without restriction on the size of the initial data.

Step 1 (Asymptotic form of the fluctuations). Multiply the Boltzmann equation (6.5) by ε , so that

$$\varepsilon^2 \partial_t g_\varepsilon + \varepsilon v \cdot \nabla_x g_\varepsilon + \mathcal{L}_M g_\varepsilon = \varepsilon \mathcal{Q}_M(g_\varepsilon, g_\varepsilon).$$

Passing to the limit as $\varepsilon \rightarrow 0$ in the above equation, we arrive at

$$\mathcal{L}_M g = 0.$$

Hence, for a.e. $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^3$, $g(t, x, \cdot) \in \operatorname{Ker} \mathcal{L}_M$, which means that g is of the form

$$g(t, x, v) = \rho(t, x) + u(t, x) \cdot v + \theta(t, x) \frac{1}{2} (|v|^2 - 3). \quad (6.13)$$

1 *Step 2* (Passing to the limit in the local conservation laws). Passing to the limit in (6.7) 1
 2 we arrive at 2

$$3 \quad \operatorname{div}_x \langle v g \rangle = 0, \quad 3$$

$$4 \quad \operatorname{div}_x \langle v \otimes v g \rangle = 0, \quad 4$$

$$5 \quad \operatorname{div}_x \langle v |v|^2 g \rangle = 0. \quad 5$$

6 Together with the relation (6.13), the first and the last relation reduce to the incompress- 6
 7 ibility condition for u 7

$$8 \quad \operatorname{div}_x u = 0. \quad (6.14) \quad 8$$

9 Together with the relation (6.13), the second relation gives 9

$$10 \quad \nabla_x (\rho + \theta) = 0. \quad 10$$

11 Since $g \in L^\infty(\mathbb{R}_+; L^2(\mathbb{R}^3, M dv))$, this implies the Boussinesq relation 11

$$12 \quad \rho + \theta = 0. \quad (6.15) \quad 12$$

13 With this last relation, the asymptotic form of g becomes 13

$$14 \quad g(t, x, v) = u(t, x) \cdot v + \theta(t, x) \frac{1}{2} (|v|^2 - 5). \quad (6.16) \quad 14$$

15 It remains to derive the motion and heat equations from the local conservation laws. To 15
 16 do this, we recast the local conservation law of momentum as 16

$$17 \quad \partial_t \langle v g_\varepsilon \rangle + \operatorname{div}_x \frac{1}{\varepsilon} \langle A g_\varepsilon \rangle + \nabla_x \frac{1}{\varepsilon} \left\langle \frac{1}{3} |v|^2 g_\varepsilon \right\rangle = 0, \quad 17$$

18 and we combine the local conservation laws of mass and energy into 18

$$19 \quad \partial_t \left\langle \frac{1}{2} (|v|^2 - 5) g_\varepsilon \right\rangle + \operatorname{div}_x \frac{1}{\varepsilon} \langle B g_\varepsilon \rangle = 0. \quad 19$$

20 One easily checks with (6.16) that 20

$$21 \quad \langle v g_\varepsilon \rangle \rightarrow \langle v g \rangle = u, \quad (6.17) \quad 21$$

$$22 \quad \left\langle \frac{1}{2} (|v|^2 - 5) g_\varepsilon \right\rangle \rightarrow \left\langle \frac{1}{2} (|v|^2 - 5) g \right\rangle = \frac{5}{2} \theta \quad 22$$

23 in $C(\mathbb{R}_+; \mathcal{D}'(\mathbb{R}^3))$ as $\varepsilon \rightarrow 0$. 23

Next we pass to the limit in the flux terms. Since \mathcal{L}_M is self-adjoint, one has

$$\frac{1}{\varepsilon} \langle A g_\varepsilon \rangle = \frac{1}{\varepsilon} \langle (\mathcal{L}_M \widehat{A}) g_\varepsilon \rangle = \left\langle \widehat{A} \frac{1}{\varepsilon} \mathcal{L}_M g_\varepsilon \right\rangle.$$

Then we eliminate the term $\frac{1}{\varepsilon} \mathcal{L}_M g_\varepsilon$ with (6.5)

$$\left\langle \widehat{A} \frac{1}{\varepsilon} \mathcal{L}_M g_\varepsilon \right\rangle = \langle \widehat{A} \mathcal{Q}_M(g_\varepsilon, g_\varepsilon) \rangle - \langle \widehat{A}(\varepsilon \partial_t + v \cdot \nabla_x) g_\varepsilon \rangle$$

so that, passing to the limit as $\varepsilon \rightarrow 0$ leads to

$$\frac{1}{\varepsilon} \langle A g_\varepsilon \rangle \rightarrow \langle \widehat{A} \mathcal{Q}_M(g, g) \rangle - \langle \widehat{A} v \cdot \nabla_x g \rangle \quad \text{in } \mathcal{D}'(\mathbb{R}_+^* \times \mathbb{R}^3). \quad (6.18)$$

Likewise

$$\frac{1}{\varepsilon} \langle B g_\varepsilon \rangle \rightarrow \langle \widehat{B} \mathcal{Q}_M(g, g) \rangle - \langle \widehat{B} v \cdot \nabla_x g \rangle \quad \text{in } \mathcal{D}'(\mathbb{R}_+^* \times \mathbb{R}^3), \quad (6.19)$$

as $\varepsilon \rightarrow 0$.

With (6.16), one easily finds that

$$\begin{aligned} \langle \widehat{A} v \cdot \nabla_x g \rangle &= v \left(\nabla_x u + (\nabla_x u)^T - \frac{2}{3} (\operatorname{div}_x u) I \right), \\ \langle \widehat{B} v \cdot \nabla_x g \rangle &= \frac{5}{2} \kappa \nabla_x \theta, \end{aligned} \quad (6.20)$$

where v and κ are given by (6.9).

The nonlinear term is slightly more difficult. Its computation involves in particular the following classical lemma.

LEMMA 6.2. *Let $\phi \in \operatorname{Ker} \mathcal{L}_M$, then*

$$\mathcal{Q}_M(\phi, \phi) = \frac{1}{2} \mathcal{L}_M(\phi^2).$$

PROOF. Differentiate twice the relation

$$\mathcal{B}(\mathcal{M}_{(\rho, u, \theta)}, \mathcal{M}_{(\rho, u, \theta)}) = 0$$

with respect to ρ , u and θ . See, for instance, [25] or [9] for the missing details. \square

Then

$$\langle \widehat{A} \mathcal{Q}_M(g, g) \rangle = \frac{1}{2} \langle \widehat{A} \mathcal{L}_M(g^2) \rangle = \frac{1}{2} \langle (\mathcal{L}_M \widehat{A}) g^2 \rangle = \frac{1}{2} \langle A g^2 \rangle = u \otimes u - \frac{1}{3} |u|^2 I, \quad (6.21)$$

1 and likewise

$$2 \quad \langle \widehat{B} \mathcal{Q}_M(g, g) \rangle = \frac{1}{2} \langle \widehat{B} \mathcal{L}_M(g^2) \rangle = \frac{1}{2} \langle (\mathcal{L}_M \widehat{B}) g^2 \rangle = \frac{1}{2} \langle B g^2 \rangle = \frac{5}{2} u \theta. \quad (6.22)$$

3
4
5 Observe that

$$6 \quad \operatorname{div}_x \left(u \otimes u - \frac{1}{3} |u|^2 I \right) = \operatorname{div}_x (u \otimes u) - \frac{1}{3} \nabla_x |u|^2,$$

7
8
9 while

$$10 \quad \operatorname{div}_x \left(\nabla_x u + (\nabla_x u)^T - \frac{2}{3} (\operatorname{div}_x u) I \right) = \Delta_x u + \nabla_x (\operatorname{div}_x u) - \frac{2}{3} \nabla_x (\operatorname{div}_x u)$$

$$11 \quad = \Delta_x u,$$

12 because of the divergence-free condition on u .

13 Gathering (6.17)–(6.22), we arrive at

$$14 \quad \partial_t u + \operatorname{div}_x (u \otimes u) - \nu \Delta_x u = 0 \quad \text{modulo gradients,}$$

$$15 \quad \partial_t \theta + \operatorname{div}_x (u \theta) - \kappa \Delta_x \theta = 0. \quad \square$$

16 6.2. The incompressible Stokes and Euler limits

17 By the same moment method, one can derive other incompressible models from the Boltzmann equation. We just state the results below without giving the proofs (which are anyway simpler than that of the Navier–Stokes limit).

18
19
20 **6.2.1. The Stokes limit.** The Stokes–Fourier system is the linearization about $u = 0$ and $\theta = 0$ of the Navier–Stokes–Fourier system. Thus, in order to derive the Stokes–Fourier system from the Boltzmann equation, one keeps the same scaling as for the incompressible Navier–Stokes limit on the Boltzmann equation, i.e.,

$$21 \quad Kn = St = \varepsilon \ll 1$$

22 and one scales the Mach number as

$$23 \quad Ma = \eta_\varepsilon = o(\varepsilon).$$

24 In other words, we start from the following Cauchy problem

$$25 \quad \varepsilon \partial_t F_\varepsilon + v \cdot \nabla_x F_\varepsilon = \frac{1}{\varepsilon} \mathcal{B}(F_\varepsilon, F_\varepsilon), \quad (t, x, v) \in \mathbb{R}_+^* \times \mathbb{R}^3 \times \mathbb{R}^3,$$

$$26 \quad F_\varepsilon(t, x, v) \rightarrow M \quad \text{as } |x| \rightarrow +\infty, \quad (6.23)$$

1 with initial condition

$$2 \quad F_\varepsilon|_{t=0} = \mathcal{M}_{(1-\eta_\varepsilon\theta^{\text{in}}, \frac{\eta_\varepsilon u^{\text{in}}}{1-\eta_\varepsilon\theta^{\text{in}}}, \frac{1}{1-\eta_\varepsilon\theta^{\text{in}}})}. \quad (6.24)$$

3
4
5
6 In this subsection, the number density fluctuation g_ε is defined to be

$$7 \quad g_\varepsilon = \frac{F_\varepsilon - M}{\eta_\varepsilon M}.$$

8
9
10
11 **THEOREM 6.3** (Bardos, Golse and Levermore [9]). *For each $\varepsilon > 0$, let F_ε be a solution*
12 *of (6.23)–(6.24). Assume that*

$$13 \quad \frac{F_\varepsilon - M}{\eta_\varepsilon M} \rightarrow g \quad \text{in the sense of distributions on } \mathbb{R}_+^* \times \mathbb{R}^3,$$

14
15
16
17 that F_ε satisfies the local conservation laws of mass, momentum and energy, and that

$$18 \quad \langle v g_\varepsilon \rangle \rightarrow \langle v g \rangle \quad \text{and} \quad \langle (|v|^2 - 5) g_\varepsilon \rangle \rightarrow \langle (|v|^2 - 5) g \rangle \quad \text{in } C(\mathbb{R}_+, \mathcal{D}'(\mathbb{R}^3))$$

19
20
21 while

$$22 \quad \mathcal{L}_M g_\varepsilon \rightarrow \mathcal{L}_M g \quad \text{in } \mathcal{D}'(\mathbb{R}_+^* \times \mathbb{R}^3)$$

23
24
25
26 and all formally small terms vanish in the sense of distributions as $\varepsilon \rightarrow 0$. Assume further
27 that

$$28 \quad \langle v \otimes v g_\varepsilon \rangle \rightarrow \langle v \otimes v g \rangle, \quad \langle B g_\varepsilon \rangle \rightarrow \langle B g \rangle,$$

$$29 \quad \langle \widehat{A} \otimes v g_\varepsilon \rangle \rightarrow \langle \widehat{A} \otimes v g \rangle \quad \text{and} \quad \langle \widehat{B} \otimes v g_\varepsilon \rangle \rightarrow \langle \widehat{B} \otimes v g \rangle$$

30
31
32 in the sense of distributions on $\mathbb{R}_+^* \times \mathbb{R}^3$.

33 Then g is of the form

$$34 \quad g(t, x, v) = u(t, x) \cdot v + \theta(t, x) \frac{1}{2} (|v|^2 - 5),$$

35
36
37
38 where (u, θ) satisfy the incompressible Stokes–Fourier system

$$39 \quad \partial_t u + \nabla_x p = \nu \Delta_x u, \quad \operatorname{div}_x u = 0,$$

$$40 \quad \partial_t \theta = \kappa \Delta_x \theta, \quad (6.25)$$

41
42
43
44
45 with ν and κ given by formula (6.9).

6.2.2. The incompressible Euler limit. The incompressible Euler equations are formally the inviscid limit of the Navier–Stokes equations. Thus, in order to derive the incompressible Euler equations from the Boltzmann equation, one chooses a scaling that increases the strength of the nonlinear term. In other words, one takes

$$St = Ma = \varepsilon \quad \text{while} \quad Kn = \eta_\varepsilon = o(\varepsilon).$$

In fact, this is consistent with the von Karman relation, which relates the Mach, Knudsen and Reynolds numbers as follows

$$Kn = \frac{Ma}{Re}. \quad (6.26)$$

Indeed, one gets $Re = \varepsilon/\eta_\varepsilon \rightarrow +\infty$ as $\varepsilon \rightarrow 0$: therefore, the limiting equation obtained in this way is incompressible (since $Ma \rightarrow 0$) and inviscid (since $Re \rightarrow +\infty$).

In other words, we start from the following Cauchy problem

$$\begin{aligned} \varepsilon \partial_t F_\varepsilon + v \cdot \nabla_x F_\varepsilon &= \frac{1}{\eta_\varepsilon} \mathcal{B}(F_\varepsilon, F_\varepsilon), \quad (t, x, v) \in \mathbb{R}_+^* \times \mathbb{R}^3 \times \mathbb{R}^3, \\ F_\varepsilon(t, x, v) &\rightarrow M \quad \text{as } |x| \rightarrow +\infty, \end{aligned} \quad (6.27)$$

with initial condition

$$F_\varepsilon|_{t=0} = \mathcal{M}_{(1-\varepsilon\theta^{\text{in}}, \frac{\varepsilon u^{\text{in}}}{1-\varepsilon\theta^{\text{in}}}, \frac{1}{1-\varepsilon\theta^{\text{in}}})}. \quad (6.28)$$

In this subsection, the number density fluctuation g_ε is defined to be

$$g_\varepsilon = \frac{F_\varepsilon - M}{\varepsilon M}.$$

THEOREM 6.4 (Bardos, Golse and Levermore [9]). *For each $\varepsilon > 0$, let F_ε be a solution of (6.27)–(6.28). Assume that*

$$\frac{F_\varepsilon - M}{\varepsilon M} \rightarrow g \quad \text{in the sense of distributions on } \mathbb{R}_+^* \times \mathbb{R}^3,$$

that F_ε satisfies the local conservation laws of mass, momentum and energy, and that

$$\langle v g_\varepsilon \rangle \rightarrow \langle v g \rangle \quad \text{and} \quad \langle (|v|^2 - 5) g_\varepsilon \rangle \rightarrow \langle (|v|^2 - 5) g \rangle \quad \text{in } C(\mathbb{R}_+, \mathcal{D}'(\mathbb{R}^3))$$

while

$$\mathcal{L}_M g_\varepsilon \rightarrow \mathcal{L}_M g \quad \text{in } \mathcal{D}'(\mathbb{R}_+^* \times \mathbb{R}^3)$$

1 and all formally small terms vanish in the sense of distributions as $\varepsilon \rightarrow 0$. Assume further
2 that

$$3 \quad \langle v \otimes v g_\varepsilon \rangle \rightarrow \langle v \otimes v g \rangle, \quad \langle B g_\varepsilon \rangle \rightarrow \langle B g \rangle,$$

$$4 \quad \langle \widehat{A} \mathcal{Q}_M(g_\varepsilon, g_\varepsilon) \rangle \rightarrow \langle \widehat{A} \mathcal{Q}_M(g_\varepsilon, g_\varepsilon) \rangle \quad \text{and} \quad \langle \widehat{B} \mathcal{Q}_M(g_\varepsilon, g_\varepsilon) \rangle \rightarrow \langle \widehat{B} \mathcal{Q}_M(g_\varepsilon, g_\varepsilon) \rangle$$

5
6
7 in the sense of distributions on $\mathbb{R}_+^* \times \mathbb{R}^3$.

8 Then g is of the form

$$9 \quad g(t, x, v) = u(t, x) \cdot v + \theta(t, x) \frac{1}{2} (|v|^2 - 5),$$

10
11 where (u, θ) satisfy the system

$$12 \quad \partial_t u + \operatorname{div}_x(u \otimes u) + \nabla_x p = 0, \quad \operatorname{div}_x u = 0,$$

$$13 \quad \partial_t \theta + \operatorname{div}_x(u \theta) = 0. \quad (6.29)$$

14 6.3. Other incompressible models

15
16 There are many possible variants of the incompressible Navier–Stokes–Fourier limit de-
17 scribed above. To begin with, it is possible to include a conservative force in the Boltzmann
18 equation. The scaling is as follows. Start from equation

$$19 \quad \varepsilon \partial_t F_\varepsilon + v \cdot \nabla_x F_\varepsilon - \varepsilon \nabla_x \phi(x) \cdot \nabla_v F_\varepsilon = \frac{1}{\varepsilon} \mathcal{B}(F_\varepsilon, F_\varepsilon),$$

20
21 where $\phi \equiv \phi(x)$ is a given (smooth) potential. Writing the (x, v) -derivative in the
22 Boltzmann equation above as a Poisson bracket, i.e.,

$$23 \quad v \cdot \nabla_x F_\varepsilon - \varepsilon \nabla_x \phi(x) \cdot \nabla_v F_\varepsilon = \left\{ \frac{1}{2} |v|^2 + \varepsilon \phi(x); F_\varepsilon \right\}$$

24
25 suggests to seek the solution F_ε in the form

$$26 \quad F_\varepsilon(t, x, v) = e^{-\varepsilon \phi(x)} M(v) (1 + \varepsilon g_\varepsilon(t, x, v)).$$

27
28 Indeed,

$$29 \quad e^{-\varepsilon \phi(x)} M(v) = \frac{1}{(2\pi)^{3/2}} e^{-|v|^2/2 - \varepsilon \phi(x)}$$

30
31 is both a Maxwellian and an element of the nullspace of the Poisson bracket $\left\{ \frac{1}{2} |v|^2 + \right.$
32 $\left. \varepsilon \phi(x); \cdot \right\}$.
33
34
35
36
37
38
39
40
41
42
43
44
45

The same formal argument as above shows that

$$g_\varepsilon(t, x, v) \rightarrow u(t, x) \cdot v + \theta(t, x) \frac{1}{2}(|v|^2 - 5),$$

where u and θ satisfy

$$\partial_t u + \operatorname{div}_x(u \otimes u) + \nabla_x p - \theta \nabla_x \phi = \nu \Delta_x u,$$

$$\partial_t \theta + \operatorname{div}_x(u \theta) + \frac{2}{5} u \cdot \nabla_x \phi = \kappa \Delta_x \theta,$$

with

$$\nu = \frac{1}{10} \langle \widehat{A} : A \rangle \quad \text{and} \quad \kappa = \frac{2}{15} \langle \widehat{B} \cdot B \rangle$$

as in the incompressible Navier–Stokes–Fourier limit theorem above. Of course, this is in agreement with the discussion in Section 2.7. We refer the interested reader to [8] for more details on the formal derivation.

In still another variant of the Navier–Stokes–Fourier limit presented above, it is possible to recover viscous heating terms as in Section 2.6. As explained in that subsection, viscous heating terms should appear when the fluctuations of velocity field are of the order of the square root of temperature fluctuations. In [13], Levermore and Bardos used the following very elegant approach: start from the Boltzmann equation in the Navier–Stokes scaling (6.2) and seek the number density F_ε as

$$F_\varepsilon(t, x, v) = M(v) \left(1 + \varepsilon g_\varepsilon^-(t, x, v) + \varepsilon^2 g_\varepsilon^+(t, x, v) \right),$$

where g_ε^- is odd in v while g_ε^+ is even in v . Because the Boltzmann collision integral is rotation invariant (see Section 3.6.2 and especially (3.51)),

$$\mathcal{B}(\Phi^+, \Phi^+) \text{ and } \mathcal{B}(\Phi^-, \Phi^-) \text{ are even in } v, \text{ while}$$

$$\mathcal{B}(\Phi^-, \Phi^+) \text{ and } \mathcal{B}(\Phi^+, \Phi^-) \text{ are odd in } v.$$

Levermore and Bardos gave a formal argument showing that

$$g_\varepsilon^-(t, x, v) \rightarrow u(t, x) \cdot v,$$

while

$$g_\varepsilon^+(t, x, v) \rightarrow \rho(t, x) + \frac{1}{2} (|u(t, x)|^2 + 3\theta(t, x)) \left(\frac{1}{3} |v|^2 - 1 \right) \\ + \frac{1}{2} A : u(t, x) \otimes u(t, x) - \widehat{A} : \nabla_x u(t, x),$$

1 where ρ , u and θ satisfy the relations

$$2 \quad \operatorname{div}_x u = 0, \quad p = \rho + \theta, \quad 3$$

$$4 \quad \partial_t u + \operatorname{div}_x(u \otimes u) + \nabla_x p = \nu \Delta_x u, \quad 5$$

$$6 \quad \partial_t \left(\frac{5}{2} \theta + \frac{1}{2} |u|^2 - p \right) + \operatorname{div}_x \left(u \left(\frac{5}{2} \theta + \frac{1}{2} |u|^2 \right) \right) \quad 7$$

$$8 \quad = \frac{5}{2} \kappa \Delta_x \theta + \mu \operatorname{div}_x \left((\nabla_x u + (\nabla_x u)^T) \cdot u \right). \quad 9$$

10 In the system above, ν and κ are given by the same formulas as in the Navier–Stokes–
11 Fourier limit theorem, i.e., (6.9). 12

13 7. Mathematical theory of the Cauchy problem for hydrodynamic models 14

15 In this section we have gathered a few mathematical results bearing on the various hydrodynamic models that appear as limits of the Boltzmann equation. We shall leave aside the compressible Navier–Stokes system, since its derivation from the Boltzmann equation leads to dissipation terms that are of the order of the Knudsen number, and therefore vanish in the hydrodynamic limit. Put in other words, the compressible Navier–Stokes system is an asymptotic expansion of the Boltzmann equation in the Knudsen number, and not a limit thereof. Hence, a mathematical treatment of this limit would involve existence results on the compressible Navier–Stokes system that are uniform in the Reynolds and Péclet numbers, which is beyond current knowledge on this model at the time of this writing. 16

17 7.1. The Stokes and acoustic systems 18

19 We begin with the simplest hydrodynamic models 20

- 21 • the Stokes–Fourier system, and 22
- 23 • the acoustic system, 23

24 which are variants of the heat and the wave equations. 24

25 **7.1.1. The Stokes–Fourier system.** Consider the Stokes–Fourier system 25

$$26 \quad \partial_t u + \nabla_x p = \nu \Delta_x u, \quad \operatorname{div}_x u = 0, \quad (t, x) \in \mathbb{R}_+^* \times \mathbb{R}^3, \quad 27$$

$$28 \quad \partial_t \theta = \kappa \Delta_x \theta, \quad 28$$

$$29 \quad (u, \theta)|_{t=0} = (u^{\text{in}}, \theta^{\text{in}}), \quad 29$$

30 where ν and $\kappa > 0$. 30

31 **THEOREM 7.1.** For each $(u^{\text{in}}, \theta^{\text{in}}) \in L^2(\mathbb{R}^3)$ such that $\operatorname{div}_x u^{\text{in}} = 0$, there exists a unique
32 solution (u, θ) of the Stokes–Fourier system such that 32

$$33 \quad (u, \theta) \in C(\mathbb{R}_+; L^2(\mathbb{R}^3)) \quad \text{and} \quad p \in C(\mathbb{R}_+^*; L^2(\mathbb{R}^3)/\mathbb{R}). \quad 33$$

1 In addition, the pressure p is constant and

$$2 \quad (u, \theta) \in C^\infty(\mathbb{R}_+^* \times \mathbb{R}^3).$$

3
4
5 PROOF. Applying the divergence to both sides of the motion equation gives

$$6 \quad \Delta_x p = 0;$$

7
8
9 since for each $t > 0$ the function $p(t, \cdot) \in L^2(\mathbb{R}^3)/\mathbb{R}$ is harmonic, it is a constant in
10 variable x . Since the divergence operator commutes with the heat operator on \mathbb{R}^3 , the
11 Stokes–Fourier system above reduces to a system of uncoupled heat equations, whence the
12 announced result. \square

13
14 **7.1.2. The acoustic system.** Consider the acoustic system

$$15 \quad \partial_t \rho + \operatorname{div}_x u = 0,$$

$$16 \quad \partial_t u + \nabla_x(\rho + \theta) = 0, \quad (t, x) \in \mathbb{R}_+^* \times \mathbb{R}^3,$$

$$17 \quad \frac{3}{2} \partial_t \theta + \operatorname{div}_x u = 0,$$

$$18 \quad (\rho, u, \theta)|_{t=0} = (\rho^{\text{in}}, u^{\text{in}}, \theta^{\text{in}}).$$

19
20
21
22 THEOREM 7.2. For each $(\rho^{\text{in}}, u^{\text{in}}, \theta^{\text{in}}) \in L^2(\mathbb{R}^3)$, there exists a unique solution (ρ, u, θ)
23 of the acoustic system such that

$$24 \quad (\rho, u, \theta) \in C(\mathbb{R}_+; L^2(\mathbb{R}^3)).$$

25
26
27 PROOF. Applying the Helmholtz decomposition to $u(t, \cdot)$ gives

$$28 \quad u(t, \cdot) = u^s(t, \cdot) - \nabla_x \phi(t, \cdot), \quad \operatorname{div}_x u_s = 0.$$

29
30
31 Hence the acoustic system becomes

$$32 \quad \partial_t \rho - \Delta_x \phi = 0, \quad \frac{3}{2} \partial_t \theta - \Delta_x \phi = 0,$$

$$33 \quad \partial_t \phi - \rho - \theta = 0, \quad \partial_t u_s = 0.$$

34
35
36 Hence

$$37 \quad \partial_t \phi - \rho - \theta = 0, \quad \partial_t(\rho + \theta) - \frac{5}{3} \Delta_x \phi = 0,$$

$$38 \quad \partial_t \left(\rho - \frac{3}{2} \theta \right) = 0.$$

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

1 Therefore, letting $\psi = \rho + \theta$, we arrive at

$$2 \quad \partial_{tt}\phi - \frac{5}{3}\Delta_x\phi = 0,$$

$$3 \quad \partial_{tt}\psi - \frac{5}{3}\Delta_x\psi = 0.$$

4 This is a system of two uncoupled wave equations; then, (ρ, u, θ) is reconstructed from
5 ψ and $\mu \equiv \mu(x) = \rho(t, x) - \frac{3}{2}\theta(t, x)$ by the formulas

$$6 \quad \rho(t, x) = \frac{3}{5}\psi(t, x) + \frac{2}{5}\mu(x),$$

$$7 \quad \theta(t, x) = \frac{2}{5}(\psi(t, x) - \mu(x)),$$

8 while

$$9 \quad u(t, x) = u^s(0, x) - \nabla_x\phi(t, x).$$

10 Applying the classical theory of the Cauchy problem for the wave equations satisfied by
11 ϕ and ψ leads to the announced result. \square

12 7.2. The incompressible Navier–Stokes equations

13 Next, we consider the incompressible Navier–Stokes equations. The mathematical theory
14 of the Cauchy problem for these equations was developed by J. Leray in the early 1930s.
15 Here is a quick summary of his results, see for instance [30,45,86] for more information
16 on this subject.

17 Consider the Navier–Stokes equations

$$18 \quad \partial_t u + \operatorname{div}_x(u \otimes u) + \nabla_x p = \nu \Delta_x u, \quad \operatorname{div}_x u = 0, \quad (t, x) \in \mathbb{R}_+^* \times \mathbb{R}^D,$$

$$19 \quad u|_{t=0} = u^{\text{in}},$$

20 where $\nu > 0$.

21 We begin with the three-dimensional case.

22 **THEOREM 7.3** (Leray [80]). *For each $u^{\text{in}} \in L^2(\mathbb{R}^3)$ such that $\operatorname{div}_x u^{\text{in}} = 0$, there exists
23 a solution in the sense of distributions to the Cauchy problem for the Navier–Stokes equa-
24 tions such that*

$$25 \quad u \in C(\mathbb{R}_+; w - L^2(\mathbb{R}^3)) \cap L^2(\mathbb{R}_+; H^1(\mathbb{R}^3)).$$

Moreover, the function $t \mapsto \|u(t, \cdot)\|_{L^2}$ is nonincreasing on \mathbb{R}_+ and satisfies, for each $t > 0$, the Leray energy inequality

$$\frac{1}{2} \|u(t)\|_{L^2}^2 + \int_0^t \int_{\mathbb{R}^D} \nu |\nabla_x u(s, x)|^2 dx ds \leq \frac{1}{2} \|u^{\text{in}}\|_{L^2}^2.$$

A solution of the Navier–Stokes equations in the sense of distributions that belongs to $C(\mathbb{R}_+; w - L^2(\mathbb{R}^3)) \cap L^2(\mathbb{R}_+; H^1(\mathbb{R}^3))$ and satisfies the Leray energy inequality is called a “Leray solution”.

In fact, a modification of Leray’s original argument allows constructing weak solutions that satisfy the *local* energy inequality

$$\partial_t \frac{1}{2} |u|^2 + \operatorname{div}_x \left(u \left(\frac{1}{2} |u|^2 + p \right) \right) + \nu |\nabla_x u|^2 \leq \nu \Delta_x \left(\frac{1}{2} |u|^2 \right) \quad (7.1)$$

in the sense of distributions on $\mathbb{R}_+^* \times \mathbb{R}^3$.

It is not known whether Leray solutions are uniquely determined by their initial data; however, Leray was able to prove that regular solutions of the Navier–Stokes equations are unique within the class of Leray solutions.

THEOREM 7.4 (Leray [80]). *Let $u^{\text{in}} \in L^2(\mathbb{R}^3)$ such that $\operatorname{div}_x u^{\text{in}} = 0$. Assume that there exists a classical solution,*

$$v \in C^1(\mathbb{R}_+; H^1(\mathbb{R}^3)) \cap C(\mathbb{R}_+; H^2(\mathbb{R}^3)) \quad \text{such that} \quad \nabla_x v \in L^\infty(\mathbb{R}_+ \times \mathbb{R}^3),$$

of the Navier–Stokes equations with initial condition

$$v|_{t=0} = u^{\text{in}}.$$

Then, any Leray solution u of the Navier–Stokes equations with initial data

$$u|_{t=0} = u^{\text{in}}$$

coincides with v a.e.

Whether the space dimension is $D = 2$ or $D = 3$ leads to fundamental differences in the regularity theory for the Navier–Stokes equations.

THEOREM 7.5 (Leray [79]). *Let $u^{\text{in}} \in L^2(\mathbb{R}^2)$ such that $\operatorname{div}_x u^{\text{in}} = 0$. Then there exists a unique weak solution u to the Navier–Stokes equations with initial data*

$$u|_{t=0} = u^{\text{in}}$$

such that

$$u \in C(\mathbb{R}_+; L^2(\mathbb{R}^2)) \cap L^2(\mathbb{R}_+; H^1(\mathbb{R}^2)).$$

1 *Furthermore, this solution is smooth for $t > 0$*

$$2 \quad u \in C^\infty(\mathbb{R}_+^* \times \mathbb{R}^2)$$

3 *and satisfies the energy equality*

$$4 \quad \frac{1}{2} \|u(t)\|_{L^2}^2 + \int_0^t \int_{\mathbb{R}^D} v |\nabla_x u(s, x)|^2 dx ds = \frac{1}{2} \|u^{\text{in}}\|_{L^2}^2$$

5 *for each $t \geq 0$.*

6
7
8
9
10
11
12 In space dimension 3, it is an outstanding open problem to determine whether Leray
13 solutions with C^∞ initial data remain C^∞ for all times.

14 What is known to this date is the following partial regularity theorem which improves
15 on an earlier result by Scheffer [112].

16
17 **THEOREM 7.6** (Caffarelli, Kohn and Nirenberg [21]). *In space dimension 3, let u be a
18 Leray solution of the Navier–Stokes equations that satisfies the local variant of Leray’s
19 energy inequality (7.1). Let the singular set of u be*

$$20 \quad S(u) = \{(t, x) \in \mathbb{R}_+ \times \mathbb{R}^3 \mid u \text{ is not bounded in a neighborhood of } (t, x)\}.$$

21
22 *Then, $S(u)$ has parabolic Hausdorff dimension less than 1.*

23
24
25 This definition of the singular set $S(u)$ comes from a bootstrap argument due to
26 Serrin [113] showing that, if a Leray solution u of the Navier–Stokes equations is bounded
27 in $B((t, x), R)$, then u is C^∞ in $B((t, x), R/2)$.

28 The parabolic Hausdorff dimension is defined through coverings with translates of
29 $(-r^2, r^2) \times B(0, r)$ in $\mathbb{R} \times \mathbb{R}^3$ (the usual Hausdorff dimension being defined with balls
30 for the Euclidean metric of \mathbb{R}^4).

31 This result implies that the singular set $S(u)$ must be smaller than a curve in space–
32 time: in other words singularities of solutions to the Navier–Stokes equations in space
33 dimension 3 are rare (if they exist at all).

34 35 36 **7.3. The incompressible Navier–Stokes–Fourier system**

37
38 By mimicking the compactness method in the proof of the Leray existence theorem, we
39 can also treat the case of the Navier–Stokes–Fourier system

$$40 \quad \partial_t u + \text{div}_x(u \otimes u) + \nabla_x p = \nu \Delta_x u, \quad \text{div}_x u = 0, \quad (t, x) \in \mathbb{R}_+^* \times \mathbb{R}^3,$$

$$41 \quad \partial_t \theta + \text{div}_x(u\theta) = \kappa \Delta_x \theta, \quad (t, x) \in \mathbb{R}_+^* \times \mathbb{R}^3,$$

$$42 \quad (u, \theta)|_{t=0} = (u^{\text{in}}, \theta^{\text{in}}),$$

$$43 \quad$$

1 where $\kappa > 0$ and $\nu > 0$. 1

2 The analogue of Leray's theorem for the Navier–Stokes–Fourier system is the following 2
3 one. 3

4
5 **THEOREM 7.7.** *For each $(u^{\text{in}}, \theta^{\text{in}}) \in L^2(\mathbb{R}^3)$ such that $\operatorname{div}_x u^{\text{in}} = 0$, there exists a solution 5
6 in the sense of distributions to the Cauchy problem for the Navier–Stokes–Fourier system 6
7 such that 7*

$$8 \quad (u, \theta) \in C(\mathbb{R}_+; w - L^2(\mathbb{R}^3)) \cap L^2(\mathbb{R}_+; H^1(\mathbb{R}^3)). \quad 9$$

10
11 *This solution satisfies, for each $t > 0$ 11*

$$12 \quad \frac{1}{2} \|u(t)\|_{L^2}^2 + \int_0^t \int_{\mathbb{R}^D} \nu |\nabla_x u(s, x)|^2 dx ds \leq \frac{1}{2} \|u^{\text{in}}\|_{L^2}^2, \quad 13$$

$$14 \quad \frac{1}{2} \|\theta(t)\|_{L^2}^2 + \int_0^t \int_{\mathbb{R}^D} \kappa |\nabla_x \theta(s, x)|^2 dx ds \leq \frac{1}{2} \|\theta^{\text{in}}\|_{L^2}^2. \quad 15$$

16 7.4. The compressible Euler system 16

17
18 The compressible Euler system is a quasilinear hyperbolic system. The existence and 18
19 uniqueness theory for this system is not entirely satisfying in its present state, especially in 19
20 space dimension greater than or equal to 2. More information on the theory of hyperbolic 20
21 system of conservation laws can be found for instance in [32,78] and [19]. 21

22
23 Consider the Cauchy problem for the compressible Euler system (with perfect mon- 22
24 atomic gas equation of state) 23
24
25
26

$$27 \quad \begin{aligned} 27 \quad \partial_t \rho + \operatorname{div}_x(\rho u) &= 0, & 27 \\ 28 \quad \partial_t(\rho u) + \operatorname{div}_x(\rho u \otimes u) + \nabla_x(\rho \theta) &= 0, & 28 \\ 29 \quad \partial_t \left(\rho \left(\frac{1}{2} |u|^2 + \frac{3}{2} \theta \right) \right) + \operatorname{div}_x \left(\rho \left(\frac{1}{2} |u|^2 + \frac{5}{2} \theta \right) \right) &= 0. & 29 \end{aligned}$$

30
31 We begin with a local existence and uniqueness result which is a particular case of a gen- 30
32 eral theorem on quasilinear *symmetrizable systems*. The theory of symmetric hyperbolic 31
33 system was developed very early by Friedrichs; the importance of the notion of symmetriz- 32
34 able systems was recognized by Godunov [52], and then by Friedrichs and Lax [43] – see 33
35 also [46] and [81] for more information on the theory of hyperbolic systems. The result 34
36 below comes from [91]; the case of general systems is studied in [78] and [72]. 35

36
37
38
39
40
41
42
43 **THEOREM 7.8.** *Let $D \geq 1$ and $(\rho^{\text{in}}, u^{\text{in}}, \theta^{\text{in}}) \in H^m(\mathbb{R}^D)$ with $m > D/2 + 1$. There exists 43
44 $T > 0$ and a unique solution $(\rho, u, \theta) \in C([0, T]; H^m(\mathbb{R}^D)) \cap C^1([0, T]; H^{m-1}(\mathbb{R}^D))$ of 44
45 the compressible Euler system in the sense of distributions on $(0, T) \times \mathbb{R}^D$. 45*

1 Such solutions are regular (since $H^m(\mathbb{R}^D) \subset C^{\lfloor m-D/2 \rfloor}(\mathbb{R}^D)$ by Sobolev embedding). 1
 2 In general, one does not expect that regular solutions to (7.2) may exist for all times. Sing- 2
 3 gularities – such as shock waves – are expected to appear in finite time for a large class of 3
 4 smooth initial data. Here is an interesting result in this direction, due to Sideris [114]. 4
 5

6 **THEOREM 7.9.** *Let $D = 3$, and let $R > 0$. Pick the initial data $(\rho^{\text{in}}, u^{\text{in}}, \theta^{\text{in}}) \in C^\infty(\mathbb{R}^3)$ 6
 7 to be such that $\rho^{\text{in}}, \theta^{\text{in}} > 0$ on \mathbb{R}^3 with 7
 8*

$$\begin{aligned} \text{supp}(\rho^{\text{in}} - 1) &\subset B(0, R), & \text{supp}(u^{\text{in}}) &\subset B(0, R), \\ \text{supp}(\theta^{\text{in}} - 1) &\subset B(0, R) & \text{and } \rho^{\text{in}} &\geq (\theta^{\text{in}})^{3/2} \text{ on } \mathbb{R}^3. \end{aligned}$$

9
 10
 11
 12 Assume further the existence of $R_0 \in (0, R)$ such that 13
 14

$$\int_{|x|>r} \frac{(|x| - r)^2}{|x|} (\rho^{\text{in}}(x) - 1) dx > 0$$

15
 16
 17 and 18

$$\int_{|x|>r} \frac{(|x|^2 - r^2)}{|x|^3} \rho^{\text{in}}(x) x \cdot u^{\text{in}}(x) dx \geq 0$$

19
 20
 21
 22 for each $r \in (R_0, R)$. Then the life-span T of the C^1 solution to (7.2) with such initial data 23
 24 is finite. 24
 25

26 In dimension greater than or equal to 2, there is no satisfying theory of weak solutions 26
 27 that could extend classical solutions after blow-up time. 27

28 At variance, in the one-dimensional case, there is a rather complete theory of weak solu- 28
 29 tions, that are obtained as superpositions of interacting Riemann problems (i.e., a Cauchy 29
 30 problem for (5.6) whose initial data is a step function with only one jump). Liu studied 30
 31 the compressible Euler system in space dimension 1 written in Lagrangian coordinates. 31
 32 Denoting by a the Lagrangian particle label, this system reads 32
 33

$$\begin{aligned} \partial_t V - \partial_a U &= 0, \\ \partial_t U + \partial_a \left(\frac{\Theta}{V} \right) &= 0, \\ \partial_t \left(\frac{1}{2} U^2 + \frac{3}{2} \Theta \right) + \partial_a \left(\frac{\Theta U}{V} \right) &= 0, \\ (V, U, \Theta)|_{t=0} &= (V^{\text{in}}, U^{\text{in}}, \Theta^{\text{in}}), \end{aligned} \tag{7.3}$$

34
 35
 36
 37
 38
 39
 40
 41 with the notation 42
 43

$$V(t, a) = \frac{1}{\rho(t, X(t, 0, a))},$$

$$U(t, a) = u(t, X(t, 0, a)),$$

$$\Theta(t, a) = \theta(t, X(t, 0, a)),$$

where $t \mapsto X(t, 0, a)$ is the path of the particle which is at the position a at time $t = 0$.

The following existence result is based on Glimm's algorithm [51] for constructing BV solutions to hyperbolic systems of conservation laws in the one-dimensional case that are global in time for initial data of small total variation.

THEOREM 7.10 (Liu [89]). *Assume that $(V^{\text{in}}, u^{\text{in}}, \theta^{\text{in}}) \in BV(\mathbb{R})$ with $\theta^{\text{in}} \geq \theta_* > 0$ while $V^{\text{in}} \leq V^*$ on \mathbb{R} . There exists $\eta_0 > 0$ such that the Cauchy problem (7.3) has a global weak solution provided that $TV(V^{\text{in}}, u^{\text{in}}, \theta^{\text{in}}) \leq \eta_0$. Moreover, this solution is “entropic”, i.e.,*

$$\partial_t \ln(V^{2/3} \Theta) \geq 0.$$

(In other words, the entropy density cannot decrease along particle paths.)

7.5. The incompressible Euler equations

There are essentially two main directions in the mathematical theory of the incompressible Euler equations:

- the PDE viewpoint, and
- the geometric viewpoint.

In the geometric viewpoint, Euler's equations for incompressible fluids in the periodic box \mathbb{T}^D are the equations of geodesics on the group of volume preserving diffeomorphisms of \mathbb{T}^D , endowed with the metric defined by the kinetic energy. We shall say nothing of this part of the theory, for which we refer the reader to the beautiful book by Arnold and Khesin [4].

Instead, we shall just recall a few results on the Euler equations as nonlinear PDEs on \mathbb{T}^D . The reader is advised to read [92] and [86] for more information on that topic.

The incompressible Euler equations are

$$\partial_t u + \operatorname{div}_x(u \otimes u) + \nabla_x p = 0, \quad \operatorname{div}_x u = 0, \quad (t, x) \in \mathbb{R}_+^* \times \mathbb{T}^D,$$

$$u|_{t=0} = u^{\text{in}}.$$

We begin with a local existence result for classical solutions in the three-dimensional case, due to Kato.

THEOREM 7.11 (Kato [71]). *Let $u^{\text{in}} \in H^m(\mathbb{T}^3)$, $m \in \mathbb{N}$, $m \geq 3$, such that $\operatorname{div}_x u^{\text{in}} = 0$. Then, there exists $T > 0$ and a unique local solution u of the incompressible Euler equations with initial data u^{in} , such that*

$$u \in C([0, T]; H^m(\mathbb{T}^3)) \cap AC_{\text{loc}}([0, T]; H^{m-1}(\mathbb{T}^3)).$$

Whether the life-span of such solutions is finite is an outstanding open problem in the theory of nonlinear PDEs. What is known is that the blow-up time, if finite, does not depend on the regularity index m , as shown by the following beautiful result.

THEOREM 7.12 (Beale, Kato and Majda [15]). *Under the same assumptions as in the previous theorem, if T is finite, then*

- either

$$\int_0^T \|\operatorname{curl}_x u(t, \cdot)\|_{L^\infty} dt = +\infty,$$

- or $u \in C([0, T]; H^m(\mathbb{T}^3))$.

(In the second case, the solution u can be extended to an interval of time $[0, T')$ with $T' > T$.)

In the two-dimensional case, there is global existence and uniqueness of a classical solution to the Cauchy problem for the incompressible Euler equations.

THEOREM 7.13 (Yudovich [123]). *Let $u^{\text{in}} \in H^m(\mathbb{T}^2)$, $m \in \mathbb{N}$, $m \geq 3$, such that $\operatorname{div}_x u^{\text{in}} = 0$. Then there exists a unique global solution u of the incompressible Euler equations with initial data u^{in} such that*

$$u \in C(\mathbb{R}_+; H^m(\mathbb{T}^2)) \cap C^1(\mathbb{R}_+; H^{m-1}(\mathbb{T}^2)).$$

A good reference on the two-dimensional case of the incompressible Euler equations is [29].

In view of the importance of the vorticity field $\operatorname{curl}_x u$ for the regularity of the solution to the incompressible Euler equations, the main difference between the two- and the three-dimensional cases comes from the following observation.

If one considers the two-dimensional flow as given by a three-dimensional velocity field u of the form

$$u(t, x) = \begin{pmatrix} u_1(t, x_1, x_2) \\ u_2(t, x_1, x_2) \\ 0 \end{pmatrix},$$

the vorticity field is

$$\operatorname{curl}_x u(t, x) = \begin{pmatrix} 0 \\ 0 \\ \omega(t, x_1, x_2) \end{pmatrix},$$

where

$$\omega(t, x_1, x_2) = \partial_{x_1} u_2(t, x_1, x_2) - \partial_{x_2} u_1(t, x_1, x_2).$$

1 The scalar quantity ω satisfies the transport equation 1

$$2 \quad \partial_t \omega + u \cdot \nabla_x \omega = 0. \quad 3$$

4
5 The maximum principle holds for the transport equation above, so that 5

$$6 \quad \|\omega\|_{L^\infty(\mathbb{R}_+ \times \mathbb{T}^2)} = \|\omega(0, \cdot)\|_{L^\infty(\mathbb{T}^2)}. \quad 7$$

8
9 Hence, in the two-dimensional case, if u^{in} is sufficiently regular, the vorticity is globally 10
11 bounded and therefore, by the Beale–Kato–Majda criterion, the regularity of the initial data 12
12 is propagated for all times.

13 In the three-dimensional case, the vorticity $\text{curl}_x u$ is a *vector* field that satisfies the 13
14 analogue of the scalar transport equation above for vectors 14

$$15 \quad \partial_t \text{curl}_x u + (u \cdot \nabla_x) \text{curl}_x u = \nabla_x u \cdot \text{curl}_x u. \quad 16$$

17
18 The length of the vector $\text{curl}_x u$ can be amplified, or damped, by the matrix $\nabla_x u$: this mech- 18
19 anism is called “vortex stretching” and so far, there is no satisfying method for controlling 19
20 it. Therefore, there is no a priori bound on the vorticity as in the two-dimensional case, and 20
21 this is why the question of global existence or finite-time blow-up for classical solutions to 21
22 the incompressible Euler equations in the three-dimensional case remains very much open. 22
23

24 Finally, we conclude this section with an important class of global solutions on the 24
25 periodic box to the incompressible Euler equations. Choose 25

$$26 \quad U(t, x) = \begin{pmatrix} u_1(t, x_1, x_2) \\ u_2(t, x_1, x_2) \\ w(t, x_1, x_2) \end{pmatrix}, \quad (t, x) \in \mathbb{R}_+ \times \mathbb{T}^3, \quad 27$$

28
29
30
31 where 31

$$32 \quad u(t, x_1, x_2) = \begin{pmatrix} u_1(t, x_1, x_2) \\ u_2(t, x_1, x_2) \end{pmatrix} \quad 33$$

34
35
36 is a C^1 -solution of the two-dimensional incompressible Euler equations on $\mathbb{R}_+ \times \mathbb{T}^2$. If 36
37 w satisfies the transport equation 37

$$38 \quad \partial_t w + \text{div}_x(wu) = 0, \quad (t, x) \in \mathbb{R}_+^* \times \mathbb{T}^2, \quad 39$$

40
41
42 the vector field U solves the three-dimensional incompressible Euler equation on $\mathbb{R}_+ \times \mathbb{T}^3$. 42
43 Such solutions are referred to as *2D–3C solutions of the incompressible Euler equations* – 43
44 see, in particular, Section S4.3, pp. 150–153 of [86] for an interesting application of 2D–3C 44
45 solutions to the problem of a priori estimates on the Euler equations. 45

8. Mathematical theory of the Cauchy problem for the Boltzmann equation

8.1. Global classical solutions for “small” data

In this subsection, we briefly review two early existence theories for the Boltzmann equation:

- Ukai’s theory for perturbations of uniform Maxwellian states; and
- the Illner–Shinbrot theory for small perturbations of the vacuum state.

Consider the Cauchy problem for the Boltzmann equation

$$\begin{aligned} \partial_t F + v \cdot \nabla_x F &= \mathcal{B}(F, F), \quad (t, x, v) \in \mathbb{R}_+^* \times \mathbb{R}^3 \times \mathbb{R}^3, \\ F|_{t=0} &= F^{\text{in}}. \end{aligned}$$

8.1.1. Small perturbations of the vacuum. The collision kernel b in the collision integral is supposed to satisfy

$$0 \leq b(z, \omega) \leq C(1 + |z|) \quad \text{for a.e. } (z, \omega) \in \mathbb{R}^3 \times \mathbb{S}^2.$$

THEOREM 8.1 (Illner and Shinbrot [70]). *Pick $c > 0$; there exists $\eta > 0$ such that for each $C \in (0, \eta)$ and each initial data F^{in} satisfying*

$$0 \leq F^{\text{in}}(x, v) \leq C e^{-c(|x|^2 + |v|^2)}, \quad x, v \in \mathbb{R}^3,$$

then the Cauchy problem for the Boltzmann equation with initial data F^{in} has a unique global (mild) solution.

The key to this result is to dominate the solution F of the Boltzmann equation with a Maxwellian traveling wave $e^{-c|x-tv|^2}$.

Obviously this result is not useful in the context of incompressible hydrodynamic limits: it bears on rarefied clouds of gas that expand in the vacuum, and therefore never approach any global Maxwellian equilibrium.

8.1.2. Small perturbations of a global Maxwellian state. The following result, due to Ukai, is the first global existence and uniqueness result proved on the (space inhomogeneous) Boltzmann equation. It bears on the dynamics of a gas whose state is a perturbation of a global Maxwellian equilibrium. It uses a detailed spectral analysis of the linearization of the collision integral at the background Maxwellian state, see [40].

THEOREM 8.2 (Ukai [116]). *Assume that $b(z, \omega) = |z \cdot \omega|$ (hard sphere case). Let $s > 3/2$, and $\beta > 3$; define*

$$\|f\|_{s,\beta} = \sup_{v \in \mathbb{R}^3} (1 + |v|)^\beta \|f(\cdot, v)\|_{H^s(\mathbb{R}^3)}$$

1 *and let*

$$2 \quad H_\beta^s = \{f \in L_{\text{loc}}^\infty(\mathbb{R}_v^3; H_x^s) \mid \|f\|_{s,\beta} < +\infty\}.$$

3
4
5 *There exists $\eta > 0$ such that, for any $f^{\text{in}} \equiv f^{\text{in}}(x, v)$ satisfying*

$$6 \quad \|f^{\text{in}}\|_{s,\beta} < \eta \quad \text{and} \quad f^{\text{in}} \geq -\mathcal{M}_{(1,0,1)}^{1/2},$$

7
8
9 *the Cauchy problem for the Boltzmann equation with initial data*

$$10 \quad F^{\text{in}} = \mathcal{M}_{(1,0,1)} + \mathcal{M}_{(1,0,1)}^{1/2} f^{\text{in}}$$

11
12
13
14 *has a unique solution F such that*

$$15 \quad F \in L^\infty(\mathbb{R}_+; H_\beta^s) \cap C(\mathbb{R}_+; H_{\beta-\varepsilon}^{s-\varepsilon}) \cap C^1(\mathbb{R}_+; H_{\beta-1-\varepsilon}^{s-1-\varepsilon})$$

16
17
18 *for each $\varepsilon > 0$.*

19
20
21 Ukai's theory describes the evolution of number densities that are close to a uniform
22 Maxwellian state, and therefore, one could think of using it in the context of incompressible
23 hydrodynamic limits. However, one should bear in mind that the parameter η that monitors
24 the size of the initial number density fluctuation is not uniform in the Knudsen number, so
25 that applying Ukai's ideas to derive, say, the incompressible Navier–Stokes equations from
26 the Boltzmann equation requires nontrivial modifications, due to Bardos and Ukai [14].

27 Hence, for the purpose of deriving hydrodynamic models from the Boltzmann equation,
28 it is desirable to have at one's disposal a global existence theory based on a priori estimates
29 that are uniform in the Knudsen number. The only such existence theory so far is the
30 DiPerna–Lions theory of weak solutions of the Boltzmann equation that is described below.

31 32 33 8.2. The DiPerna–Lions theory

34
35 As already mentioned in our presentation of the Boltzmann equation, the collision integral
36 is local in t and x and an integral operator in v . In other words, the collision integral acts as
37 a multiplication operator in the variables (t, x) , and as kind of convolution in the variable v .

38 On the other hand, the natural a priori bound for the number density is

$$39 \quad F \in L_t^\infty(L \ln L_x).$$

40
41
42 For such an F , expressions like

$$43 \quad F^2 \quad \text{or} \quad F \int_{\mathbb{R}^3} F \, dv$$

44

45

are defined only as measurable functions, and not as distributions. Hence the collision integral does not define a distribution for all number densities that satisfy the natural a priori bounds on solutions of the Boltzmann equation.

To get around this, DiPerna and Lions proposed to use the following notion of solution.

DEFINITION 8.3. A nonnegative function $F \in C(\mathbb{R}_+; L^1(\mathbb{R}^3 \times \mathbb{R}^3))$ is a renormalized solution of the Boltzmann equation if

$$\frac{\mathcal{B}(F, F)}{\sqrt{1+F}} \in L^1_{\text{loc}}(dt dx dv),$$

and if, for each $\Gamma \in C^1(\mathbb{R}_+)$ such that

$$\Gamma'(Z) \leq \frac{C}{\sqrt{1+Z}} \quad \text{for all } Z \geq 0,$$

one has

$$(\partial_t + v \cdot \nabla_x) \Gamma(F) = \Gamma'(F) \mathcal{B}(F, F)$$

in the sense of distributions on $\mathbb{R}_+^* \times \mathbb{R}^3 \times \mathbb{R}^3$.

In this subsection, we shall consider collision kernels that satisfy the following weak cut-off assumption

$$\lim_{|v| \rightarrow +\infty} \frac{1}{1+|v|^2} \int_{|v_*| < R} \int_{\mathbb{S}^2} b(v-v_*, \omega) d\omega dv_* = 0 \quad \text{for each } R > 0. \quad (8.1)$$

The following result was proved by DiPerna and Lions in [37].

THEOREM 8.4. Let $F^{\text{in}} \equiv F(x, v) \geq 0$ a.e. on $\mathbb{R}^3 \times \mathbb{R}^3$ satisfy

$$\iint_{\mathbb{R}^3 \times \mathbb{R}^3} (1 + |x|^2 + |v|^2 + |\ln F^{\text{in}}|) F^{\text{in}} dx dv < +\infty.$$

Then, there exists a renormalized solution of the Boltzmann equation such that $F|_{t=0} = F^{\text{in}}$. This renormalized solution satisfies

- the continuity equation (local conservation of mass)

$$\partial_t \int_{\mathbb{R}^3} F dv + \operatorname{div}_x \int_{\mathbb{R}^3} v F dv = 0;$$

- the global conservation of momentum: for each $t \geq 0$,

$$\iint_{\mathbb{R}^3 \times \mathbb{R}^3} v F(t) dv dx = \iint_{\mathbb{R}^3 \times \mathbb{R}^3} v F^{\text{in}} dv dx;$$

- the energy inequality: for each $t \geq 0$,

$$\iint_{\mathbb{R}^3 \times \mathbb{R}^3} \frac{1}{2} |v|^2 F(t) \, dv \, dx \leq \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \frac{1}{2} |v|^2 F^{\text{in}} \, dv \, dx;$$

- and the H inequality: for each $t > 0$, one has

$$\begin{aligned} & \iint_{\mathbb{R}^3 \times \mathbb{R}^3} F \ln F(t) \, dv \, dx \\ & + \int_0^t \iint_{\mathbb{R}^3} \iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{R}^3} (F' F'_* - F F_*) \ln \left(\frac{F' F'_*}{F F_*} \right) b \, dv \, dv_* \, d\omega \, dx \, ds \\ & \leq \iint_{\mathbb{R}^3 \times \mathbb{R}^3} F^{\text{in}} \ln F^{\text{in}} \, dv \, dx. \end{aligned}$$

A complete description of the proof of the DiPerna–Lions theorem is beyond the scope of the present work. We shall just explain the main ideas in it.

8.2.1. The role of the normalizing nonlinearity. Assume that $F \equiv F(t, x, v) \geq 0$ a.e. on $\mathbb{R}_+ \times \mathbb{R}^3 \times \mathbb{R}^3$ satisfies

$$\iint_{\mathbb{R}^3 \times \mathbb{R}^3} (1 + |v|^2) F(t, x, v) \, dv \, dx \leq C \quad \text{for a.e. } t \geq 0, \quad (8.2)$$

and

$$\int_0^T \iint_{\mathbb{R}^3} \iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{R}^3} (F' F'_* - F F_*) \ln \left(\frac{F' F'_*}{F F_*} \right) b \, dv \, dv_* \, d\omega \, dx \, ds \leq C_T \quad (8.3)$$

for each $T > 0$. A solution to the Boltzmann equation satisfies the first inequality (by the global conservation law of energy) and the second estimate by the entropy production bound deduced from Boltzmann's H -theorem.

Then

$$\frac{|\mathcal{B}(F, F)|}{\sqrt{1 + F}} \in L^1_{\text{loc}}(\mathbb{R}_+ \times \mathbb{R}^3 \times \mathbb{R}^3).$$

Indeed, we first recall the elementary inequality

$$(\sqrt{X} - \sqrt{Y})^2 \leq \frac{1}{4} (X - Y) (\ln X - \ln Y) \quad \text{for each } X, Y > 0.$$

Then

$$\begin{aligned} |\sqrt{F' F'_*} - \sqrt{F F_*}| &= (\sqrt{F' F'_*} + \sqrt{F F_*}) |\sqrt{F' F'_*} - \sqrt{F F_*}| \\ &\leq (\sqrt{F' F'_*} - \sqrt{F F_*})^2 + 2\sqrt{F F_*} |\sqrt{F' F'_*} - \sqrt{F F_*}|. \end{aligned}$$

Hence

$$\begin{aligned}
& \int_0^T \int_{\mathbb{R}^3} \int_{|v| < R} \frac{|\mathcal{B}(F, F)|}{\sqrt{1+F}} dv dx dt \\
& \leq \int_0^T \int_{\mathbb{R}^3} \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \iint_{\mathbb{R}^3 \times \mathbb{R}^3} (\sqrt{F'F'_*} - \sqrt{FF_*})^2 b dv dv_* d\omega dx ds \\
& \quad + 2 \left(\int_0^T \int_{\mathbb{R}^3} \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \iint_{\mathbb{R}^3 \times \mathbb{R}^3} (\sqrt{F'F'_*} - \sqrt{FF_*})^2 b dv dv_* d\omega dx ds \right)^{1/2} \\
& \quad \times \left(\int_0^T \iint_{\mathbb{R}^3 \times \mathbb{R}^3} F_* \left(\iint_{|v| < R, |\omega|=1} b(v - v_*, \omega) d\omega dv \right) dv_* dx ds \right)^{1/2}.
\end{aligned}$$

By using the entropy production bound (8.3) and the weak cut-off assumption above on the collision kernel b , this inequality becomes

$$\begin{aligned}
& \int_0^T \int_{\mathbb{R}^3} \int_{|v| < R} \frac{|\mathcal{B}(F, F)|}{\sqrt{1+F}} dv dx dt \\
& \leq C_T + 2\sqrt{C_T} \\
& \quad \times \left(\int_0^T \iint_{\mathbb{R}^3 \times \mathbb{R}^3} F_* \left(\iint_{|v| < R, |\omega|=1} b(v - v_*, \omega) d\omega dv \right) dv_* dx ds \right)^{1/2} \\
& \leq C_T + 2\sqrt{C_T} \left(\int_0^T \iint_{\mathbb{R}^3 \times \mathbb{R}^3} F_* O(1 + |v_*|^2) dv_* dx ds \right)^{1/2} \\
& \leq C_T + 2\sqrt{C_T O(C)}.
\end{aligned}$$

In the sequel, we restrict our attention for simplicity to the case of bounded collision kernels

$$0 < b(z, \omega) \leq C_b, \quad (z, \omega) \in \mathbb{R}^3 \times \mathbb{S}^2. \quad (8.4)$$

8.2.2. Compactness by velocity averaging. A renormalized solution of the Boltzmann equation is obtained as a limit point of the sequence F_n of solutions to the regularized Boltzmann equation

$$\begin{aligned}
\partial_t F_n + v \cdot \nabla_x F_n &= \frac{\mathcal{B}(F_n, F_n)}{1 + (1/n) \int_{\mathbb{R}^3} F_n dv}, \\
F_n|_{t=0} &= F^{\text{in}}
\end{aligned} \quad (8.5)$$

in the weak topology of $L^1_{\text{loc}}(\mathbb{R}_+; L^1(\mathbb{R}^3 \times \mathbb{R}^3))$. As in the case of all nonlinear PDEs, the main difficulty is to pass to the limit in the nonlinear term – here, in the collision integral. This requires a class of compactness results on the Boltzmann equation that are somehow adapted to the collision integral. In particular, since the collision integral acts as a

convolution operator in the variable v and a multiplication operator in the variables t and x , one should seek compactness “with respect to the t and x variables only” – a notion which, of course, remains to be defined.

The appropriate class of compactness theorems was discovered a few years before being applied to the Boltzmann equation. They are known as “velocity averaging results”, and were first introduced by Golse, Perthame and Sentis in [56] within the context of the diffusion approximation of the neutron transport equation. (Independently, analogous regularity results for the coefficients of the spherical harmonic expansion of the solution to the free transport equation were announced in [1].)

Obviously, whatever compactness in the strong topology of L^1_{loc} is to be found on a sequence of solutions to the Boltzmann equation (regularized or not) has to come from the streaming (free transport) operator $\partial_t + v \cdot \nabla_x$.

Being hyperbolic, the transport operator $v \cdot \nabla_x$ propagates singularities along characteristics. Therefore, at first sight it seems hopeless that one might obtain any regularizing effect from the free streaming part of the Boltzmann equation, or of any other similar kinetic model. One can think of the following elementary example.

EXAMPLE. Let $f \in L^2(\mathbb{R})$; for a.e. $x, v \in \mathbb{R}^2$, define $F(x, v) = f(v_2 x_1 - v_1 x_2)$. Clearly, $F \in L^2_{\text{loc}}(\mathbb{R}^2 \times \mathbb{R}^2)$ and $v \cdot \nabla_x F = 0$. However, since f can be any function in $L^2(\mathbb{R})$, $F \notin H^s_{\text{loc}}(\mathbb{R}^2 \times \mathbb{R}^2)$ for any $s > 0$, although $v \cdot \nabla_x F \in L^2_{\text{loc}}(\mathbb{R}^2 \times \mathbb{R}^2)$.

The key to obtaining regularizing effects from the transport operator $v \cdot \nabla_x$ is to seek those effects not on the number density itself, but on velocity averages thereof, in other words, on the macroscopic densities. Here is the prototype of all velocity averaging results.

THEOREM 8.5 (Golse, Perthame and Sentis [56]). *Let F_ε be a bounded family in $L^2(\mathbb{R}^D \times \mathbb{R}^D)$. Assume that the family $v \cdot \nabla_x F_\varepsilon$ is also bounded in $L^2(\mathbb{R}^D \times \mathbb{R}^D)$. Then, for each $\phi \in L^2(\mathbb{R}^D)$, the family of moments ρ_ε defined by*

$$\rho_\varepsilon(x) = \int_{\mathbb{R}^D} F_\varepsilon(x, v) \phi(v) \, dv$$

is relatively compact in $L^2_{\text{loc}}(\mathbb{R}^D)$.

PROOF. Set $G_\varepsilon = F_\varepsilon + v \cdot \nabla_x F_\varepsilon$, and let \widehat{F}_ε and \widehat{G}_ε denote respectively the Fourier transforms of F_ε and G_ε in the x variable only. The assumptions on F_ε imply that the family G_ε is bounded in $L^2(\mathbb{R}^D \times \mathbb{R}^D)$, and that

$$\widehat{\rho}_\varepsilon(\xi) = \int_{\mathbb{R}^D} \frac{\widehat{G}_\varepsilon(\xi, v) \phi(v) \, dv}{1 + i v \cdot \xi}.$$

We need to study how $\widehat{\rho}_\varepsilon(\xi)$ decays for $|\xi|$ large. By the Cauchy–Schwarz inequality

$$|\widehat{\rho}_\varepsilon(\xi)|^2 \leq \frac{1}{m(\xi)} \int_{\mathbb{R}^D} |\widehat{G}_\varepsilon(\xi, v)|^2 \, dv,$$

1 with

$$2 \quad \frac{1}{m(\xi)} := \int_{\mathbb{R}^D} \frac{|\phi(v)|^2 dv}{1 + |v \cdot \xi|^2} = \int_{\mathbb{R}^D} \frac{|\phi(v)|^2 dv}{1 + |v \cdot \omega|^2 |\xi|^2},$$

3 where $\omega = \xi/|\xi| \in \mathbb{S}^{D-1}$ for all $\xi \in \mathbb{R}^D \setminus \{0\}$. The latter integral is a decreasing family indexed by $|\xi|$ of continuous functions of ω ; this family vanishes pointwise in ω as $|\xi| \rightarrow +\infty$ by dominated convergence. By Dini's theorem, it vanishes uniformly in $\omega \in \mathbb{S}^{D-1}$, and therefore $m(\xi) \rightarrow +\infty$ as $|\xi| \rightarrow +\infty$. Since the family

$$4 \quad \int_{\mathbb{R}^D} |\hat{\rho}_\varepsilon(\xi)|^2 m(\xi) d\xi \leq \iint_{\mathbb{R}^D \times \mathbb{R}^D} |\widehat{G}_\varepsilon(\xi, v)|^2 d\xi dv$$

5 is bounded by Plancherel's theorem, ρ_ε is relatively compact in $L^2_{\text{loc}}(\mathbb{R}^D)$ (by a variant of Rellich's compactness theorem). \square

6 Since the operator $(I + v \cdot \nabla_x)^{-1}$ (which maps G on the solution F of $F + v \cdot \nabla_x F = G$) is a contraction mapping on both $L^1(\mathbb{R}^D \times \mathbb{R}^D)$ and $L^\infty(\mathbb{R}^D \times \mathbb{R}^D)$, the velocity averaging result above also holds in L^p for all $p \in (1, +\infty)$ by interpolation. However, it fails in L^1 , as the following example shows. (It also fails in L^∞ , see [55], p. 124.)

7 EXAMPLE ([55], pp. 123–124). Consider G_ε , a bounded family of $L^1(\mathbb{R}^D \times \mathbb{R}^D)$, and for each ε , let F_ε be the solution to $F_\varepsilon + v \cdot \nabla_x F_\varepsilon = G_\varepsilon$. Assume that $G_\varepsilon \rightarrow \delta_0 \otimes \delta_{v^*}$ weakly, where $|v^*| = 1$. Then both F_ε and $v \cdot \nabla_x F_\varepsilon$ are bounded in $L^1(\mathbb{R}^D \times \mathbb{R}^D)$ and

$$8 \quad \rho_\varepsilon(x) = \int_{\mathbb{R}^D} F_\varepsilon(x, v) dv = \int_0^{+\infty} \int_{\mathbb{R}^D} e^{-t} G_\varepsilon(x - tv, v) dv dt$$

9 so that, for any test function $\phi \in C_c(\mathbb{R}^D)$,

$$10 \quad \int_{\mathbb{R}^D} \rho_\varepsilon(x) \phi(x) dx \rightarrow \int_0^{+\infty} e^{-t} \phi(tv^*) dt$$

11 as $\varepsilon \rightarrow 0$. Hence ρ_ε converges weakly to a density carried by the half-line $\mathbb{R}_+ v^*$, so that in particular the family ρ_ε is not relatively compact in $L^1_{\text{loc}}(\mathbb{R}^D)$.

12 This example rests on the possible build-up of concentrations in F_ε and $v \cdot \nabla_x F_\varepsilon$. If such concentrations are ruled out, the same interpolation argument as above entails the following L^1 variant of velocity averaging.

13 PROPOSITION 8.6 (Golse, Lions, Perthame and Sentis [55]). *Let F_ε be a family of measurable functions on $\mathbb{R}^D \times \mathbb{R}^D$ such that, for each compact subset K of \mathbb{R}^D , both families F_ε and $v \cdot \nabla_x F_\varepsilon$ are uniformly integrable on $K \times \mathbb{R}^D$. Then the family ρ_ε defined by*

$$14 \quad \rho_\varepsilon(x) = \int_{\mathbb{R}^D} F_\varepsilon(x, v) dv$$

1 is relatively compact in $L^1_{\text{loc}}(\mathbb{R}^D)$. 1

2
3 PROOF. Let $\chi \equiv \chi(x)$ belongs to $C_c^\infty(\mathbb{R}^D)$. Set 3

$$4 \quad G_\varepsilon(x, v) = \chi(x)F_\varepsilon(x, v) + v \cdot \nabla_x(\chi(x)F_\varepsilon(x, v)) \quad 4$$

5 and, for each $\lambda > 0$, decompose $\chi\rho_\varepsilon$ as follows 5

$$6 \quad \chi\rho_\varepsilon = \chi\rho_{\lambda, \varepsilon}^> + \chi\rho_{\lambda, \varepsilon}^< \quad 6$$

7 with 7

$$8 \quad \chi\rho_{\lambda, \varepsilon}^> = \int_{|v| \leq \lambda} (I + v \cdot \nabla_x)^{-1} (G_\varepsilon \mathbb{1}_{|G_\varepsilon| > \lambda}) \, dv + \int_{|v| > \lambda} \chi F_\varepsilon \, dv, \quad 8$$

$$9 \quad \chi\rho_{\lambda, \varepsilon}^< = \int_{|v| \leq \lambda} (I + v \cdot \nabla_x)^{-1} (G_\varepsilon \mathbb{1}_{|G_\varepsilon| \leq \lambda}) \, dv. \quad 9$$

10
11 The assumptions on F_ε imply that G_ε is uniformly integrable, so that $G_\varepsilon \mathbb{1}_{|G_\varepsilon| > \lambda} \rightarrow 0$ and $\chi F_\varepsilon \mathbb{1}_{|v| > \lambda} \rightarrow 0$ in $L^1(\mathbb{R}^D \times \mathbb{R}^D)$ uniformly in ε as $\lambda \rightarrow +\infty$; thus $\chi\rho_{\lambda, \varepsilon}^> \rightarrow 0$ in $L^1(\mathbb{R}^D)$ uniformly in ε as $\lambda \rightarrow +\infty$. 10

11 On the other hand, for each λ , the family $G_\varepsilon \mathbb{1}_{|G_\varepsilon| > \lambda}$ indexed by ε is bounded in $L^2(\mathbb{R}^D)$; 11
12 thus, by velocity averaging in L^2 , $\chi\rho_{\lambda, \varepsilon}^<$ is relatively compact in $L^2(\mathbb{R}^D)$ – and therefore 12
13 in $L^1(\mathbb{R}^D)$, since it has support in $\text{supp } \chi$ which is compact. 13

14 Hence $\chi\rho_\varepsilon$ is relatively compact in $L^1(\mathbb{R}^D)$. Since χ is arbitrary, this eventually implies 14
15 that ρ_ε is relatively compact in $L^1_{\text{loc}}(\mathbb{R}^D)$. \square 15

16 In fact, by a further interpolation argument, one can get rid of the assumption of uniform 16
17 integrability on derivatives. 17

18 THEOREM 8.7 (Golse and Saint-Raymond [60]). Let F_ε be a family of measurable func- 18
19 tions on $\mathbb{R}^D \times \mathbb{R}^D$ such that, for each compact subset K of \mathbb{R}^D , the family F_ε is uniformly 19
20 integrable on $K \times \mathbb{R}^D$, while $v \cdot \nabla_x F_\varepsilon$ is bounded in $L^1(K \times \mathbb{R}^D)$. Then the family ρ_ε 20
21 defined by 21

$$22 \quad \rho_\varepsilon(x) = \int_{\mathbb{R}^D} F_\varepsilon(x, v) \, dv \quad 22$$

23 is relatively compact in $L^1_{\text{loc}}(\mathbb{R}^D)$. 23

24 PROOF. Without loss of generality, assume that all the F_ε are supported in $K \times \mathbb{R}^D$. Write 24

$$25 \quad \rho_\varepsilon(x) = \lambda \int_{\mathbb{R}^D} (\lambda I + v \cdot \nabla_x)^{-1} F_\varepsilon(x, v) \, dv \quad 25$$

$$26 \quad + \int_{\mathbb{R}^D} (\lambda I + v \cdot \nabla_x)^{-1} (v \cdot \nabla_x F_\varepsilon)(x, v) \, dv. \quad 26$$

1 Since

$$2 \quad 3 \quad 4 \quad \left\| (\lambda I + v \cdot \nabla_x)^{-1} \right\|_{\mathcal{L}(L^1_{x,v})} \leq \frac{1}{\lambda},$$

5 the second term on the right-hand side of the equality above is $O(1/\lambda)$ in L^1_x uniformly
6 in ε , while the first term is strongly relatively compact in L^1_x for each $\lambda > 0$ by the previous
7 proposition. Hence the family ρ_ε is strongly relatively compact in L^1_x . \square

9 There are several extensions and variants of the velocity averaging results recalled here,
10 see for instance [36,38,47–49,106]. Except for the extension of the above results to the
11 evolution problem, which is needed in the construction of renormalized solutions to the
12 Boltzmann equation, we shall not discuss these extensions in the present notes, but refer
13 the interested reader to Chapter 1 of [18] for a survey of that theory as of 2000.

14 Here is the analogue of the L^1 -variant of velocity averaging for evolution problems.

15
16 **THEOREM 8.8.** *Consider $F_\varepsilon \equiv F_\varepsilon(t, x, v)$, a family of measurable functions on $\mathbb{R}_+ \times$
17 $\mathbb{R}^D \times \mathbb{R}^D$ such that, for each $T > 0$ and each compact $K \subset \mathbb{R}^D$, F_ε is uniformly integrable
18 on $[0, T] \times K \times \mathbb{R}^D$ while $(\partial_t + v \cdot \nabla_x)F_\varepsilon$ is bounded in $L^1([0, T] \times K \times \mathbb{R}^D)$. Then the
19 family ρ_ε defined by*

$$20 \quad 21 \quad 22 \quad \rho_\varepsilon(t, x) = \int_{\mathbb{R}^D} F_\varepsilon(t, x, v) \, dv$$

23
24 is relatively compact in $L^1_{\text{loc}}(\mathbb{R}_+ \times \mathbb{R}^D)$.

25
26 The proof is a straightforward variant of the arguments for the steady transport operator
27 $v \cdot \nabla_x$ given above.

28
29 **8.2.3. Conclusion.** Let us briefly explain how the renormalization procedure is combined
30 with compactness by velocity averaging in the proof of the DiPerna–Lions theorem.

31 Choose as normalizing nonlinearity the function

$$32 \quad 33 \quad 34 \quad \Gamma_\delta(Z) = \frac{1}{\delta} \ln(1 + \delta Z), \quad \delta > 0,$$

35 and consider the truncated Boltzmann equation (8.5). We leave it to the reader to verify
36 that, under the condition (8.4), the map

$$37 \quad 38 \quad 39 \quad F \mapsto \frac{\mathcal{B}(F, F)}{1 + (1/n) \int_{\mathbb{R}^3} F \, dv}$$

40
41 is Lipschitz continuous on $L^1(\mathbb{R}_x^3 \times \mathbb{R}_v^3)$, so that the truncated Boltzmann equation (8.5)
42 has a global solution. Because the truncation factor

$$43 \quad 44 \quad 45 \quad \frac{1}{1 + (1/n) \int_{\mathbb{R}^3} F \, dv}$$

is independent of v , the symmetries of the collision integral that imply the local conservation laws of mass, momentum and energy, and the H -theorem hold also for the truncated collision integral

$$\frac{\mathcal{B}(F, F)}{1 + (1/n) \int_{\mathbb{R}^3} F \, dv}$$

so that, in particular,

$$\begin{aligned} & \iint_{\mathbb{R}^3 \times \mathbb{R}^3} (1 + |x - tv|^2 + |v|^2) F_n(t, x, v) \, dx \, dv \\ &= \iint_{\mathbb{R}^3 \times \mathbb{R}^3} (1 + |x|^2 + |v|^2) F^{\text{in}}(x, v) \, dx \, dv \end{aligned}$$

for all $t \geq 0$, while

$$\iint_{\mathbb{R}^3 \times \mathbb{R}^3} F_n \ln F_n(t, x, v) \, dx \, dv \leq \iint_{\mathbb{R}^3 \times \mathbb{R}^3} F^{\text{in}} \ln F^{\text{in}}(x, v) \, dx \, dv.$$

As explained in Case 4 of Section 3.3, this implies the existence of a positive constant C such that

$$\begin{aligned} & \iint_{\mathbb{R}^3 \times \mathbb{R}^3} (1 + |x|^2 + |v|^2) F_n(t, x, v) \, dx \, dv \leq C(1 + t^2) \quad \text{and} \\ & \iint_{\mathbb{R}^3 \times \mathbb{R}^3} F_n |\ln F_n|(t, x, v) \, dx \, dv \leq C(1 + t^2) \quad \text{for all } t \geq 0. \end{aligned} \tag{8.6}$$

Hence F_n is weakly relatively compact in $L^1_{\text{loc}}(\mathbb{R}_+ \times \mathbb{R}^3 \times \mathbb{R}^3)$ by the Dunford–Pettis theorem. Since, for each $\delta > 0$, one has

$$\Gamma_\delta(Z) \leq Z \quad \text{for each } Z \geq 0,$$

the sequence of functions $\Gamma_\delta(F_n)$ is also weakly relatively compact in $L^1_{\text{loc}}(\mathbb{R}_+; L^1(\mathbb{R}^3 \times \mathbb{R}^3))$. On the other hand, the discussion in Section 8.2.1 shows that

$$(\partial_t + v \cdot \nabla_x) \Gamma_\delta(F_n) = \frac{\mathcal{B}(F_n, F_n)}{(1 + \delta F_n)(1 + (1/n) \int_{\mathbb{R}^3} F_n \, dv)}$$

is bounded in $L^1_{\text{loc}}(\mathbb{R}_+ \times \mathbb{R}^3 \times \mathbb{R}^3)$. By velocity averaging (Theorem 8.8), this implies that the sequence

$$\int_{\mathbb{R}^3} \Gamma_\delta(F_n) \, dv$$

1 is strongly relatively compact in $L^1_{\text{loc}}(\mathbb{R}_+; L^1(\mathbb{R}^3))$. On the other hand, because of the
 2 bound (8.6),

$$\int_{\mathbb{R}^3} F_n \, dv - \int_{\mathbb{R}^3} \Gamma_\delta(F_n) \, dv \rightarrow 0$$

7 in $L^1_{\text{loc}}(\mathbb{R}_+; L^1(\mathbb{R}^3))$ as $\delta \rightarrow 0$, uniformly in $n \geq 1$. Hence we conclude that the sequence

$$\int_{\mathbb{R}^3} F_n \, dv$$

12 is strongly relatively compact in $L^1_{\text{loc}}(\mathbb{R}_+; L^1(\mathbb{R}^3))$.

Hence, modulo extraction of a subsequence

$$F_n \rightarrow F \quad \text{weakly in } L^1_{\text{loc}}(\mathbb{R}_+ \times \mathbb{R}^3 \times \mathbb{R}^3)$$

17 while

$$\int_{\mathbb{R}^3} F_n \, dv \rightarrow \int_{\mathbb{R}^3} F \, dv \quad \text{strongly in } L^1_{\text{loc}}(\mathbb{R}_+ \times \mathbb{R}^3).$$

22 Since the collision integral acts as a convolution in the v -variable and a multiplication op-
 23 erator in the t and x variables, this compactness theorem implies that, for each $\phi \in C_c(\mathbb{R}^3)$,

$$\int_{\mathbb{R}^3} \mathcal{B}(F_n, F_n) \phi \, dv \rightarrow \int_{\mathbb{R}^3} \mathcal{B}(F, F) \phi \, dv \quad \text{in measure on } [0, T] \times K$$

28 for each $T > 0$ and each compact $K \subset \mathbb{R}^3$. At first sight, this is not enough to pass to the
 29 limit in the sense of distributions in both sides of the truncated Boltzmann equation (8.5).

30 Instead, one integrates the truncated Boltzmann equation along characteristics by treat-
 31 ing the gain term in the truncated collision integral as a source term. One easily sees that
 32 the limit F is a supersolution of the limiting Boltzmann equation integrated along charac-
 33 teristics; notice that this does not make use of the renormalization procedure. That F is a
 34 subsolution is more involved, we refer to [37] for a complete proof.

37 8.3. Variants of the DiPerna–Lions theory

38
 39 The original DiPerna–Lions theorem considers a cloud of gas that expands in the vacuum,
 40 without any restriction on its degree of rarefaction, i.e., for an initial number density of
 41 arbitrary size that has finite mass, energy, second moment in x and entropy. As explained
 42 in Section 8.1.1, this kind of situation is not compatible with incompressible hydrodynamic
 43 limits.

44 For that purpose, we describe below two variants of the DiPerna–Lions theory that are
 45 particularly relevant in the context of incompressible hydrodynamic limits.

8.3.1. The periodic box. The first variant of the DiPerna–Lions theory that we discuss here is the case of the spatial domain \mathbb{T}^3 (the periodic box). The collision kernel b satisfies the weak cut-off condition (8.1). Let M be the centered reduced Gaussian

$$M(v) = \frac{1}{(2\pi)^{3/2}} e^{-|v|^2/2}.$$

We shall say that $F \in C(\mathbb{R}_+; L^1(\mathbb{R}^3 \times \mathbb{R}^3))$ is a *renormalized solution* of the Boltzmann equation

$$\begin{aligned} \partial_t F + v \cdot \nabla_x F &= \mathcal{B}(F, F), \quad (t, x, v) \in \mathbb{R}_+^* \times \mathbb{T}^3 \times \mathbb{R}^3, \\ F|_{t=0} &= F^{\text{in}}, \end{aligned}$$

relative to M if and only if, for each normalizing nonlinearity $\Gamma \in C^1(\mathbb{R}_+)$ such that

$$\Gamma'(Z) \leq \frac{C}{\sqrt{1+Z}}, \quad Z \geq 0,$$

one has

$$M(\partial_t + v \cdot \nabla_x) \Gamma\left(\frac{F}{M}\right) = \Gamma'\left(\frac{F}{M}\right) \mathcal{B}(F, F)$$

in the sense of distributions on $\mathbb{R}_+^* \times \mathbb{T}^3 \times \mathbb{R}^3$.

THEOREM 8.9. *Let $F^{\text{in}} \geq 0$ a.e. be a measurable function such that $H(F^{\text{in}}|M) < +\infty$. There exists a renormalized solution F relative to M of the Cauchy problem for the Boltzmann equation with initial data F^{in} . This solution satisfies*

- the continuity equation

$$\partial_t \int_{\mathbb{R}^3} F \, dv + \operatorname{div}_x \int_{\mathbb{R}^3} v F \, dv = 0;$$

- the following variant of the local conservation of momentum

$$\partial_t \int_{\mathbb{R}^3} v F \, dv + \operatorname{div}_x \int_{\mathbb{R}^3} v \otimes v F \, dv + \operatorname{div}_x m = 0,$$

where $m \in L^\infty(\mathbb{R}_+; \mathcal{M}(\mathbb{T}^3, M_3(\mathbb{R})))$ with values in nonnegative symmetric matrices;

- the following energy relation

$$\begin{aligned} & \iint_{\mathbb{T}^3 \times \mathbb{R}^3} \frac{1}{2} |v|^2 F(t, x, v) \, dx \, dv + \frac{1}{2} \iint_{\mathbb{T}^3 \times \mathbb{R}^3} \operatorname{trace}(m(t)) \\ &= \iint_{\mathbb{T}^3 \times \mathbb{R}^3} \frac{1}{2} |v|^2 F^{\text{in}}(x, v) \, dx \, dv \end{aligned}$$

1 for each $t > 0$;

- 2 • and the H inequality: for each $t > 0$, one has

$$\begin{aligned} & \frac{1}{4} \int_0^t \int_{\mathbb{T}^3} \int \int \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{R}^3} (F' F'_* - F F_*') \ln \left(\frac{F' F'_*}{F F_*'} \right) b \, dv \, dv_* \, d\omega \, dx \, ds \\ & \leq H(F^{\text{in}} | M) - H(F(t) | M) - \frac{1}{2} \int \text{trace}(m(t)). \end{aligned}$$

3
4
5
6
7
8
9
10 The above result – especially the existence of the defect measure m – is due to Lions
11 and Masmoudi [88].

12 **8.3.2. The Euclidean space with uniform Maxwellian state at infinity.** The next variant
13 of the DiPerna–Lions theory that we consider is the case of a spatial domain that is the
14 Euclidean space with uniform Maxwellian equilibrium at infinity. Consider the Cauchy
15 problem
16

$$\begin{aligned} & \partial_t F + v \cdot \nabla_x F = \mathcal{B}(F, F), \quad (t, x, v) \in \mathbb{R}_+^* \times \mathbb{R}^3 \times \mathbb{R}^3, \\ & F(t, x, v) \rightarrow M \quad \text{as } |x| \rightarrow +\infty, \\ & F|_{t=0} = F^{\text{in}}, \end{aligned}$$

17
18
19
20
21
22 where M is the centered reduced Gaussian

$$M(v) = \frac{1}{(2\pi)^{3/2}} e^{-|v|^2/2}.$$

23
24
25 Here again, the collision kernel b satisfies the weak cut-off assumption (8.1). The notion
26 of “renormalized solution relative to M ” of the Cauchy problem above is the same as in
27 the case of the periodic box.

28
29
30
31 **THEOREM 8.10 (Lions [83]).** Let $F^{\text{in}} \geq 0$ a.e. be a measurable function such that
32 $H(F^{\text{in}} | M) < +\infty$. There exists a renormalized solution F relative to M of the Cauchy
33 problem for the Boltzmann equation with initial data F^{in} . This solution satisfies
34

- 35 • the continuity equation

$$\partial_t \int_{\mathbb{R}^3} F \, dv + \text{div}_x \int_{\mathbb{R}^3} v F \, dv = 0;$$

- 36
37
38
39
40 • and the H -inequality: for each $t > 0$, one has

$$\begin{aligned} & \frac{1}{4} \int_0^t \int_{\mathbb{R}^3} \int \int \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{R}^3} (F' F'_* - F F_*') \ln \left(\frac{F' F'_*}{F F_*'} \right) b \, dv \, dv_* \, d\omega \, dx \, ds \\ & \leq H(F^{\text{in}} | M) - H(F(t) | M). \end{aligned}$$

9. The Hilbert expansion method: Application to the compressible Euler limit

In this section we describe a first method for deriving hydrodynamic models from the Boltzmann equation. In spite of its numerous shortcomings (which we shall discuss at the end of the present section), this method is extremely robust, and can be applied to various kinetic models other than the Boltzmann equation.

Start from the Boltzmann equation in the compressible Euler scaling

$$\partial_t F_\varepsilon + v \cdot \nabla_x F_\varepsilon = \frac{1}{\varepsilon} \mathcal{B}(F_\varepsilon, F_\varepsilon), \quad (t, x, v) \in \mathbb{R}_+^* \times \mathbb{T}^3 \times \mathbb{R}^3. \quad (9.1)$$

For simplicity, we only consider in this section the case of a hard sphere gas, so that the collision kernel b in Boltzmann's collision integral \mathcal{B} is given by the expression

$$b(z, \omega) = |z \cdot \omega|, \quad (z, \omega) \in \mathbb{R}^3 \times \mathbb{S}^2.$$

Consider next the compressible Euler system for a perfect monatomic gas

$$\begin{aligned} \partial_t \rho + \operatorname{div}_x(\rho u) &= 0, \\ \partial_t(\rho u) + \operatorname{div}_x(\rho u \otimes u) + \nabla_x(\rho \theta) &= 0, \\ \partial_t \left(\rho \left(\frac{1}{2} |u|^2 + \frac{3}{2} \theta \right) \right) + \operatorname{div}_x \left(\rho u \left(\frac{1}{2} |u|^2 + \frac{5}{2} \theta \right) \right) &= 0, \\ (\rho, u, \theta)|_{t=0} &= (\rho^{\text{in}}, u^{\text{in}}, \theta^{\text{in}}), \end{aligned} \quad (9.2)$$

where

$$\rho^{\text{in}}, u^{\text{in}} \text{ and } \theta^{\text{in}} \in H^5(\mathbb{T}^3), \quad \rho^{\text{in}} \geq 0 \text{ and } \theta^{\text{in}} > 0 \text{ on } \mathbb{T}^3. \quad (9.3)$$

Let (ρ, u, θ) be the solution to (9.2) predicted by Theorem 7.8 under the assumption (9.3), and call $T > 0$ its lifespan. Finally, define the local Maxwellian

$$E(t, x, v) = \mathcal{M}_{(\rho(t,x), u(t,x), \theta(t,x))}.$$

THEOREM 9.1 (Caffisch [22]). *There exists ε_0 such that, for each $\varepsilon \in (0, \varepsilon_0)$, there is a unique solution F_ε of the Boltzmann equation (9.1) on $[0, T] \times \mathbb{T}^3 \times \mathbb{R}^3$ satisfying the estimate*

$$\sup_{0 \leq t \leq T'} \|F(t, \cdot, \cdot) - E(t, \cdot, \cdot)\|_{L^2(\mathbb{T}^3 \times \mathbb{R}^3)} = O(\varepsilon) \quad \text{as } \varepsilon \rightarrow 0^+$$

for each $T' < T$.

We shall not give a complete proof of this result, which is fairly technical. Instead, we simply discuss below the main ideas in it.

1 First, the solution F_ε is sought as a truncated Hilbert expansion, plus a remainder term 1

$$2 \quad F_\varepsilon(t, x, v) = \sum_{k=0}^6 \varepsilon^k F_k(t, x, v) + \varepsilon^3 R_\varepsilon(t, x, v), \quad (9.4) \quad 3$$

4 such that the last term in the truncated expansion satisfies 5

$$6 \quad \int_{\mathbb{R}^3} \begin{pmatrix} 1 \\ v \\ |v|^2 \end{pmatrix} F_6 dv = 0. \quad (9.5) \quad 7$$

8 We recall from our discussion in Section 5.1 that the projection of a term F_k in Hilbert's 8
 9 expansion on the space of collision invariants is determined by postulating the existence of 9
 10 the next term in that expansion, i.e., F_{k+1} . In the truncated expansion above, we obviously 10
 11 do not postulate the existence of a F_7 , so that there is a certain amount of arbitrariness in 11
 12 the choice of F_6 , which is resolved by condition (9.5). The other terms F_k , $k = 0, \dots, 5$, 12
 13 are computed as in Section 5.1. 13
 14

15 Next we write an equation for the remainder R_ε : inserting the right-hand side of (9.4) in 15
 16 the scaled Boltzmann equation (9.1) we arrive at 16
 17

$$18 \quad (\partial_t + v \cdot \nabla_x) R_\varepsilon = \frac{2}{\varepsilon} \mathcal{B}(F_0, R_\varepsilon) + 2\mathcal{B}\left(\sum_{k=1}^6 \varepsilon^{k-1} F_k, R_\varepsilon\right) + \varepsilon^2 \mathcal{B}(R_\varepsilon, R_\varepsilon) \quad 18$$

$$19 \quad + \sum_{k+l \geq 7} \varepsilon^{k+l-4} \mathcal{B}(F_k, F_l) - \varepsilon^3 (\partial_t + v \cdot \nabla_x) F_6. \quad (9.6) \quad 19$$

20 Indeed, for $k = 0, \dots, 5$, the terms F_k are chosen as explained in Section 5.1, so that 20
 21

$$22 \quad (\partial_t + v \cdot \nabla_x) \sum_{k=0}^5 \varepsilon^k F_k = \sum_{l+m \leq 6} \varepsilon^{l+m-1} \mathcal{B}(F_l, F_m). \quad 22$$

23 Notice that, in (9.6), the nonlinear term is multiplied by an ε^2 factor: hence (9.6) is a 23
 24 weakly nonlinear equation. However, the linear term in (9.6) is 24
 25

$$26 \quad \frac{2}{\varepsilon} \mathcal{B}(F_0, R_\varepsilon) \quad 26$$

27 at leading order, which defines a nonpositive operator in a weighted L^2 space in the 27
 28 v -variable. Specifically, the operator $R \mapsto \mathcal{B}(F_0, R)$ is self-adjoint in $L^2(\mathbb{R}^3; F_0^{-1} dv)$. 28
 29 A slight difficulty in this setting is that the weight F_0^{-1} depends on (t, x) . 29
 30

31 To avoid this, we use a slightly different definition of the linearized collision integral 31
 32 than in Section 3.6. Define 32
 33

$$34 \quad \mathbf{L}_M \phi = -2M^{-1/2} \mathcal{B}(M, M^{1/2} \phi), \quad 34$$

where M is a Maxwellian density. As in Section 3.6, the operator \mathbf{L}_M is split into

$$\mathcal{L}_M \phi(v) = a(|v|)\phi(v) - \mathbf{K}_M \phi(v),$$

where

$$a_M(|v|) = \iint_{\mathbb{R}^3 \times \mathbb{S}^2} |(v - v_*) \cdot \omega| M_* dv_* d\omega$$

and

$$\mathbf{K}_M \phi(v) = \iint_{\mathbb{R}^3 \times \mathbb{S}^2} (\sqrt{M} \phi_* - \sqrt{M'_*} \phi' - \sqrt{M''_*} \phi''_*) |(v - v_*) \cdot \omega| \sqrt{M_*} dv_* d\omega.$$

The properties of \mathbf{L}_M are summarized in the following theorem.

THEOREM 9.2. *The operator \mathbf{K}_M is compact on $L^2(\mathbb{R}^3)$, while $a_M(|v|) \sim a^*|v|$ as $|v| \rightarrow +\infty$. Hence the operator \mathbf{L}_M is an unbounded self-adjoint Fredholm operator on $L^2(\mathbb{R}^3)$ with domain $L^2(\mathbb{R}^3; (1 + |v|) dv)$ and nullspace*

$$\text{Ker } \mathbf{L}_M = \text{span}\{\sqrt{M}, \sqrt{M}v_1, \sqrt{M}v_2, \sqrt{M}v_3, \sqrt{M}|v|^2\}.$$

The advantage in using \mathbf{L}_M instead of the operator $R \mapsto \mathcal{B}(F_0, R)$ is that the former operator is self-adjoint on the unweighted space $L^2(\mathbb{R}^3)$.

Formulating (9.6) in terms of the operator \mathbf{L}_M leads to an equation of the form

$$\begin{aligned} & (\partial_t + v \cdot \nabla_x)(E^{-1/2} R_\varepsilon) + \frac{1}{\varepsilon} \mathbf{L}_E(E^{-1/2} R_\varepsilon) \\ &= \varepsilon^2 \mathbf{Q}_E(E^{-1/2} R_\varepsilon, E^{-1/2} R_\varepsilon) + 2E^{-1/2} \mathcal{B}\left(\sum_{k=1}^6 \varepsilon^{k-1} F_k, R_\varepsilon\right) \\ &+ \sum_{k+l \geq 7} \varepsilon^{k+l-4} E^{-1/2} \mathcal{B}(F_k, F_l) - \varepsilon^3 E^{-1/2} (\partial_t + v \cdot \nabla_x) F_6 \\ &+ \frac{1}{2} R_\varepsilon (\partial_t + v \cdot \nabla_x) \ln E, \end{aligned} \tag{9.7}$$

where E is the local Maxwellian whose parameters solve (9.2) and the quadratic operator \mathbf{Q}_M is defined by

$$\mathbf{Q}_M(\phi, \phi) = M^{-1/2} \mathcal{B}(\sqrt{M} \phi, \sqrt{M} \phi).$$

The last term on the right-hand side of (9.7) is the most annoying one, since it grows like

$$\frac{1}{2} |v|^3 \frac{|\nabla_x \theta|}{\theta^2} R_\varepsilon;$$

1 in particular, it cannot be controlled by the damping part of the linearized collision integral, 1
 2 since $a_M(|v|) = O(|v|)$ as $|v| \rightarrow +\infty$. 2

3 To overcome this difficulty, R. Caflisch introduced a new Maxwellian state \mathbf{M} defined 3
 4 by 4

$$5 \quad \mathbf{M}(v) = \mathcal{M}_{(1,0,\hat{\theta})}(v) = \frac{1}{(2\pi\hat{\theta})^{3/2}} e^{-|v|^2/2\hat{\theta}}, \quad 5$$

6 where 6

$$7 \quad \hat{\theta} = 2\|\theta\|_{L^\infty}. \quad 7$$

8 Hence, there exists a constant C , that depends on $\|\rho\|_{L^\infty}$, $\|u\|_{L^\infty}$ and $\|1/\theta\|_{L^\infty}$ such that 8
 9

$$10 \quad E \leq CM. \quad 10$$

11 Next, one decomposes R_ε as 11

$$12 \quad R_\varepsilon = \sqrt{E}r_\varepsilon + \sqrt{\mathbf{M}}q_\varepsilon, \quad 12$$

13 where the new unknowns r_ε and q_ε are governed by the coupled system 13
 14

$$15 \quad \begin{aligned} 15 \quad (\partial_t + v \cdot \nabla_x)r_\varepsilon &= -\frac{1}{\varepsilon}\mathbf{L}_E r_\varepsilon + \mathbb{1}_{|v| \leq c} \frac{1}{\varepsilon} \sigma \mathbf{K}_M q_\varepsilon, & 15 \\ 16 \quad (\partial_t + v \cdot \nabla_x)q_\varepsilon &= -\mathbf{M}^{-1/2} r_\varepsilon (\partial_t + v \cdot \nabla_x) E^{1/2} - \frac{1}{\varepsilon} (a_M - \mathbb{1}_{|v| > c} \mathbf{K}_M) q_\varepsilon & 16 \\ &+ 2\mathbf{M}^{-1/2} \mathcal{B} \left(\sum_{k=1}^6 \varepsilon^{k-1} F_k, \mathbf{M}^{1/2} (\sigma r_\varepsilon + q_\varepsilon) \right) & 17 \\ &+ 2\mathbf{Q}_M (\sigma r_\varepsilon + q_\varepsilon, \sigma r_\varepsilon + q_\varepsilon) + \varepsilon^2 s, & 18 \end{aligned} \quad (9.8)$$

19 where 19

$$20 \quad \sigma = \sqrt{\frac{E}{\mathbf{M}}}, \quad 20$$

21 where c is a truncation parameter to be chosen below, and the source term s is 21
 22

$$23 \quad s = \mathbf{M}^{-1/2} \sum_{k+l \geq 7} \varepsilon^{k+l-6} \mathcal{B}(F_k, F_l) - \varepsilon \mathbf{M}^{-1/2} (\partial_t + v \cdot \nabla_x) F_6. \quad 23$$

24 This system is solved for the initial condition 24
 25

$$26 \quad r_\varepsilon|_{t=0} = q_\varepsilon|_{t=0} = 0 \quad 26$$

1 by a fixed point argument in the norm 1

$$2 \quad \|\phi\|_{r,s} = \sup_{\xi \in \mathbb{R}^3} (1 + |\xi|)^r \|f(\cdot, \xi)\|_{H^s(\mathbb{R}^3)} \quad 2$$

3 for $s > 3/2$. We refer to [22] for the complete proof. 3

4 REMARKS. Several remarks are in order. 4

- 5 1. The construction in Caglioli's theorem leads to a solution of the Boltzmann equation 5
- 6 that exists and approximates the Maxwellian built on the solution of the compressible 6
- 7 Euler system for as long as the solution of that system exists and remains smooth. 7
- 8 This is very satisfying: should there be a blow-up in finite time in the solution of the 8
- 9 Boltzmann equation, it cannot happen before the onset of singularities in the Euler 9
- 10 system. 10
- 11 2. There is however a rather unpleasant feature in Caglioli's construction: the solution 11
- 12 of the Boltzmann equation so constructed is, in general, not everywhere nonnegative, 12
- 13 and therefore loses physical meaning. This is most easily seen on the initial data for 13
- 14 F_ε in the form (9.4): in Caglioli's paper, $R_\varepsilon|_{t=0} = 0$, so that, in the particular case 14
- 15 of Maxwell's molecules, $F_\varepsilon|_{t=0}$ is a polynomial in v that is not (at least in general) 15
- 16 everywhere nonnegative. It could be that an improvement of Caglioli's ansatz with 16
- 17 initial layers as in [73] helps avoiding this; however, there is no mention of this diffi- 17
- 18 culty in either [22] or [73]. 18
- 19 3. The interested reader is invited to compare Caglioli's theorem with an earlier re- 19
- 20 sult by Nishida [103], who obtained the compressible Euler limit of the Boltzmann 20
- 21 equation by an abstract Cauchy–Kowalewski argument (in the style of Nirenberg and 21
- 22 Ovsyannikov, see, for instance, [102]). Nishida's result is as follows: consider the 22
- 23 scaled Boltzmann equation (9.1) with an initial data F^{in} analytic in x with enough 23
- 24 decay in v that is a perturbation of some absolute Maxwellian. Then, there exists 24
- 25 a family of solutions of the scaled Boltzmann parametrized by $\varepsilon > 0$ that lives on 25
- 26 some interval of time independent of $\varepsilon > 0$. This family of solutions converges to 26
- 27 the Maxwellian built on an analytic solution of the compressible Euler system in the 27
- 28 vanishing ε limit. However, the lifespan of Nishida's family of solutions of the scaled 28
- 29 Boltzmann equation is not known to coincide with the blow-up time of the limiting 29
- 30 smooth solution of the compressible Euler system. See also [118] for an improved 30
- 31 variant of Nishida's result. 31
- 32 32 32
- 33 33 33
- 34 34 34
- 35 35 35
- 36 36 36

37 10. The relative entropy method: Application to the incompressible Euler limit 37

38 As explained in Section 9, all methods based on asymptotic expansions in the Knudsen 38

39 number apply only to situations where the solutions of both the Boltzmann equation and the 39

40 limiting hydrodynamic equation are smooth. The present section introduces a new method 40

41 for cases where only the solution of the target equation (i.e., the hydrodynamic model) is 41

42 smooth. 42

43 43

44 44

45 45

1 We shall explain how this method applies to the incompressible Euler limit of the Boltzmann equation. Consider the Boltzmann equation in the incompressible Euler scaling

$$\begin{aligned} \varepsilon \partial_t F_\varepsilon + v \cdot \nabla_x F_\varepsilon &= \frac{1}{\varepsilon^q} \mathcal{B}(F_\varepsilon, F_\varepsilon), \quad (t, x, v) \in \mathbb{R}_+^* \times \mathbb{T}^3 \times \mathbb{R}^3, \\ F_\varepsilon|_{t=0} &= \mathcal{M}(1, \varepsilon u^{\text{in}}, 1), \end{aligned} \tag{10.1}$$

2 where $q > 1$ and $u^{\text{in}} \equiv u^{\text{in}}(x)$ is a divergence-free vector field on \mathbb{T}^3 . Here, the collision kernel b is supposed to satisfy assumption (3.55) (i.e., to come from a hard cut-off potential) as well as the additional condition

$$\inf_{(z, \omega) \in \mathbb{R}^3 \times \mathbb{S}^2} \frac{b(z, \omega)}{|(z/|z|) \cdot \omega|} > 0. \tag{10.2}$$

3 Throughout this section, we denote by M the centered reduced Gaussian distribution in v , i.e.,

$$M(v) = \mathcal{M}_{(1,0,1)}(v) = \frac{1}{(2\pi)^{3/2}} e^{-|v|^2/2}.$$

4 Next, we introduce a new concept of limit that is especially well adapted to all incompressible hydrodynamic limits of the Boltzmann equation.

5 **DEFINITION 10.1** (Bardos, Golse and Levermore [10]). A family $g_\varepsilon \equiv g_\varepsilon(x, v)$ of $L^1_{\text{loc}}(\mathbb{T}^3 \times \mathbb{R}^3; M \, dx \, dv)$ is said to converge entropically to g at rate ε if

- $1 + \varepsilon g_\varepsilon \geq 0$ a.e. on $\mathbb{T}^3 \times \mathbb{R}^3$ for each $\varepsilon > 0$,
- $g_\varepsilon \rightarrow g$ weakly in $L^1_{\text{loc}}(\mathbb{T}^3 \times \mathbb{R}^3; M \, dx \, dv)$,
- and

$$\frac{1}{\varepsilon^2} H(M(1 + \varepsilon g_\varepsilon) | M) \rightarrow \frac{1}{2} \iint_{\mathbb{T}^3 \times \mathbb{R}^3} g(x, v)^2 \, dx \, dv$$

as $\varepsilon \rightarrow 0$.

6 After these preliminaries, we can state the incompressible Euler limit theorem.

7 **THEOREM 10.2** (Saint-Raymond [110]). Assume that $u^{\text{in}} \in H^3(\mathbb{T}^3)$ is a divergence-free vector field, and let u be the maximal solution of the incompressible Euler equations

$$\begin{aligned} \partial_t u + u \cdot \nabla_x u + \nabla_x p &= 0, \quad \text{div}_x u = 0, \\ u|_{t=0} &= u^{\text{in}} \end{aligned}$$

8 on $[0, T) \times \mathbb{T}^3$. For $\varepsilon > 0$, let F_ε be a renormalized solution relative to M of the scaled Boltzmann equation (10.1). Then, for each $t \in [0, T)$,

$$\frac{F_\varepsilon(t, x, v) - M(v)}{\varepsilon M(v)} \rightarrow u(t, x) \cdot v$$

1 *entropically at rate ε as $\varepsilon \rightarrow 0$.* 1

2
3 The relative entropy method was used for the first time to derive incompressible hy- 3
4 drodynamic models from the Boltzmann equation in Chapter 2 of [18], [54] and in [88] 4
5 – however, the results obtained in these references were incomplete since the proofs used 5
6 additional controls not known to be satisfied by renormalized solutions of the Boltzmann 6
7 equation. 7

8 Of course, if u is a 2D–3C solution of the incompressible Euler equations (see Sec- 8
9 tion 7.5), then $T = +\infty$ and the incompressible Euler limit is global. 9

10 Let us now describe the main ideas in the relative entropy method.

11 First, assuming that F_ε is a classical solution to the scaled Boltzmann equation (10.1), 11
12 we compute 12

$$13 \frac{d}{dt} \frac{1}{\varepsilon^2} H(F_\varepsilon | \mathcal{M}_{(1, \varepsilon u, 1)}) = -\frac{1}{\varepsilon^2} \int_{\mathbb{T}^3} D(u) : \int_{\mathbb{R}^3} (v - \varepsilon u)^{\otimes 2} F_\varepsilon \, dv \, dx \quad 14$$

$$15 + \frac{1}{\varepsilon} \int_{\mathbb{T}^3} \nabla_x p \cdot \int_{\mathbb{R}^3} (v - \varepsilon u) F_\varepsilon \, dv \, dx. \quad 16$$

17
18
19 Now, this identity is not known to be true if F_ε is a renormalized solution. What is known 19
20 instead is the following variant of it (see Theorem 8.10); for each $t \in [0, T)$, one has 20

$$21 \frac{1}{\varepsilon^2} H(F_\varepsilon | \mathcal{M}_{(1, \varepsilon u, 1)})(t) + \frac{1}{\varepsilon} \int_{\mathbb{T}^3} \text{trace}(m_\varepsilon(t)) \quad 22$$

$$23 \leq -\frac{1}{\varepsilon^2} \int_0^t \int_{\mathbb{T}^3} D(u) : \int_{\mathbb{R}^3} (v - \varepsilon u)^{\otimes 2} F_\varepsilon \, dv \, dx \, ds \quad 24 \quad (10.3)$$

$$25 + \frac{1}{\varepsilon} \int_0^t \int_{\mathbb{T}^3} \nabla_x p \cdot \int_{\mathbb{R}^3} (v - \varepsilon u) F_\varepsilon \, dv \, dx \, ds - \frac{1}{\varepsilon} \int_0^t \int_{\mathbb{T}^3} D(u) : m_\varepsilon(s) \, ds. \quad 26$$

27
28
29 The key argument is the following lemma. 30

31
32 LEMMA 10.3. *For each $T' \in [0, T)$ there exists $C_{T'} \geq 1$ such that* 32

$$33 \frac{1}{\varepsilon^2} \int_0^t \int_{\mathbb{T}^3} \int_{\mathbb{R}^3} |D(u) : (v - \varepsilon u)^{\otimes 2} F_\varepsilon| \, dv \, dx \, ds \quad 34$$

$$35 \leq \frac{C_{T'}}{\varepsilon^2} \int_0^t \|D(u)\|_{L^\infty} H(F_\varepsilon | \mathcal{M}_{(1, \varepsilon u, 1)})(s) \, ds + o(1) \quad 36$$

37
38
39 *uniformly in $t \in [0, T']$ as $\varepsilon \rightarrow 0$.* 40

41
42 Define 42

$$43 X_\varepsilon(t) = \frac{1}{\varepsilon^2} H(F_\varepsilon | \mathcal{M}_{(1, \varepsilon u, 1)})(t) + \frac{1}{\varepsilon} \int_{\mathbb{T}^3} \text{trace}(m_\varepsilon(t)); \quad 44$$

1 it follows from (10.3) and Lemma 10.3 that, for each $T' \in [0, T)$ and each $t \in [0, T']$, one
2 has

$$3 \quad X_\varepsilon(t) \leq C_{T'} \|D(u)\|_{L^\infty([0, T'] \times \mathbb{T}^3)} \int_0^t X_\varepsilon(s) \, ds$$

$$4 \quad + \frac{1}{\varepsilon} \int_0^t \int_{\mathbb{T}^3} \nabla_x p \cdot \int_{\mathbb{R}^3} (v - \varepsilon u) F_\varepsilon \, dv \, dx \, ds + o(1)_{L^\infty([0, T'])}.$$

5 Hence

$$6 \quad X_\varepsilon(t) \leq e^{C_{T'} \|D(u)\|_{L^\infty([0, T'] \times \mathbb{T}^3)}} \left| \frac{1}{\varepsilon} \int_0^t \int_{\mathbb{T}^3} \nabla_x p \cdot \int_{\mathbb{R}^3} (v - \varepsilon u) F_\varepsilon \, dv \, dx \, ds \right|$$

$$7 \quad + o(1)_{L^\infty([0, T'])}. \quad (10.4)$$

8 With this inequality, one concludes as follows. The sequence of fluctuations

$$9 \quad \frac{1}{\varepsilon} (F_\varepsilon - M) \text{ is relatively compact in } w^* - L^\infty(\mathbb{R}_+; w - L^1((1 + |v|^2) \, dx \, dv))$$

10 so that, modulo extraction of a subsequence

$$11 \quad \int_{\mathbb{R}^3} F_\varepsilon \, dv \rightarrow 1 \quad \text{in } L^\infty(\mathbb{R}_+; L^1(\mathbb{T}^3))$$

12 and

$$13 \quad \frac{1}{\varepsilon} \int_{\mathbb{R}^3} v F_\varepsilon \, dv \rightarrow U \quad \text{in } w^* - L^\infty(\mathbb{R}_+; w - L^1(\mathbb{T}^3)).$$

14 Because of the local conservation of mass that is satisfied by the renormalized solution F_ε ,
15 one has

$$16 \quad \operatorname{div}_x U = 0.$$

17 Hence

$$18 \quad \frac{1}{\varepsilon} \int_0^t \int_{\mathbb{T}^3} \nabla_x p \cdot \int_{\mathbb{R}^3} (v - \varepsilon u) F_\varepsilon \, dv \, dx \, ds \rightarrow \int_0^t \int_{\mathbb{T}^3} \nabla_x p \cdot (U - u) \, dx \, ds = 0$$

19 so that

$$20 \quad X_\varepsilon(t) \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0 \text{ for each } t \in [0, T'].$$

21 By convexity and weak limit, one has

$$22 \quad \frac{1}{2} \|(U - u)(t)\|_{L^2(\mathbb{T}^3)}^2 \leq \liminf_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2} H(F_\varepsilon | \mathcal{M}_{(1, \varepsilon u, 1)})(t)$$

1 for each $t \in [0, T']$. Therefore, passing to the limit in (10.4) shows that $U = u$, as an-
2 nounced.

3 We shall not describe the proof of Lemma 10.3, which is really technical. In this proof,
4 one has to master the difficulties created by the high velocity tails of the number density;
5 this is done by using decay estimates due to Grad and Caflisch [23] on the gain term of the
6 linearized collision operator, we refer to [110] for a complete proof.

7 But the difficulties in the proof of Lemma 10.3 are special to the Boltzmann equation;
8 the main line of the relative entropy method is as described above. This method is due to
9 Yau [122] who used it for the hydrodynamic limit of Ginzburg–Landau models.

11. Applications of the moment method

14 The moment method is based on compactness results which are used to pass to the limit
15 in the local conservation laws of mass momentum and energy as the Knudsen number
16 vanishes. This method does not require estimates other than the natural bounds on mass,
17 energy, entropy and entropy production. On principle, it could therefore be used when both
18 the solutions of the scaled Boltzmann equation and of the limiting hydrodynamic equations
19 are not known to be regular.

20 First, we state the various theorems on hydrodynamic limits that can be proved in this
21 way.

11.1. The acoustic limit

25 We start from the Boltzmann equation in the acoustic scaling posed in the periodic box

$$27 \quad \partial_t F_\varepsilon + v \cdot \nabla_x F_\varepsilon = \frac{1}{\varepsilon} \mathcal{B}(F_\varepsilon, F_\varepsilon), \quad (t, x, v) \in \mathbb{R}_+^* \times \mathbb{T}^3 \times \mathbb{R}^3, \quad 28$$

$$29 \quad F_\varepsilon|_{t=0} = F_\varepsilon^{\text{in}}. \quad 30 \quad (11.1)$$

31 Assume that b satisfies the weak cut-off assumption (8.1) as well as the bound

$$33 \quad 0 < \int_{\mathbb{S}^2} b(z, \omega) d\omega \leq C_b (1 + |z|^2)^\beta \quad \text{a.e., } z \in \mathbb{R}^3, \quad 34 \quad (11.2)$$

35 for some $\beta \in [0, 1]$.

36 We assume that

$$37 \quad \iint_{\mathbb{T}^3 \times \mathbb{R}^3} F_\varepsilon^{\text{in}} dx dv = 1, \quad 38$$

$$39 \quad \iint_{\mathbb{T}^3 \times \mathbb{R}^3} v F_\varepsilon^{\text{in}} dx dv = 0, \quad 40 \quad (11.3)$$

$$41 \quad \iint_{\mathbb{T}^3 \times \mathbb{R}^3} \frac{1}{2} |v|^2 F_\varepsilon^{\text{in}} dx dv = \frac{3}{2}. \quad 42$$

1 Set M to be the centered, reduced Gaussian distribution

$$2 \quad 3 \quad 4 \quad 5 \quad M(v) = \frac{1}{(2\pi)^{3/2}} e^{-|v|^2/2}, \quad (11.4)$$

6 so that

$$7 \quad 8 \quad 9 \quad \int_{\mathbb{R}^3} M \, dx \, dv = 1,$$

$$10 \quad 11 \quad 12 \quad \int_{\mathbb{R}^3} v M \, dx \, dv = 0,$$

$$13 \quad 14 \quad 15 \quad \int_{\mathbb{R}^3} \frac{1}{2} |v|^2 M \, dx \, dv = \frac{3}{2}.$$

16 The acoustic limit of the Boltzmann equation (11.1) is given by the following theorem.

17
18 **THEOREM 11.1** (Golse and Levermore [53]). *Let $\delta_\varepsilon > 0$ be such that*

$$19 \quad 20 \quad 21 \quad \delta_\varepsilon \rightarrow 0 \quad \text{and} \quad \delta_\varepsilon |\ln \delta_\varepsilon|^{\beta/2} = o(\varepsilon^{1/2})$$

22 as $\varepsilon \rightarrow 0$. Assume that

$$23 \quad 24 \quad 25 \quad \frac{F_\varepsilon^{\text{in}}(x, v) - M}{\delta_\varepsilon M} \rightarrow \rho^{\text{in}}(x) + u^{\text{in}}(x) \cdot v + \theta^{\text{in}}(x) \frac{1}{2} (|v|^2 - 3)$$

26
27 *entropically at rate δ_ε . For each $\varepsilon > 0$, let F_ε be a renormalized solution relative to M of*

28 *the scaled Boltzmann equation (11.1) with initial data $F_\varepsilon^{\text{in}}$.*

29 *Then, for each $t \geq 0$, the family*

$$30 \quad 31 \quad 32 \quad \frac{F_\varepsilon(t, x, v) - M}{\delta_\varepsilon M} \rightarrow \rho(t, x) + u(t, x) \cdot v + \theta(t, x) \frac{1}{2} (|v|^2 - 3)$$

33
34 *entropically at rate δ_ε , where (ρ, u, θ) is a solution of the acoustic system*

$$35 \quad 36 \quad 37 \quad \partial_t \rho + \text{div}_x u = 0,$$

$$38 \quad 39 \quad \partial_t u + \nabla_x (\rho + \theta) = 0, \quad (t, x) \in \mathbb{R}_+^* \times \mathbb{T}^3,$$

$$40 \quad 41 \quad \frac{3}{2} \partial_t \theta + \text{div}_x u = 0,$$

42 with initial data

$$43 \quad 44 \quad 45 \quad (\rho, u, \theta)|_{t=0} = (\rho^{\text{in}}, u^{\text{in}}, \theta^{\text{in}}).$$

1 Notice that our assumptions on the family of initial data imply that 1

$$2 (\rho^{\text{in}}, u^{\text{in}}, \theta^{\text{in}}) \in L^2(\mathbb{T}^3), 2$$

3 and that 3

$$4 \int_{\mathbb{T}^3} \rho^{\text{in}}(x) dx = 0, \quad \int_{\mathbb{T}^3} u^{\text{in}}(x) dx = 0 \quad \text{and} \quad \int_{\mathbb{T}^3} \theta^{\text{in}}(x) dx = 0. 4$$

5 In particular, the existence and uniqueness theory for the acoustic system described in 5
6 Section 7.1.2 applies here, and the solution (ρ, u, θ) satisfies 6

$$7 (\rho, u, \theta) \in C(\mathbb{R}_+; L^2(\mathbb{T}^3)) 7$$

8 and 8

$$9 \int_{\mathbb{T}^3} \rho^{\text{in}}(x) dx = 0, \quad \int_{\mathbb{T}^3} u^{\text{in}}(x) dx = 0 \quad \text{and} \quad \int_{\mathbb{T}^3} \theta^{\text{in}}(x) dx = 0. 9$$

10 11.2. The Stokes–Fourier limit 10

11 We start from the Boltzmann equation in the Stokes scaling posed in the periodic box 11

$$12 \begin{aligned} 12 \quad \varepsilon \partial_t F_\varepsilon + v \cdot \nabla_x F_\varepsilon &= \frac{1}{\varepsilon} \mathcal{B}(F_\varepsilon, F_\varepsilon), \quad (t, x, v) \in \mathbb{R}_+^* \times \mathbb{T}^3 \times \mathbb{R}^3, & 12 \\ 13 \quad F_\varepsilon|_{t=0} &= F_\varepsilon^{\text{in}}. & 13 \end{aligned} \quad (11.5)$$

14 Assume that the collision kernel b comes from a hard cut-off potential, i.e., that it sat- 14
15 isfies (5.2) for some $\alpha \in [0, 1]$. Assume further that, for each $\varepsilon > 0$, the initial data $F_\varepsilon^{\text{in}}$ 15
16 satisfies the relations (11.3). 16

17 **THEOREM 11.2** (Golse and Levermore [53]). *Let $\delta_\varepsilon > 0$ be such that* 17

$$18 \delta_\varepsilon \rightarrow 0 \quad \text{and} \quad \delta_\varepsilon |\ln \delta_\varepsilon|^\alpha = o(\varepsilon) 18$$

19 as $\varepsilon \rightarrow 0$. Assume that 19

$$20 \frac{F_\varepsilon^{\text{in}}(x, v) - M}{\delta_\varepsilon M} \rightarrow u^{\text{in}}(x) \cdot v + \theta^{\text{in}}(x) \frac{1}{2} (|v|^2 - 5) 20$$

21 entropically at rate δ_ε , where u^{in} satisfies 21

$$22 \operatorname{div}_x u^{\text{in}} = 0 22$$

1 and where M is the centered, reduced Gaussian distribution (11.4). For each $\varepsilon > 0$, let
 2 F_ε be a renormalized solution relative to M of the scaled Boltzmann equation (11.5) with
 3 initial data $F_\varepsilon^{\text{in}}$.

4 Then, for each $t \geq 0$, the family

$$5 \quad \frac{F_\varepsilon(t, x, v) - M}{\delta_\varepsilon M} \rightarrow u(t, x) \cdot v + \theta(t, x) \frac{1}{2} (|v|^2 - 5)$$

6 entropically at rate δ_ε , where (u, θ) is a solution of the Stokes–Fourier system

$$7 \quad \begin{aligned} \partial_t u + \nabla_x p &= v \Delta_x u, & \operatorname{div}_x u &= 0, & (t, x) &\in \mathbb{R}_+^* \times \mathbb{T}^3, \\ \partial_t \theta &= \kappa \Delta_x \theta, \end{aligned}$$

8 with initial data

$$9 \quad (u, \theta)|_{t=0} = (u^{\text{in}}, \theta^{\text{in}}).$$

10 The viscosity and heat conductivity are given by

$$11 \quad \nu = \frac{1}{10} \int_{\mathbb{R}^3} \widehat{A} : AM \, dv, \quad \kappa = \frac{2}{15} \int_{\mathbb{R}^3} \widehat{B} \cdot BM \, dv,$$

12 see also formula (6.9) for expressions of these quantities in terms of the functions \mathbf{a} and \mathbf{b}
 13 defined in (6.12).

14 We recall that \widehat{A} and \widehat{B} are defined in terms of

$$15 \quad A(v) = v \otimes v - \frac{1}{3} |v|^2 I, \quad B = \frac{1}{2} (|v|^2 - 5) v$$

16 by

$$17 \quad \begin{aligned} \mathcal{L}_M \widehat{A} &= A, & \widehat{A} &\perp \operatorname{Ker} \mathcal{L}_M, \\ \mathcal{L}_M \widehat{B} &= B, & \widehat{B} &\perp \operatorname{Ker} \mathcal{L}_M. \end{aligned}$$

18 Our assumptions on the family of initial data imply that

$$19 \quad (u^{\text{in}}, \theta^{\text{in}}) \in L^2(\mathbb{T}^3),$$

20 while

$$21 \quad \int_{\mathbb{T}^3} u^{\text{in}}(x) \, dx = 0 \quad \text{and} \quad \int_{\mathbb{T}^3} \theta^{\text{in}}(x) \, dx = 0.$$

Hence, the existence and uniqueness theory for the Stokes–Fourier system described in Section 7.1.1 applies here, and the solution (u, θ) satisfies

$$(u, \theta) \in C(\mathbb{R}_+; L^2(\mathbb{T}^3)) \cap C^\infty(\mathbb{R}_+^* \times \mathbb{R}^3)$$

and

$$\int_{\mathbb{T}^3} u^{\text{in}}(x) \, dx = 0 \quad \text{and} \quad \int_{\mathbb{T}^3} \theta^{\text{in}}(x) \, dx = 0.$$

11.3. The Navier–Stokes–Fourier limit

We start from the Boltzmann equation in the Navier–Stokes scaling, posed in the Euclidean space, with Maxwellian equilibrium at infinity

$$\begin{aligned} \varepsilon \partial_t F_\varepsilon + v \cdot \nabla_x F_\varepsilon &= \frac{1}{\varepsilon} \mathcal{B}(F_\varepsilon, F_\varepsilon), \quad (t, x, v) \in \mathbb{R}_+^* \times \mathbb{R}^3 \times \mathbb{R}^3, \\ F_\varepsilon(t, x, v) &\rightarrow M \quad |x| \rightarrow +\infty, \\ F_\varepsilon|_{t=0} &= F_\varepsilon^{\text{in}}, \end{aligned} \tag{11.6}$$

where M is the centered, reduced Gaussian distribution (11.4).

Assume that the collision kernel b comes from a hard cut-off potential, i.e., that it satisfies (5.2) for some $\alpha \in [0, 1]$.

THEOREM 11.3 (Golse and Saint-Raymond [61,62]). *Assume that*

$$\frac{F_\varepsilon^{\text{in}}(x, v) - M}{\varepsilon M} \rightarrow u^{\text{in}}(x) \cdot v + \theta^{\text{in}}(x) \frac{1}{2}(|v|^2 - 5)$$

entropically at rate ε , where u^{in} satisfies

$$\operatorname{div}_x u^{\text{in}} = 0.$$

For each $\varepsilon > 0$, let F_ε be a renormalized solution relative to M of the scaled Boltzmann equation (11.5) with initial data $F_\varepsilon^{\text{in}}$.

Then the family

$$\left(\frac{1}{\varepsilon} \int_{\mathbb{R}^3} v F_\varepsilon(t, x, v) \, dv, \frac{1}{\varepsilon} \int_{\mathbb{R}^3} \left(\frac{1}{3} |v|^2 - 1 \right) (F_\varepsilon(t, x, v) - M) \, dv \right)$$

is weakly relatively compact in $L^1_{\text{loc}}(\mathbb{R}_+ \times \mathbb{R}^3)$, and each of its limit points as $\varepsilon \rightarrow 0$ is

$$(u, \theta) \in C(\mathbb{R}_+, w - L^2(\mathbb{R}^3)) \cap L^2(\mathbb{R}_+; H^1(\mathbb{R}^3)),$$

1 *a solution of the Navier–Stokes–Fourier system* 1

2

3

$$\partial_t u + \operatorname{div}_x(u \otimes u) + \nabla_x p = \nu \Delta_x u, \quad \operatorname{div}_x u = 0, \quad (t, x) \in \mathbb{R}_+^* \times \mathbb{T}^3,$$

4

5

$$\partial_t \theta + \operatorname{div}_x(u \theta) = \kappa \Delta_x \theta,$$

6

7

8

9

$$(u, \theta)|_{t=0} = (u^{\text{in}}, \theta^{\text{in}}).$$

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

The viscosity and heat conductivity are given by

$$\nu = \frac{1}{10} \int_{\mathbb{R}^3} \widehat{A} : AM \, dv, \quad \kappa = \frac{2}{15} \int_{\mathbb{R}^3} \widehat{B} \cdot BM \, dv$$

again, see (6.9) for expressions of these quantities in terms of the functions **a** and **b** defined in (6.12). Moreover, this solution (u, θ) satisfies, for each $t > 0$, the inequality

$$\begin{aligned} & \frac{1}{2} \|u(t)\|_{L^2}^2 + \frac{5}{4} \|\theta(t)\|_{L^2}^2 + \int_0^t \int_{\mathbb{R}^3} (\nu |\nabla_x u|^2 + \kappa |\nabla_x \theta|^2) \, dx \, ds \\ & \leq \frac{1}{2} \|u^{\text{in}}\|_{L^2}^2 + \frac{5}{4} \|\theta^{\text{in}}\|_{L^2}^2. \end{aligned}$$

In particular, if $\theta^{\text{in}} = 0$, this theorem shows that any weak limit point of

$$\frac{1}{\varepsilon} \int_{\mathbb{R}^3} v F_\varepsilon(t, x, v) \, dv$$

in $L_{\text{loc}}^1(\mathbb{R}_+ \times \mathbb{R}^3)$ as $\varepsilon \rightarrow 0$ is a Leray solution of the Navier–Stokes equations with initial data u^{in} .

This theorem explains the following observation by Lions: “[...] the global existence of [renormalized] solutions [...] can be seen as the analogue for Boltzmann’s equation to the pioneering work on the Navier–Stokes equations by J. Leray” (see [84], p. 432).

11.4. Sketch of the proof of the Navier–Stokes–Fourier limit by the moment method

The proof of the Navier–Stokes–Fourier limit theorem above involves many ideas developed in a sequence of papers over the past 15 years:

- the BGL program was defined in [10]; this reference provided the general entropy and entropy production estimates used to control the number density fluctuation and its distance to local equilibrium; as a result, the evolution Stokes and the steady Navier–Stokes motion equations were derived under the assumption that the renormalized solutions to the Boltzmann equation considered satisfy the local conservation of momentum as well as a nonlinear compactness estimate (for the Navier–Stokes limit) that will be described below in more details;

- 1 • under the same assumptions as in [10], Lions and Masmoudi [87] were able to de- 1
 2 rive the evolution Navier–Stokes motion equations, by a kind of “compensated com- 2
 3 pactness” argument bearing on fast oscillating acoustic waves; they also introduced 3
 4 a slightly modified notion of renormalized solution which led them to a complete 4
 5 derivation of the evolution Stokes motion equation; 5
- 6 • in [11], it was observed for the first time that the local conservation law of momen- 6
 7 tum could be *proved* in the hydrodynamic limit, thereby relieving the need for as- 7
 8 suming that local conservation law at the level of the renormalized solutions of the 8
 9 Boltzmann equation; this led to a complete proof of the acoustic limit for bounded 9
 10 collision kernels; a more complete understanding of how the local conservation laws 10
 11 of both momentum and energy could be proved in the hydrodynamic limit was even- 11
 12 tually reached in [53]; the latter reference provided an essentially optimal derivation 12
 13 of the Stokes motion and energy equations as well as a derivation of the acoustic sys- 13
 14 tem that allowed for the most general hard cut-off potentials; however, the acoustic 14
 15 system was established only under some unphysical restriction on the scaling of the 15
 16 number density fluctuation; 16
- 17 • in [107,109], Saint-Raymond gave a complete derivation of both the Navier–Stokes 17
 18 motion and energy equations for the BGK model with constant relaxation time; her 18
 19 proof was based on obtaining for the first time some weaker analogue of the nonlinear 19
 20 compactness assumption used in [10]; 20
- 21 • finally, a complete derivation of the incompressible Navier–Stokes motion and heat 21
 22 equations from the Boltzmann equation was proposed for the first time in [61] for 22
 23 bounded collision kernels (such as occurring in the case of cut-off Maxwell mole- 23
 24 cules); this reference used all the methods constructed in the previous works men- 24
 25 tioned above, together with a new velocity averaging method specific to the L^1 case 25
 26 and that amplified Saint-Raymond’s observation in [107]; this result was later ex- 26
 27 tended to all hard cut-off potentials (including hard spheres) in [62]. 27

28 It is this last reference that we describe below; although its scope is more general than 28
 29 that of [61], it involves a new idea for handling unbounded collision kernels that actually 29
 30 simplifies the discussion in [61]. 30
 31

32 **11.4.1. A priori estimates.** In this subsection, we quickly list the a priori estimates on 32
 33 the family F_ε of solutions of the Boltzmann equation that are uniform in $\varepsilon > 0$. As can be 33
 34 seen from Theorem 8.10, the only such estimate comes from the DiPerna–Lions variant of 34
 35 Boltzmann’s H -theorem, 35
 36

$$\begin{aligned}
 & H(F_\varepsilon(t)|M) + \frac{1}{4\varepsilon^2} \int_0^t \int_{\mathbb{R}^3} \int \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F'_\varepsilon F'_{\varepsilon*} - F_\varepsilon F_{\varepsilon*}) \\
 & \quad \times \ln\left(\frac{F'_\varepsilon F'_{\varepsilon*}}{F_\varepsilon F_{\varepsilon*}}\right) b \, dv \, dv_* \, d\omega \, dx \, ds \\
 & \leq H(F_\varepsilon^{\text{in}}|M). \tag{11.7}
 \end{aligned}$$

44 We shall further transform (11.7) with the two following inequalities. 44
 45

1 *Pointwise inequalities.* For each ξ and $\eta \in \mathbb{R}_+^*$,

$$2 \quad 3 \quad 4 \quad 5 \quad 6 \quad 7 \quad 8 \quad 9 \quad 10 \quad 11 \quad 12 \quad 13 \quad 14 \quad 15 \quad 16 \quad 17 \quad 18 \quad 19 \quad 20 \quad 21 \quad 22 \quad 23 \quad 24 \quad 25 \quad 26 \quad 27 \quad 28 \quad 29 \quad 30 \quad 31 \quad 32 \quad 33 \quad 34 \quad 35 \quad 36 \quad 37 \quad 38 \quad 39 \quad 40 \quad 41 \quad 42 \quad 43 \quad 44 \quad 45$$

$$(\sqrt{\xi} - 1)^2 \leq \frac{1}{4}(\xi \ln \xi - \xi + 1) \quad (11.8)$$

while

$$(\sqrt{\xi} - \sqrt{\eta})^2 \leq \frac{1}{4}(\xi - \eta)(\ln \xi - \ln \eta). \quad (11.9)$$

Since $(F_\varepsilon - M)/(\varepsilon M)$ converges entropically at rate ε , one has

$$H(F_\varepsilon^{\text{in}} | M) \leq C^{\text{in}} \varepsilon^2. \quad (11.10)$$

This bound and the relative entropy inequality (11.7) entail the following entropy bound for each $t > 0$

$$H(F_\varepsilon(t) | M) \leq C^{\text{in}} \varepsilon^2 \quad (11.11)$$

and the following entropy production bounds

$$\int_0^{+\infty} \int_{\mathbb{R}^3} \int \int \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} (F'_\varepsilon F'_{\varepsilon*} - F_\varepsilon F_{\varepsilon*}) \ln \left(\frac{F'_\varepsilon F'_{\varepsilon*}}{F_\varepsilon F_{\varepsilon*}} \right) b \, dv \, dv_* \, d\omega \, dx \, dt \leq 4C^{\text{in}} \varepsilon^4. \quad (11.12)$$

Introducing the relative number density and relative number density fluctuations,

$$G_\varepsilon = \frac{F_\varepsilon}{M} \quad \text{and} \quad g_\varepsilon = \frac{F_\varepsilon - M}{\varepsilon M}, \quad (11.13)$$

the two pointwise inequalities (11.8) and (11.9) convert the entropy and entropy production bounds into the two uniform a priori estimates

$$\int_{\mathbb{R}^3} \langle (\sqrt{G_\varepsilon(t)} - 1)^2 \rangle dx \leq \frac{1}{4} C^{\text{in}} \varepsilon^2 \quad (11.14)$$

and

$$\int_0^{+\infty} \int_{\mathbb{R}^3} \langle \langle (\sqrt{G'_\varepsilon G'_{\varepsilon*}} - \sqrt{G_\varepsilon G_{\varepsilon*}})^2 \rangle \rangle dx \, dt \leq C^{\text{in}} \varepsilon^4. \quad (11.15)$$

The importance of the two a priori bounds above (11.14) and (11.15) in the derivation of the Navier–Stokes limit cannot be overestimated. In fact, various analogues of these bounds were used earlier in the context of nonlinear diffusion limits, see [12] and especially [59].

The importance of the entropy and entropy dissipation bounds for hydrodynamic limits of diffusive type was noticed. We recall that

$$\langle \phi \rangle = \int_{\mathbb{R}^3} \phi(v) M(v) dv,$$

and we further introduce the notation

$$\langle\langle \Phi \rangle\rangle = \iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} \Phi(v, v_*, \omega) d\mu(v, v_*, \omega),$$

where

$$d\mu(v, v_*, \omega) = b(v - v_*, \omega) M(v) dv M(v_*) dv_* d\omega.$$

In the sequel, we outline the main ideas in the derivation of the Navier–Stokes motion equation in Theorem 11.3, since the derivation of the heat equation is essentially analogous.

Of course, this proof more or less follows the formal argument presented in Section 6.1. However, several key properties of the solutions to the scaled Boltzmann equation used in this formal argument – such as, for instance, the local conservation laws of momentum and energy – are not known to be satisfied by renormalized solutions. Hence the proof sketched below differs noticeably from the formal argument in several places, yet the general idea remains essentially the same.

11.4.2. Normalizing functions. As explained in Section 8.2, the Boltzmann equation can be equivalently renormalized with any admissible nonlinearity whose derivative saturates the quadratic growth of the collision integral.

Throughout the proof of the Navier–Stokes limit theorem, we shall essentially use two kinds of normalizing nonlinearities

- compactly supported nonlinearities that coincide with the identity near the reference Maxwellian state, and
- variants of the maximal, i.e., square-root renormalization.

Nonlinearities of the first kind are used to define the renormalized form of the Boltzmann equation in which one passes to the vanishing ε limit, while the square-root normalization is used to establish compactness properties of the family of solutions to the scaled Boltzmann equation.

The first kind of normalizing nonlinearities is defined through the class of bump functions $\gamma \in C^\infty(\mathbb{R}_+)$ such that

$$\gamma|_{[0,3/2]} \equiv 1, \quad \gamma|_{[2,+\infty)} \equiv 0, \quad \gamma \text{ is nonincreasing on } \mathbb{R}_+. \quad (11.16)$$

The Boltzmann equation is then renormalized with the nonlinearity

$$\Gamma(Z) = (Z - 1)\gamma(Z);$$

1 later on, we denote

$$2 \hat{\gamma}(Z) = \frac{d}{dZ}((Z-1)\gamma(Z)) = \Gamma'(Z). \quad (11.17)$$

3 The scaled Boltzmann equation renormalized with Γ is put in the form

$$4 \partial_t(g_\varepsilon \gamma_\varepsilon) + \frac{1}{\varepsilon} v \cdot \nabla_x(g_\varepsilon \gamma_\varepsilon) = \frac{1}{\varepsilon^3} \hat{\gamma}_\varepsilon \mathcal{Q}_M(G_\varepsilon, G_\varepsilon), \quad (11.18)$$

5 where we have denoted

$$6 \gamma_\varepsilon = \gamma(G_\varepsilon), \quad \hat{\gamma}_\varepsilon = \hat{\gamma}(G_\varepsilon),$$

7 and where \mathcal{Q} designates the Boltzmann collision integral intertwined with the multiplication by M (see (6.6)),

$$8 \mathcal{Q}(G, G) = M^{-1} \mathcal{B}(MG, MG).$$

9 Later on, we shall pass to the limit in the momentum equation deduced from (11.18).

10 The second class of normalizing nonlinearities that we shall use to establish compactness properties of the number density fluctuations G_ε is defined as

$$11 \Gamma_\zeta(Z) = \sqrt{\zeta + Z}, \quad \zeta > 0,$$

12 where the parameter ζ will later be adapted to ε .

13 **11.4.3. Governing equations for moments of g_ε .** As explained in Theorem 8.10, renormalized solutions to the Boltzmann equation satisfy the local conservation of mass (i.e., the continuity equation); in terms of the number density fluctuation g_ε , this local conservation law is expressed as

$$14 \varepsilon \partial_t \langle g_\varepsilon \rangle + \operatorname{div}_x \langle v g_\varepsilon \rangle = 0. \quad (11.19)$$

15 Now, the entropy bound (11.11) implies that

$$16 (1 + |v|^2)g_\varepsilon \text{ is relatively compact in } w - L^1_{\text{loc}}(dt dx; L^1(M dv)). \quad (11.20)$$

17 Before saying a few words on (11.20), let us explain how we use it. Modulo extraction of a subsequence, one has

$$18 g_\varepsilon \rightarrow g \quad \text{in } w - L^1_{\text{loc}}(dt dx; L^1((1 + |v|^2)M dv))$$

19 and hence

$$20 \langle g_\varepsilon \rangle \rightarrow \langle g \rangle \quad \text{and} \quad \langle v g_\varepsilon \rangle \rightarrow \langle v g \rangle \quad \text{in } w - L^1_{\text{loc}}(dt dx).$$

1 Passing to the limit as $\varepsilon \rightarrow 0$ in (11.19) leads to 1

$$2 \quad \operatorname{div}_x \langle v g \rangle = 0, \quad 2$$

3 so that, denoting 3

$$4 \quad u = \langle v g \rangle \quad 4$$

5 the relation above is the incompressibility condition in the Navier–Stokes equations, i.e., 5

$$6 \quad \operatorname{div}_x u = 0. \quad 6$$

7 Let us go back to (11.20) and explain how it follows from the entropy bound (11.11), 7
8 see [10] for more details on this. Define 8

$$9 \quad h(z) = (1+z) \ln(1+z) - z, \quad z > -1. \quad 9$$

10 In terms of h , the entropy bound (11.11) is expressed as 10

$$11 \quad \frac{1}{\varepsilon^2} \int_{\mathbb{R}^3} \langle h(\varepsilon g_\varepsilon(t)) \rangle dx \leq C^{\text{in}}. \quad 11$$

12 If the entropy bound (11.11) was equivalent to an $L^p(M dv dx)$ bound for some $p > 1$, 12
13 Hölder’s inequality would imply that $(1 + |v|)^2 g_\varepsilon$ is bounded in $L^\infty(dt; L^r(M dv dx))$ for 13
14 some $r > 1$, since $(1 + |v|^2) \in L^q(M dv)$ for each $q \in (1, +\infty)$. However, the entropy 14
15 control (11.11) on g_ε is weaker than an $L^p(M dv dx)$ bound. But (11.20) follows from a 15
16 careful use of Young’s inequality 16

$$17 \quad p|z| \leq \frac{\alpha}{\varepsilon^2} h(\varepsilon z) + \frac{1}{\alpha} h^*(p), \quad p > 0, z > -1, 0 < \varepsilon < \alpha, \quad 17$$

18 where 18

$$19 \quad h^*(p) = e^p - p - 1 \quad 19$$

20 designates the Legendre dual of h . The compactness (11.20) follows from replacing z 20
21 with g_ε and p with $\frac{1}{4}(1 + |v|^2)$ in the inequality above, letting then $\alpha \rightarrow 0$ in the inequality 21
22 so obtained. 22

23 Let us now explain how the motion equation in the Navier–Stokes system is derived 23
24 from the Boltzmann equation. This is of course the main part in the proof, and it involves 24
25 several technicalities. 25

26 In particular, we shall need truncations in the velocity variable at a level that is tied to ε . 26
27 For each function $\xi \equiv \xi(v)$ and each $K > 6$, we define 27

$$28 \quad \xi_{K\varepsilon}(v) = \xi(v) \mathbb{1}_{|v|^2 \leq K|\ln \varepsilon|}. \quad (11.21) \quad 28$$

1 Multiplying each side of the scaled, renormalized Boltzmann equation (11.18) by each
2 component of v_{K_ε} and averaging in v leads to

$$3 \quad \partial_t \langle v g_\varepsilon \gamma_\varepsilon \rangle + \operatorname{div}_x \mathbf{F}_\varepsilon(A) + \nabla_x \cdot \left\langle \frac{1}{\varepsilon} |v|_{K_\varepsilon}^2 g_\varepsilon \gamma_\varepsilon \right\rangle = \mathbf{D}_\varepsilon(v), \quad (11.22)$$

4 where $\mathbf{F}_\varepsilon(A)$ is the truncated, renormalized traceless part of the momentum flux

$$5 \quad \mathbf{F}_\varepsilon(A) = \frac{1}{\varepsilon} \langle A_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle \quad (11.23)$$

6 while $\mathbf{D}_\varepsilon(v)$ is the momentum conservation defect

$$7 \quad \mathbf{D}_\varepsilon(v) = \frac{1}{\varepsilon^3} \langle \langle v_{K_\varepsilon} \hat{\gamma}_\varepsilon (G'_\varepsilon G'_{\varepsilon*} - G_\varepsilon G_{\varepsilon*}) \rangle \rangle. \quad (11.24)$$

8 Notice that truncating large velocities in the number density, or large values thereof (which
9 is what the renormalization procedure does) break the symmetries in the collision integral
10 leading to the local conservation of momentum (see Proposition 3.1): this accounts for
11 the defect $\mathbf{D}_\varepsilon(v)$ on the right-hand side of (11.22). As $\varepsilon \rightarrow 0$, $v_{K_\varepsilon} \rightarrow v$ while $G_\varepsilon \rightarrow 1$ so
12 that $\hat{\gamma}_\varepsilon \rightarrow 1$; hence, the missing symmetries are restored in the integrand defining $\mathbf{D}_\varepsilon(v)$.
13 Hence, one can hope that $\mathbf{D}_\varepsilon(v) \rightarrow 0$ as $\varepsilon \rightarrow 0$.

14 In fact, the strategy for establishing the Navier–Stokes limit theorem consists of the
15 following three steps.

16 *Step 1.* Prove that, modulo extraction of a subsequence

$$17 \quad \langle v g_\varepsilon \gamma_\varepsilon \rangle \rightarrow \langle v g \rangle = u \quad \text{in } w - L^1_{\text{loc}}(dt \, dx),$$

18 while

$$19 \quad P \langle v g_\varepsilon \gamma_\varepsilon \rangle \rightarrow u \quad \text{in } C(\mathbb{R}_+; \mathcal{D}'(\mathbb{R}^3)),$$

20 where P denotes the Leray projection, i.e., the orthogonal projection on divergence-free
21 vector fields in $L^2(\mathbb{R}^3)$.

22 *Step 2.* Likewise, prove that

$$23 \quad \mathbf{D}_\varepsilon(v) \rightarrow 0 \quad \text{in } L^1_{\text{loc}}(dt \, dx).$$

24 *Step 3.* Finally prove that

$$25 \quad P(\operatorname{div}_x \mathbf{F}_\varepsilon(A)) \rightarrow P \operatorname{div}_x(u \otimes u) - \nu \Delta_x u \quad \text{in } \mathcal{D}'(\mathbb{R}_+^* \times \mathbb{R}^3).$$

26 Once these three steps are completed, one applies P to both sides of (11.22), which gives

$$27 \quad \partial_t P \langle v g_\varepsilon \gamma_\varepsilon \rangle + P(\operatorname{div}_x \mathbf{F}_\varepsilon(A)) = P \mathbf{D}_\varepsilon(v). \quad (11.25)$$

1 Taking limits in each term as $\varepsilon \rightarrow 0$ shows that u satisfies the Navier–Stokes motion equation. 2
 2 As for the initial condition, observe that it is guaranteed by the uniform convergence 3
 3 in t , i.e., by the second statement in Step 1. 4

5 **11.4.4. Vanishing of the momentum conservation defect.** We start with Step 2, i.e., we 5
 6 explain how to prove the following proposition. 6
 7

8 PROPOSITION 11.4. *Under the same assumptions as in Theorem 11.3,* 8
 9

$$10 \quad \mathbf{D}_\varepsilon(v) \rightarrow 0 \quad \text{in } L^1_{\text{loc}}(dt \, dx). \quad 10$$

11
 12 First, we start from the elementary formula 12
 13

$$14 \quad \begin{aligned} 14 \quad G'_\varepsilon G'_{\varepsilon*} - G_\varepsilon G_{\varepsilon*} & \\ 15 \quad &= (\sqrt{G'_\varepsilon G'_{\varepsilon*}} - \sqrt{G_\varepsilon G_{\varepsilon*}})(\sqrt{G'_\varepsilon G'_{\varepsilon*}} + \sqrt{G_\varepsilon G_{\varepsilon*}}) \\ 16 \quad &= (\sqrt{G'_\varepsilon G'_{\varepsilon*}} - \sqrt{G_\varepsilon G_{\varepsilon*}})^2 + 2\sqrt{G_\varepsilon G_{\varepsilon*}}(\sqrt{G'_\varepsilon G'_{\varepsilon*}} - \sqrt{G_\varepsilon G_{\varepsilon*}}) \end{aligned} \quad 15$$

17
 18 and split the momentum conservation defect as 18
 19

$$20 \quad \mathbf{D}_\varepsilon(v) = \mathbf{D}_\varepsilon^1(v) + \mathbf{D}_\varepsilon^2(v) \quad 20$$

21
 22 with 22
 23

$$24 \quad \mathbf{D}_\varepsilon^1(v) = \frac{1}{\varepsilon^3} \langle \langle v_{K_\varepsilon} \hat{\gamma}_\varepsilon (\sqrt{G'_\varepsilon G'_{\varepsilon*}} - \sqrt{G_\varepsilon G_{\varepsilon*}})^2 \rangle \rangle \quad 24$$

25
 26 and 26
 27

$$28 \quad \mathbf{D}_\varepsilon^2(v) = \frac{2}{\varepsilon^3} \langle \langle v_{K_\varepsilon} \hat{\gamma}_\varepsilon \sqrt{G_\varepsilon G_{\varepsilon*}} (\sqrt{G'_\varepsilon G'_{\varepsilon*}} - \sqrt{G_\varepsilon G_{\varepsilon*}}) \rangle \rangle. \quad 28$$

29
 30 That $\mathbf{D}_\varepsilon^1(v) \rightarrow 0$ in $L^1_{\text{loc}}(dt \, dx)$ follows from the entropy production estimate (11.15). 30
 31 Setting 31
 32

$$33 \quad \mathcal{E}_\varepsilon = \frac{1}{\varepsilon^2} \sqrt{G_\varepsilon G_{\varepsilon*}} (\sqrt{G'_\varepsilon G'_{\varepsilon*}} - \sqrt{G_\varepsilon G_{\varepsilon*}}) \quad 33$$

34
 35 we further split $\mathbf{D}_\varepsilon^2(v)$ as 35
 36

$$37 \quad \begin{aligned} 37 \quad \mathbf{D}_\varepsilon^2(v) &= -\frac{2}{\varepsilon} \langle \langle v \mathbb{1}_{|v|^2 > K_\varepsilon} \hat{\gamma}_\varepsilon \mathcal{E}_\varepsilon \rangle \rangle + \frac{2}{\varepsilon} \langle \langle v \hat{\gamma}_\varepsilon (1 - \hat{\gamma}_{\varepsilon*} \hat{\gamma}'_{\varepsilon*}) \mathcal{E}_\varepsilon \rangle \rangle \\ 38 \quad &+ \frac{1}{\varepsilon} \langle \langle (v + v_*) \hat{\gamma}_\varepsilon \hat{\gamma}_{\varepsilon*} \hat{\gamma}'_{\varepsilon*} \mathcal{E}_\varepsilon \rangle \rangle. \end{aligned} \quad 37$$

1 The first term is easily mastered by the entropy production estimate (11.15) and the fol- 1
 2 lowing classical estimate on the tail of Gaussian integrals 2

$$3 \int_{\mathbb{R}^N} e^{-|v|^2/2} |v|^\alpha \mathbb{1}_{|v|^2 > R} dv = O(R^{(a+N)/2-1} e^{-R/2}) \text{ as } R \rightarrow +\infty. 4$$

5
 6
 7 Observe that the integrand in the third term has the same symmetries as the original col- 7
 8 lision integrand (before truncation in $|v|$ and renormalization). It is also mastered by a 8
 9 combination of the entropy production estimate (11.15) with the Gaussian tail estimate 9
 10 above. 10

11 The most difficult part in the analysis of the momentum conservation defect is by far the 11
 12 second term in the decomposition of $\mathbf{D}_\varepsilon^2(v)$ above. That it vanishes in $L^1_{\text{loc}}(dt dx)$ as $\varepsilon \rightarrow 0$ 12
 13 ultimately relies upon the following estimate. 13

14
 15 *Nonlinear compactness estimate.* 15

$$16 (1 + |v|)^\alpha \left(\frac{\sqrt{G_\varepsilon} - 1}{\varepsilon} \right)^2 \text{ is uniformly integrable on } [0, T] \times K \times \mathbb{R}^3 \quad (11.26) 17$$

18
 19
 20 for the measure $dt dx M dv$, for each $T > 0$ and each compact $K \subset \mathbb{R}^3$, where α is the 21
 22 relative velocity exponent that appears in the hard cut-off assumption (3.55) on the collision 22
 23 kernel b . 23

24 We shall not give further details on the proof that $\mathbf{D}_\varepsilon^2(v) \rightarrow 0$ in $L^1_{\text{loc}}(dt dx)$, which is 24
 25 based on the above nonlinear compactness estimate together with the entropy production 25
 26 bound (11.15). 26

27 Let us however say a few words on the nonlinear compactness estimate itself. The rela- 27
 28 tive entropy bound (11.11) is essentially as good as an $L^\infty(dt; L^2(M dv dx))$ bound on g_ε 28
 29 on the set of (t, x, v) 's such that $G_\varepsilon(t, x, v) = O(1)$. Elsewhere, it essentially reduces to 29
 30 an $O(\varepsilon)$ bound in $L^\infty(dt; L^1(M dv dx))$, which is quite not enough for the Navier–Stokes 30
 31 limit. This is why the first works on this limit *assumed* some variant of this nonlinear 31
 32 compactness estimate. For instance, in either [10] or [87], it was assumed that 32
 33

$$34 (1 + |v|^2) \frac{g_\varepsilon^2}{1 + G_\varepsilon} \text{ is uniformly integrable on } [0, T] \times K \times \mathbb{R}^3, \quad (11.27) 35$$

36
 37 whereas all that was known on this quantity was the estimate 37
 38

$$39 (1 + |v|^2) \frac{g_\varepsilon^2}{1 + G_\varepsilon} = O(|\ln \varepsilon|) \text{ in } L^1_{\text{loc}}(dt dx; L^1(M dv)) 40$$

41
 42 (see [10]). This led to a decomposition of the number density fluctuation as 43
 44

$$45 g_\varepsilon = g_\varepsilon^b + \varepsilon g_\varepsilon^\sharp, 45$$

1 where the “good” part of the fluctuation is

$$2 \quad g_\varepsilon^b = \frac{g_\varepsilon}{0.5 + 0.5G_\varepsilon} = O(1) \quad \text{in } L^\infty(dt; L^2(M dv dx)),$$

3 while the “bad” part is

$$4 \quad g_\varepsilon^\sharp = \frac{g_\varepsilon^2}{1 + G_\varepsilon} = O(1) \quad \text{in } L^\infty(dt; L^1(M dv dx)).$$

5 In later works – for instance in [107,109] and [61] – this decomposition was slightly mod-
6 ified as follows. Pick a bump function $\gamma \in C_c^\infty(\mathbb{R}_+^*)$ such that

$$7 \quad \gamma|_{[\frac{3}{4}, \frac{5}{4}]} \equiv 1, \quad \text{supp}(\gamma) \subset \left[\frac{1}{2}, \frac{3}{2}\right] \text{ and } 0 \leq \gamma \leq 1,$$

8 and define

$$9 \quad g_\varepsilon^b = g_\varepsilon \gamma(G_\varepsilon), \quad g_\varepsilon^\sharp = \frac{1 - \gamma(G_\varepsilon)}{\varepsilon} g_\varepsilon.$$

10 It was proved in [61] that

$$11 \quad |g_\varepsilon^b|^2 \text{ is uniformly integrable on } [0, T] \times K \times \mathbb{R}^3 \quad (11.28)$$

12 for the measure $dt dx M dv$, while

$$13 \quad g_\varepsilon^\sharp = O\left(\frac{1}{\ln|\ln \varepsilon|}\right) \quad \text{in } L_{loc}^1(dt dx; L^1(M dv)). \quad (11.29)$$

14 Observe the difference between these last two controls and (11.27): with the new defin-
15 ition of g_ε^b and g_ε^\sharp , it is no longer true that $|g_\varepsilon^b|^2 \leq C g_\varepsilon^\sharp$, while $(\frac{g_\varepsilon}{1+G_\varepsilon})^2 \leq \frac{g_\varepsilon^2}{1+G_\varepsilon}$, so that
16 (11.27) actually entailed that the square of the good part in the old flat-sharp decomposi-
17 tion is uniformly integrable, even with a quadratic weight in v . In fact, the techniques in
18 [61] did not allow adding a quadratic weight in v as in (11.27), so that this compactness
19 assumption remained unproved; fortunately, it was possible to complete the proof of the
20 Navier–Stokes limit for cut-off Maxwell molecules with only the bounds (11.28)–(11.29),
21 and the weighted estimate

$$22 \quad (1 + |v|)^s (1 - \gamma(G_\varepsilon)) = O\left(\frac{\varepsilon^2}{\sqrt{\ln|\ln \varepsilon|}}\right) \quad \text{in } L_{loc}^1(dt dx; L^1(M dv)). \quad (11.30)$$

23 This control shows that the set where the bad part of the number density fluctuation domi-
24 nates is small in weighted v -space. There is a definite lack of symmetry between the con-
25 trols (11.29), bearing on large values of g_ε , and (11.30), bearing on large $|v|$'s. This lack
26 of symmetry is remedied in the most recent variant (11.26) of these nonlinear compactness
27 estimates (see [62]), we shall return to this when sketching the proof of (11.26).
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

11.4.5. The asymptotic momentum flux. With the vanishing of conservation defects (Step 2 in the proof of the Navier–Stokes limit) settled in the previous section, we turn our attention to Step 3, i.e., passing to the limit in the divergence of the momentum flux modulo gradients. This is by far the most difficult part of our analysis, and does require several preparations. In the present subsection, we reduce the momentum flux to some asymptotic normal form, to which we eventually apply compactness results to be described later.

LEMMA 11.5. *Let Π be the $L^2(M dv)$ -orthogonal projection on $\text{Ker } \mathcal{L}$; then, under the same assumptions as in Theorem 11.3,*

$$\mathbf{F}_\varepsilon(A) = \left\langle A \left(\Pi \frac{\sqrt{G_\varepsilon} - 1}{\varepsilon} \right)^2 \right\rangle - 2 \left\langle \widehat{A} \frac{1}{\varepsilon^2} \mathcal{Q}_M(\sqrt{G_\varepsilon}, \sqrt{G_\varepsilon}) \right\rangle + o(1)_{L^1_{\text{loc}}(dt dx)},$$

where we recall that the tensor field \widehat{A} is defined by

$$\widehat{A} \perp \text{Ker } \mathcal{L}_M \quad \text{and} \quad \mathcal{L}_M(\widehat{A}) = A = v \otimes v - \frac{1}{3}|v|^2 I.$$

The proof of this lemma is based upon splitting the momentum flux as

$$\begin{aligned} \mathbf{F}_\varepsilon(A) &= \frac{1}{\varepsilon} \left\langle A_{K_\varepsilon} \gamma_\varepsilon \frac{G_\varepsilon - 1}{\varepsilon} \right\rangle \\ &= \left\langle A_{K_\varepsilon} \gamma_\varepsilon \left(\frac{\sqrt{G_\varepsilon} - 1}{\varepsilon} \right)^2 \right\rangle + \frac{2}{\varepsilon} \left\langle A_{K_\varepsilon} \gamma_\varepsilon \frac{\sqrt{G_\varepsilon} - 1}{\varepsilon} \right\rangle \\ &= \mathbf{F}_\varepsilon^1(A) + \mathbf{F}_\varepsilon^2(A), \end{aligned}$$

as a consequence of the elementary identity

$$\begin{aligned} \frac{1}{\varepsilon}(G_\varepsilon - 1) &= \frac{1}{\varepsilon}(\sqrt{G_\varepsilon} - 1)(\sqrt{G_\varepsilon} + 1) \\ &= \frac{1}{\varepsilon}(\sqrt{G_\varepsilon} - 1)^2 + \frac{2}{\varepsilon}(\sqrt{G_\varepsilon} - 1). \end{aligned}$$

Then, one applies the following corollary of the nonlinear compactness estimate (11.26).

COROLLARY 11.6. *Under the same assumptions as in Theorem 11.3,*

$$\frac{\sqrt{G_\varepsilon} - 1}{\varepsilon} - \Pi \frac{\sqrt{G_\varepsilon} - 1}{\varepsilon} \rightarrow 0 \quad \text{in } L^2(dt dx; L^2((1 + |v|^\alpha)M dv))$$

as $\varepsilon \rightarrow 0$.

1 With the corollary above, one can show that the term $\mathbf{F}_\varepsilon^1(A)$ in the decomposition of the
 2 momentum flux is asymptotically close to

$$3 \left\langle A \left(\Pi \frac{\sqrt{G_\varepsilon} - 1}{\varepsilon} \right)^2 \right\rangle$$

4 (notice that the high velocity truncation is disposed of since $\Pi(\sqrt{G_\varepsilon} - 1)/\varepsilon$ has at most
 5 polynomial growth in v as $|v| \rightarrow +\infty$). That the second term $\mathbf{F}_\varepsilon^2(A)$ is close to

$$6 \left\langle \widehat{A} \frac{1}{\varepsilon^2} \mathcal{Q}_M(\sqrt{G_\varepsilon}, \sqrt{G_\varepsilon}) \right\rangle$$

7 uses Lemma 6.2.

8 Next, we explain how Lemma 11.5 is used in the proof of the Navier–Stokes limit. To
 9 begin with, since

$$10 \frac{\sqrt{G_\varepsilon} - 1}{\varepsilon} \simeq \frac{1}{2} g_\varepsilon \gamma_\varepsilon,$$

11 one has

$$12 \left\langle A \left(\Pi \frac{\sqrt{G_\varepsilon} - 1}{\varepsilon} \right)^2 \right\rangle \simeq \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle \otimes \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle - \frac{1}{3} |\langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle|^2 I.$$

13 On the other hand, the entropy production estimate (11.15) implies that, modulo extraction
 14 of a subsequence, one has

$$15 \frac{1}{\varepsilon^2} (\sqrt{G'_\varepsilon G'_{\varepsilon*}} - \sqrt{G_\varepsilon G_{\varepsilon*}}) \rightarrow q$$

16 in $w - L^2(dt dx d\mu)$. Passing to the limit in the scaled, renormalized Boltzmann equa-
 17 tion (11.18) entails the relation

$$18 \iint_{\mathbb{R}^3 \times \mathbb{S}^2} q b(v - v_*, \omega) M dv_* d\omega$$

$$19 = v \cdot \nabla_x g = \frac{1}{2} A : \nabla_x u + \text{terms that are odd in } v.$$

20 Eventually we arrive at the following asymptotic form of the momentum flux.

21 PROPOSITION 11.7. *Under the same assumptions as in Theorem 11.3, one has*

$$22 \mathbf{F}_\varepsilon(A) = \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle \otimes \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle - \frac{1}{3} |\langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle|^2 I$$

$$23 - v(\nabla_x u + (\nabla_x u)^T) + o(1)_{L^1_{\text{loc}}(dt dx)},$$

1 where

$$2 \quad u = \langle vg \rangle \quad \text{and} \quad g = \lim_{\varepsilon \rightarrow 0} g_\varepsilon \quad \text{in } w - L^1_{\text{loc}}(dt dx; L^1(M dv)).$$

3
4
5 **11.4.6. Strong compactness of $\langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle$.** In order to pass to the limit in the quadratic
6 term $\langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle$ and to conclude that

$$7 \quad P \operatorname{div}_x \left(\langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle \otimes \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle - \frac{1}{3} |\langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle|^2 I \right)$$

$$8 \quad \rightarrow P \operatorname{div}_x \left(u \otimes u - \frac{1}{3} |u|^2 I \right)$$

9
10
11 in the sense of distributions on $\mathbb{R}_+^* \times \mathbb{R}^3$ as $\varepsilon \rightarrow 0$, the weak convergence properties of g_ε
12 established so far are clearly insufficient. One needs instead some *strong* compactness
13 properties on the family $\langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle$.

14 (a) *Strong compactness in the x -variable.* Velocity averaging is the natural way to obtain
15 compactness in the space variable x for kinetic equations in the parabolic scaling (11.6).

16 For the purpose of studying the compactness of $\langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle$ in the x -variable, we use the
17 following variant of the L^2 -based velocity averaging theorem.

18
19 LEMMA 11.8. *Let ϕ_ε be a bounded family in $L^2_{\text{loc}}(dt dx; L^2(M dv))$ such that $|\phi_\varepsilon|^2$ is
20 locally uniformly integrable on $\mathbb{R}_+^* \times \mathbb{R}^3 \times \mathbb{R}^3$ for the Lebesgue measure. Assume that*

$$21 \quad (\varepsilon \partial_t + v \cdot \nabla_x) \phi_\varepsilon \text{ is bounded in } L^1_{\text{loc}}(dt dx dv).$$

22
23 Then, for each $\psi \in L^2(M dv)$, the family $\langle \phi_\varepsilon \psi \rangle$ is relatively compact in $L^2_{\text{loc}}(dt dx)$ with
24 respect to the x -variable, meaning that, for each $T > 0$ and each compact $K \subset \mathbb{R}^3$, one
25 has

$$26 \quad \iint_{[0, T] \times K} |\langle \phi_\varepsilon \psi \rangle(t, x + y) - \langle \phi_\varepsilon \psi \rangle(t, x)|^2 dt dx \rightarrow 0$$

27 as $y \rightarrow 0$ uniformly in ε .

28 See [61] for the proof, which is somewhat similar to the L^1 case of velocity averaging
29 recalled in Section 8.2.2 (especially Proposition ??).

30 Now, we apply the lemma above to

$$31 \quad \phi_\varepsilon = \frac{\sqrt{\varepsilon^c + G_\varepsilon} - 1}{\varepsilon}$$

32 since

$$33 \quad (\varepsilon \partial_t + v \cdot \nabla_x) \phi_\varepsilon = \frac{1}{\varepsilon^2} \frac{Q(G_\varepsilon, G_\varepsilon)}{2\sqrt{\varepsilon^c + G_\varepsilon}} = O(1)_{L^1_{\text{loc}}(dt dx dv)}$$

for $c \in (1, 2)$, by the entropy production estimate (11.15). Since

$$\frac{\sqrt{\varepsilon^c + G_\varepsilon} - 1}{\varepsilon} \simeq \frac{1}{2} g_\varepsilon \gamma_\varepsilon,$$

applying the velocity averaging lemma above leads to the following compactness (“in the x -variable”) result.

PROPOSITION 11.9. *Under the same assumptions as in Theorem 11.3, for each $T > 0$ and $K \subset \mathbb{R}^3$ compact, one has*

$$\iint_{[0, T] \times K} |\langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle(t, x + y) - \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle(t, x)|^2 dt dx \rightarrow 0$$

uniformly in ε as $y \rightarrow 0$.

(b) *Strong compactness in the t -variable.* It remains to obtain compactness in the time variable. As we shall see, the solenoidal part of $\langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle$ is strongly compact in the t -variable, but its orthogonal complement, which is a gradient field, is not because of high frequency oscillations in t .

PROPOSITION 11.10. *Under the assumptions of Theorem 11.3, modulo extraction of a subsequence, one has*

$$P \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle \rightarrow u$$

in $C(\mathbb{R}_+; w - L_x^2)$ and in $L_{loc}^2(dt dx)$ as $\varepsilon \rightarrow 0$.

PROOF. Indeed, Proposition 11.9 and the translation invariance of the Leray projection P imply that

$$\iint_{[0, T] \times K} |P \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle(t, x + y) - P \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle(t, x)|^2 dt dx \rightarrow 0 \quad (11.31)$$

uniformly in ε as $y \rightarrow 0$. On the other hand, the conservation law (11.25) implies that

$$\partial_t \int_{\mathbb{R}^3} P \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle \cdot \xi dx = O(1) \quad \text{in } L_{loc}^1(dt), \quad (11.32)$$

for each compactly supported, solenoidal vector field $\xi \in H^3(\mathbb{R}^3)$, since we know from Lemma 11.5 and the bounds (11.14) and (11.15), that $\mathbf{F}_\varepsilon(A)$ is bounded in $L_{loc}^1(dt dx)$. Also,

$$g_\varepsilon \gamma_\varepsilon \leq (1 + \sqrt{2}) \frac{\sqrt{G_\varepsilon} - 1}{\varepsilon}$$

1 so that (11.14) implies that

$$2 \quad \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle = O(1) \quad \text{in } B(\mathbb{R}_+; L^2(\mathbb{R}^3)), \quad (11.33) \quad 3$$

4 where $B(X, Y)$ denotes the class of bounded maps from X to Y .

5 Since the class of H^3 , compactly supported solenoidal vector fields is dense in that of
6 all H^3 solenoidal vector fields (see Appendix A of [86]), (11.33) and (11.32) imply that
7

$$8 \quad P \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle \text{ is relatively compact in } C(\mathbb{R}_+; w - L^2(\mathbb{R}^3)), \quad (11.34) \quad 9$$

10 by a variant of Ascoli's theorem that can be found in Appendix C of [86].

11 As for the $L^2_{\text{loc}}(dt dx)$ compactness, observe that (11.34) implies that

$$12 \quad P \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle \star \chi_\delta \text{ is relatively compact in } L^2_{\text{loc}}(dt dx), \quad 13$$

14 where χ_δ designates any mollifying sequence and \star is the convolution in the x -variable
15 only. Hence

$$16 \quad P \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle \cdot P \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle \star \chi_\delta \rightarrow Pu \cdot Pu \star \chi_\delta \quad 17$$

18 in $w - L^1_{\text{loc}}(dt dx)$ as $\varepsilon \rightarrow 0$. By (11.31),

$$19 \quad P \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle \star \chi_\delta \rightarrow P \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle \quad 20$$

21 in $L^2_{\text{loc}}(dt dx)$ uniformly in ε as $\delta \rightarrow 0$. With this, we conclude that

$$22 \quad |P \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle|^2 \rightarrow |Pu|^2 \quad \text{in } w - L^1_{\text{loc}}(dt dx) \quad 23$$

24 which implies that $P \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle \rightarrow Pu$ strongly in $L^2_{\text{loc}}(dt dx)$. \square 25

26 Next, consider

$$27 \quad \nabla_x \pi_\varepsilon = \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle - P \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle. \quad 28$$

29 Since

$$30 \quad \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle \rightarrow u \quad \text{in } w - L^2_{\text{loc}}(dt dx) \quad \text{and} \quad \text{div}_x u = 0 \quad 31$$

32 one has

$$33 \quad \nabla_x \pi_\varepsilon \rightarrow 0 \quad \text{in } w - L^2_{\text{loc}}(dt dx) \quad (11.35) \quad 34$$

35 as $\varepsilon \rightarrow 0$. Decompose then

$$36 \quad \begin{aligned} 37 \quad & P \text{div}_x (\langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle \otimes \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle) \\ 38 \quad & = P \text{div}_x (P \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle \otimes P \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle) + P \text{div}_x (P \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle \otimes \nabla_x \pi_\varepsilon) \\ 39 \quad & + P \text{div}_x (\nabla_x \pi_\varepsilon \otimes P \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle) + P \text{div}_x (\nabla_x \pi_\varepsilon \otimes \nabla_x \pi_\varepsilon). \end{aligned} \quad 40$$

1 By Proposition 11.10, the first term converges to $P \operatorname{div}_x(u \otimes u)$ in the sense of distribu- 1
 2 tions, while the second and third terms converge to 0 in the sense of distributions because 2
 3 of (11.35). 3

4 As for the last term, let $\zeta_\delta = \xi_\delta \star \xi_\delta \star \xi_\delta$, where ξ_δ is an approximate identity. One can 4
 5 prove that 5

$$\begin{aligned} 6 & \varepsilon \partial_t \zeta_\delta \star_x \nabla_x \pi_\varepsilon + \nabla_x \zeta_\delta \star_x \left\langle \frac{1}{3} |v|_{K_\varepsilon}^2 g_\varepsilon \gamma_\varepsilon \right\rangle \rightarrow 0 \quad \text{in } L^1_{\text{loc}}(\mathbb{R}_+; H^1_{\text{loc}}(\mathbb{R}^3)), \\ 7 & \varepsilon \partial_t \zeta_\delta \star_x \left\langle \frac{1}{3} |v|_{K_\varepsilon}^2 g_\varepsilon \gamma_\varepsilon \right\rangle + \frac{5}{3} \Delta_x \zeta_\delta \star_x \pi_\varepsilon \rightarrow 0 \quad \text{in } L^1_{\text{loc}}(\mathbb{R}_+; H^1_{\text{loc}}(\mathbb{R}^3)) \end{aligned}$$

8 as a result of (11.22), the vanishing of momentum and energy conservation defects 8
 9 (see Proposition 11.4 for the momentum, and proceed analogously for the energy) 9
 10 and the fact that $\mathbf{F}_\varepsilon(A)$ is bounded in $L^1_{\text{loc}}(dt dx)$ (see Lemma 11.5, and the bounds 10
 11 (11.14) and (11.15)). From the above system, Lions and Masmoudi observed in [87] that 11

$$\begin{aligned} 12 & \operatorname{div}_x(\nabla_x \zeta_\delta \star_x \pi_\varepsilon \otimes \nabla_x \zeta_\delta \star_x \pi_\varepsilon) \\ 13 & = \frac{1}{2} \nabla_x \left(|\nabla_x \zeta_\delta \star_x \pi_\varepsilon|^2 - \frac{5}{3} \zeta_\delta \star_x \left\langle \frac{1}{3} |v|_{K_\varepsilon}^2 g_\varepsilon \gamma_\varepsilon \right\rangle^2 \right) + o(1)_{L^1_{\text{loc}}(dt dx)}. \end{aligned}$$

14 Together with the uniform compactness “in the x -variable” proved in Proposition 11.9, this 14
 15 implies that 15
 16

$$17 \quad P \operatorname{div}_x(\nabla_x \pi_\varepsilon \otimes \nabla_x \pi_\varepsilon) \rightarrow 0 \quad 17$$

18 in the sense of distributions. Collecting the observations above, we have just proved the 18
 19 following proposition. 19
 20

21 **PROPOSITION 11.11.** *Under the assumptions of Theorem 11.3, modulo extraction of a 21
 22 subsequence, one has* 22

$$23 \quad P \operatorname{div}_x(\langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle \otimes \langle v_{K_\varepsilon} g_\varepsilon \gamma_\varepsilon \rangle) \rightarrow P \operatorname{div}_x(u \otimes u) \quad 23$$

24 *in the sense of distributions on $\mathbb{R}_+^* \times \mathbb{R}^3$ as $\varepsilon \rightarrow 0$.* 24
 25

26 With this proposition, we have completed all three steps in the proof of the Navier– 26
 27 Stokes limit (Theorem 11.3). 27
 28

29 **11.5. The nonlinear compactness estimate** 29

30 It only remains to prove the nonlinear compactness estimate (11.26), on which the two most 30
 31 important steps in the proof of the Navier–Stokes limit – i.e., the vanishing of conservation 31
 32 defects and the limiting form of the momentum flux – are based. 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45

1 This nonlinear compactness estimate results from new ideas on velocity averaging in L^1 ; 1
 2 consistently with our presentation of this subject in Section 8.2.2, we shall describe these 2
 3 ideas on the steady transport equation. 3

4 We start with a definition of the notion of partial uniform integrability in a product space. 4

5
 6 DEFINITION 11.12. Let μ and ν be positive Borel measures on \mathbb{R}^N ; we say that a fam- 6
 7 ily ϕ_ε of elements of $L^1(\mathbb{R}_x^N \times \mathbb{R}_y^N; d\mu(x) d\nu(y))$ is uniformly integrable in the variable y if 7
 8

$$9 \int_{\mathbb{R}^N} \left(\sup_{\nu(A) < \eta} \int_A |\phi_\varepsilon(x, y)| d\nu(y) \right) d\mu(x) \rightarrow 0$$

10
 11 as $\eta \rightarrow 0$, uniformly in ε . 11

12
 13 EXAMPLE. Assume that ν is a finite measure; then, for each $p > 1$, any bounded family 14
 15 in $L^1(d\mu(x); L^p(d\nu(y)))$ is uniformly integrable in y . Indeed, if ϕ_ε is any such family, one 15
 16 has 16

$$17 \int_{\mathbb{R}^N} \left(\sup_{\nu(A) < \eta} \int_A |\phi_\varepsilon(x, y)| d\nu(y) \right) d\mu(x) \leq \eta^{1/p'} \int_{\mathbb{R}^N} \|\phi_\varepsilon(x, \cdot)\|_{L^q(d\nu)} dx,$$

18
 19 where $p' = p/(p - 1)$, by applying Hölder's inequality to the inner integral. 20
 21

22 As usual, we shall say that a family $\phi_\varepsilon \in L^1(\mathbb{R}_x^N \times \mathbb{R}_y^N; d\mu(x) d\nu(y))$ is locally uni- 22
 23 formly integrable in y if, for each compact $K \subset \mathbb{R}^N \times \mathbb{R}^N$, the family $\mathbb{1}_K \phi_\varepsilon$ is uniformly 24
 25 integrable in y . 25

26 With this notion of partial uniform integrability, we can formulate an important improve- 26
 27 ment of the L^1 -variant of velocity averaging stated in Proposition 8.6 and Theorem 8.7. 27

28 THEOREM 11.13 (Golse and Saint-Raymond [60]). Let f_ε be a bounded family in 28
 29 $L^1_{\text{loc}}(\mathbb{R}^N \times \mathbb{R}^N; dx dy)$ such that 29

- 30 • the family $v \cdot \nabla_x f_\varepsilon$ is bounded in $L^1_{\text{loc}}(\mathbb{R}^N \times \mathbb{R}^N; dx dy)$, and 30
- 31 • the family f_ε is locally uniformly integrable in the variable v . 31

32 Then 32

- 33 (1) the family f_ε is locally uniformly integrable (in both variables (x, v)), and 33
- 34 (2) for each compactly supported $\phi \in L^\infty(\mathbb{R}^N)$, the family of averages 34

$$35 \int_{\mathbb{R}^N} f_\varepsilon(x, v) \phi(v) dv$$

36
 37 is relatively compact in $L^1_{\text{loc}}(\mathbb{R}^N; dx)$. 38
 39

40
 41 This result amplifies an earlier remark by Saint-Raymond who observed in [107] that, 41
 42 under the extra assumption that f_ε is bounded in $L^1_x(L^\infty_y)$, the family of averages 42
 43

$$44 \int_{\mathbb{R}^N} f_\varepsilon(x, v) \phi(v) dv$$

45

1 is locally uniformly integrable. 1

2
3 SKETCH OF THE PROOF OF THEOREM 11.13. We shall explain how to prove Saint-
4 Raymond's result under the assumptions of the theorem above. 3

5
6 *Step 1.* Let $\chi \equiv \chi(t, x, v)$ be the solution of the free transport equation 6

$$\begin{aligned} 7 \quad \partial_t \chi + v \cdot \nabla_x \chi &= 0, \quad (x, v) \in \mathbb{R}^N \times \mathbb{R}^N, t > 0, \\ 8 \quad \chi(0, x, v) &= \mathbb{1}_A(x), \quad (x, v) \in \mathbb{R}^N \times \mathbb{R}^N. \end{aligned}$$

9
10
11 Clearly, $\chi(t, x, v) = \mathbb{1}_A(x - tv)$ for each $t > 0$; it can therefore be put in the form 11

$$12 \quad \chi(t, x, v) = \mathbb{1}_{A_{t,x}}(v), \quad t > 0, (x, v) \in \mathbb{R}^N \times \mathbb{R}^N, \quad 12$$

13
14
15 where, for each $t > 0$, 15

$$16 \quad A_{t,x} = \{v \in \mathbb{R}^N \mid x - tv \in A\}. \quad 16$$

17
18
19 Further, $A_{t,x}$ is measurable and, for each $t > 0$ and $x \in \mathbb{R}^N$, one has 19

$$20 \quad |A_{t,x}| = \int_{\mathbb{R}^N} \chi(t, x, v) dv = \int_{\mathbb{R}^N} \mathbb{1}_A(x - tv) dv = \frac{1}{t^N} \int_{\mathbb{R}^N} \mathbb{1}_A(z) dz = \frac{|A|}{t^N}. \quad 20$$

21
22
23
24 *Step 2.* Without loss of generality, assume that f_ε and ϕ_ε in the statement of the theorem
25 are nonnegative, and that all the f_ε 's are supported in the same compact K of $\mathbb{R}^N \times \mathbb{R}^N$.
26 Then 26

$$\begin{aligned} 27 \quad & \int_A \int_{\mathbb{R}^N} f_\varepsilon(x, v) \phi(v) dv dx \quad 27 \\ 28 \quad &= \int_{\mathbb{R}^N} \int_{A_{t,x}} f_\varepsilon(x, v) \phi(v) dv dx \quad 28 \\ 29 \quad & - \int_0^t \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \chi(s, x, v) v \cdot \nabla_x f_\varepsilon(x, v) \phi(v) dx dv ds \quad 29 \\ 30 \quad & \quad \quad \quad (11.36) \quad 30 \\ 31 \quad & \quad \quad \quad 31 \\ 32 \quad & \quad \quad \quad 32 \\ 33 \quad & \quad \quad \quad 33 \\ 34 \quad & \quad \quad \quad 34 \end{aligned}$$

35
36 as can be seen by integrating by parts the second integral on the right-hand side of the
37 equality above. 37

38 Pick $\eta > 0$ arbitrarily small; the second integral on the right-hand side of (11.36) satisfies 38

$$39 \quad \left| \int_0^t \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \chi(s, x, v) v \cdot \nabla_x f_\varepsilon(x, v) \phi(v) dx dv ds \right| \leq t \|v \cdot \nabla_x f_\varepsilon\|_{L^1} \|\phi\|_{L^\infty} \quad 39$$

40
41
42 and therefore can be made less than η by choosing 42

$$43 \quad 0 < t < \frac{\eta}{1 + \|v \cdot \nabla_x f_\varepsilon\|_{L^1} \|\phi\|_{L^\infty}}. \quad 43$$

44
45

1 For this $t > 0$, the first integral on the right-hand side of (11.36) satisfies

$$2 \int_{\mathbb{R}^N} \int_{A_{t,x}} f_\varepsilon(x, v) \phi(v) dv dx \rightarrow 0$$

3 as $|A| \rightarrow 0$ uniformly in ε , since f_ε is uniformly integrable in v and $|A_{t,x}| = |A|/t^N$, as
 4 established in Step 1.

5 Therefore, for each $\eta > 0$, there exists $\alpha > 0$ such that $|A| < \alpha$ implies that

$$6 \int_A \int_{\mathbb{R}^N} f_\varepsilon(x, v) \phi(v) dv dx \leq 2\eta$$

7 uniformly in $\varepsilon > 0$, which entails that the family of averages

$$8 \int_{\mathbb{R}^N} f_\varepsilon(x, v) \phi(v) dv$$

9 is uniformly integrable. □

10 The nonlinear compactness estimate (11.26) will of course be obtained from state-
 11 ment (1) in Theorem 11.13. In fact, we first observe that, for each $c \in (1, 2)$,

$$12 \phi_\varepsilon^\delta = \left(\frac{\sqrt{\varepsilon^c + G_\varepsilon} - 1}{\varepsilon} \right)^2 \gamma \left(\varepsilon \delta \left(\frac{\sqrt{\varepsilon^c + G_\varepsilon} - 1}{\varepsilon} \right) \right)$$

13 satisfies

$$14 \phi_\varepsilon^\delta = O(1) \quad \text{in } L_t^\infty(L^1(M dv dx)),$$

15 while

$$16 (\varepsilon \partial_t + v \cdot \nabla_x) \phi_\varepsilon^\delta = O(1) \quad \text{in } L_{loc}^1(dt dx M dv).$$

17 We next let $\delta \rightarrow 0$ and remove the ε^c from under the square root (in that order) so that

$$18 \phi_\varepsilon^\delta \simeq \left(\frac{\sqrt{G_\varepsilon} - 1}{\varepsilon} \right)^2.$$

19 In order to apply Theorem 11.13 to ϕ_ε^δ , it remains to prove that this family is uniformly
 20 integrable in the v -variable. In fact, we prove the following results.

21 **PROPOSITION 11.14.** *Under the assumptions of Theorem 11.3, for each $T > 0$ and each
 22 compact $K \subset \mathbb{R}^3$, the family*

$$23 (1 + |v|)^\alpha \left(\frac{\sqrt{G_\varepsilon} - 1}{\varepsilon} \right)^2$$

1 *is uniformly integrable on $[0, T] \times K \times \mathbb{R}^3$ for the measure $dt dx M dv$. (We recall that α*
 2 *is the relative velocity exponent in the hard cut-off assumption on the collision kernel b ,*
 3 *see (3.55).)*

4
 5 This proposition improves upon the result in [61], that applied to cut-off Maxwell mole-
 6 cules only (i.e., to the case where $\alpha = 0$). Its proof is fairly technical, so that we shall just
 7 sketch the main idea in it.

8 Start from the identity

$$\begin{aligned} & \mathcal{L}_M \left(\frac{\sqrt{G_\varepsilon} - 1}{\varepsilon} \right) \\ &= \varepsilon \mathcal{Q}_M \left(\frac{\sqrt{G_\varepsilon} - 1}{\varepsilon}, \frac{\sqrt{G_\varepsilon} - 1}{\varepsilon} \right) - \frac{1}{\varepsilon} \mathcal{Q}_M(\sqrt{G_\varepsilon}, \sqrt{G_\varepsilon}). \end{aligned} \quad (11.37)$$

15 Next, we recall the Bardos–Caffisch–Nicolaeenco spectral gap for \mathcal{L}_M in weighted space
 16 (see Theorem 3.11)

$$\langle \phi \mathcal{L}_M \phi \rangle \geq C_0 \langle (1 + |v|)^\alpha \phi^2 \rangle, \quad \phi \in (\text{Ker } \mathcal{L}_M)^\perp$$

20 together with the following continuity estimate for \mathcal{Q} (see [57])

$$\| \mathcal{Q}_M(\phi, \phi) \|_{L^2((1+|v|)^{-\alpha} M dv)} \leq C \| \phi \|_{L^2(M dv)} \| \phi \|_{L^2((1+|v|)^\alpha M dv)}.$$

24 Using both estimates in the identity above leads to the following control

$$\begin{aligned} & \left(1 - O(\varepsilon) \left\| \frac{\sqrt{G_\varepsilon} - 1}{\varepsilon} \right\|_{L^2(M dv)} \right) \left\| \frac{\sqrt{G_\varepsilon} - 1}{\varepsilon} - \Pi \frac{\sqrt{G_\varepsilon} - 1}{\varepsilon} \right\|_{L^2((1+|v|)^\alpha M dv)} \\ & \leq O(\varepsilon)_{L^2_{t,x}} + O(\varepsilon) \left\| \frac{\sqrt{G_\varepsilon} - 1}{\varepsilon} \right\|_{L^2(M dv)}^2. \end{aligned}$$

32 This control suggests that

$$\frac{\sqrt{G_\varepsilon} - 1}{\varepsilon} \text{ is close to its hydrodynamic projection } \Pi \frac{\sqrt{G_\varepsilon} - 1}{\varepsilon}$$

37 precisely in that weighted L^2 space that appears in the statement of Proposition 11.14.
 38 Since the hydrodynamic projection $\Pi \frac{\sqrt{G_\varepsilon} - 1}{\varepsilon}$ is as regular in v as one can hope for (being
 39 a quadratic polynomial in v), this eventually entails uniform integrability in v once the
 40 difficulties related to the (t, x) dependence in the estimate above have been handled. As
 41 we already indicated above, the remaining part of the proof is too technical to be described
 42 here, and we refer the interested reader to [62] for a complete argument.

43 Instead of giving more details on this proof, we comment on the differences between
 44 the nonlinear compactness estimate (11.28)–(11.30) obtained in [61] in the case of cut-
 45 off Maxwell molecules, and the most recent variant of such controls obtained in [62],

namely (11.26). In fact, the main difference between these controls lies in the method for obtaining uniform integrability in v .

In [61], the decomposition (11.37) was replaced with

$$G_\varepsilon = \left(G_\varepsilon - \frac{\mathcal{A}^+(MG_\varepsilon, MG_\varepsilon)}{M\langle G_\varepsilon \rangle} \right) + \frac{\mathcal{A}^+(MG_\varepsilon, MG_\varepsilon)}{M\langle G_\varepsilon \rangle} \quad (11.38)$$

(in fact, with a more technical variant of (11.38) involving several truncations), where \mathcal{A}^+ was the gain part of a fictitious collision operator

$$\mathcal{A}^+(\phi, \phi) = \iint_{\mathbb{R}^3 \times \mathbb{S}^2} \phi' \phi'_* |\cos(v - v_*, \omega)| dv_* d\omega.$$

Roughly speaking the term

$$\left(G_\varepsilon - \frac{\mathcal{A}^+(MG_\varepsilon, MG_\varepsilon)}{M\langle G_\varepsilon \rangle} \right)$$

was easily controlled by the entropy production bound, while the term

$$\frac{\mathcal{A}^+(MG_\varepsilon, MG_\varepsilon)}{M\langle G_\varepsilon \rangle}$$

improved the regularity and decay in v due to estimates by Grad [63] and Caffisch [23] – that are linear analogues of the result by Lions on the smoothing properties of the gain part in the Boltzmann collision integral (Lemma 3.10).

However, the main drawback of the decomposition (11.38) was that it naturally involved the quantity

$$g_\varepsilon \gamma_\varepsilon + \frac{1 - \gamma_\varepsilon}{\varepsilon}$$

(see (2.17) in [61]), thereby leading to a dissymmetry between the roles of large v 's and large values of g_ε in the estimates (11.28)–(11.30). This, as we already mentioned, was an obstruction in upgrading (11.28)–(11.29) with a $(1 + |v|)^\alpha$ weight as in (11.26).

The method used in [62] essentially differs from that of [61]; to begin with, we do not attempt to establish that the number density F_ε is close in some sense to its associated local equilibrium M_{F_ε} (i.e., the Maxwellian with same local density, energy and momentum as F_ε), an idea that is very natural in the context of the BGK model of the Boltzmann equation, see [107, 109].

Instead, we seek to prove that some variant of the number density fluctuation, namely the quantity $\frac{\sqrt{G_\varepsilon} - 1}{\varepsilon} \simeq \frac{1}{2} g_\varepsilon$, becomes close to its associated infinitesimal Maxwellian, i.e., its $L^2(M dv)$ -orthogonal projection on $\text{Ker } \mathcal{L}_M$. This new approach naturally leads to using the Dirichlet form of the linearized collision operator, whose relative coercivity transversally to $\text{Ker } \mathcal{L}_M$ provides exactly the $(1 + |v|)^\alpha$ weight needed in the proof.

12. Conclusions and open problems

This survey of the Boltzmann equation and its hydrodynamic limits remains incomplete in several respects. As already mentioned in the Introduction, we have chosen to emphasize global theories (such as the DiPerna–Lions theory for the Boltzmann equation, or the Leray theory for the Navier–Stokes equations) for the evolution problem posed on a spatial domain without boundaries. More realistic formulations of the hydrodynamic limits of the Boltzmann equation would certainly involve boundary value problems. At the formal level, the influence of boundaries on hydrodynamic limits is discussed at length in Sone’s authoritative monograph [115].

More specifically, hydrodynamic limits of boundary value problems may, in some particular cases, involve boundary layer equations whose mathematical study is a subject of its own and has inspired a considerable amount of literature. These boundary layers are meant to forget the specific dependence of the boundary data upon the velocity variable so as to match it with the inner form of the number density, which, as explained in this survey, is a local Maxwellian or infinitesimal Maxwellian in the hydrodynamic limit. This is obviously not the place for a description of that theory, to begin with because these boundary layer are essentially steady (instead of evolution) problems. The interested reader could start with Chapter 5, Section 5 of [27] and then get acquainted with the mathematical theory of half-space problems as exposed in [6,26,31,57,58,93] and [119]. These works deal with linearized, or at best, weakly nonlinear problems. Nonlinear boundary layer equations with nonzero net mass flux at the boundary – typically in the case of a gas condensing on, or evaporating from a solid boundary – have been studied in depth numerically by Sone, Aoki and their collaborators, see Chapter 7 of [115] and the references therein. At the formal level, their theory is expected to agree with a general theory of boundary conditions for the Euler system of compressible fluids, yet to be formulated at the time of this writing.

In some cases however, boundary layer are not leading order effects. This is the case of the incompressible Stokes or Navier–Stokes limit of the Boltzmann equation for a gas in a container with specular or diffuse reflection, or Maxwell’s accommodation condition at the boundary; see [96], which extends to such a boundary value problems the results in [53]. The analogue of the DiPerna–Lions theory for the boundary value problem is due to Mischler [97], this is by no means a straightforward extension of [37]. It requires rather subtle analytical tools because natural boundary conditions for the Boltzmann equation – except in the case of a specular reflection of gas molecule at the boundary – do not interact well with the renormalization procedure which is essential in controlling the collision integral.

Let us conclude with a few open problems, in addition to those already mentioned in the body of this article.

It would be of considerable interest to derive the global BV solutions constructed by T.-P. Liu from the Boltzmann equation. As in the case of the incompressible Euler limit of the Boltzmann equation, the entropy production bound entailed by Boltzmann’s H -theorem does not balance the action of the streaming operator on the number density: the compactness of hydrodynamic moments of the number density is probably to be sought in some stability property of BV solutions of the compressible Euler system. Most likely, such a theory should use Bressan’s remarkable results in that direction (see [19,20]).

As an indication of the level of difficulty of this problem, let us simply mention that the hydrodynamic limit of the Boltzmann equation leading to a single Riemann problem has not been established so far, except in the case of a single, weak shock wave. Shock profiles for the Boltzmann equation were formally discussed in [25]; for the hard sphere gas, a complete construction of these profiles in the weakly nonlinear regime can be found in a series of important papers by Nicolaenko [99–101]. All these constructions were based on a deep understanding of the algebra of the linearized Boltzmann equation in connection with the underlying saddle point structure of subsonic states in gas dynamics. They suggested some topological structure that remains to be unraveled and could ultimately lead to a construction of shock profiles of arbitrary strength. The extension to other cut-off molecular interactions is due to Caflisch and Nicolaenko [24]. However, all these constructions led to solutions of the Boltzmann equation whose positivity was not established. This latter problem was solved only very recently by Liu and Yu [90].

Another open problem would be to improve Theorem ??, by relaxing the unphysical assumption made on the size of the number density fluctuations δ_ε to reach the physically natural condition that $\delta_\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0$. This would probably require more information on the local conservation laws of momentum and energy for renormalized solutions of the Boltzmann equation. Such information would most likely be an important prerequisite for making progress on the compressible Euler limit.

Finally, we have only considered evolution problems in this survey. Although steady problems are beyond the scope of the present book, they are perhaps even more important in some applications (such as aerodynamics). For instance, it is well known that, for any divergence free force field $f \equiv f(x) \in L^2(\Omega; \mathbb{R}^3)$, the steady incompressible Navier–Stokes equations in a smooth, bounded open domain $\Omega \subset \mathbb{R}^3$

$$\begin{aligned} -\nu \Delta_x u &= f - \nabla_x p - (u \cdot \nabla_x)u, & \operatorname{div}_x u &= 0, & x &\in \Omega, \\ u|_{\partial\Omega} &= 0 \end{aligned} \tag{12.1}$$

has at least one classical solution $u \equiv u(x) \in H^2(\Omega, \mathbb{R}^3)$, obtained by a Leray–Schauder fixed point argument (see, for instance, [74]). Unfortunately, the parallel theory for the Boltzmann equation is not as advanced: for one thing, at the time of this writing, there is no analogue of the DiPerna–Lions result for the steady Boltzmann equation with a prescribed external force field. The reader interested in those matters is advised to read the survey article by Maslova [94]; see also her book [95]; among classical references on the subject, there are some papers by Guiraud (see [65–67]), and more recent work by Arkeryd and Nouri (see, for instance, [3]). See also [117] for the case of a gas flow modeled by the Boltzmann equation around a convex body. Finally, a rather exhaustive description of such steady problems and their hydrodynamic limits (at the formal level) may be found in Sone’s book [115].

In spite of the difficulties inherent to the steady Boltzmann equation, the fact that the solutions of (12.1) are more regular than in the case of the evolution problem could be of considerable help, at least in the context of the hydrodynamic limit.

References

- [1] V.I. Agoshkov, *Spaces of functions with differential difference characteristics and smoothness of the solutions of the transport equations*, Dokl. Akad. Nauk SSSR **276** (1984), 1289–1293.
- [2] K. Aoki, C. Bardos, F. Golse and Y. Sone, *Derivation of hydrodynamic limits from either the Liouville equation or kinetic models: Study of an example*, Mathematical Analysis of Liquids and Gases, Kyoto, 1999, A. Matsumura, ed., Sūrikaiseikikenkyūsho Kōkyūroku **1146** (2000), 154–181.
- [3] L. Arkeryd and A. Nouri, *The stationary Boltzmann equation in \mathbb{R}^n with given indata*, Ann. Sc. Norm. Sup. Pisa Cl. Sci. (5) **1** (2002), 359–385.
- [4] V.I. Arnold and B.A. Khesin, *Topological Methods in Hydrodynamics*, Springer-Verlag, New York (1998).
- [5] C. Bardos and C. Mouhot, *Explicit spectral gap estimates for the linearized Boltzmann and Landau operators with hard potentials*, Rev. Mat. Iberoamericana, to appear.
- [6] C. Bardos, R. Caflisch and B. Nicolaenko, *The Milne and Kramers problems for the Boltzmann equation of a hard sphere gas*, Comm. Pure Appl. Math. **39** (1986), 323–352.
- [7] C. Bardos and F. Golse, *Différents aspects de la notion d'entropie au niveau de l'équation de Boltzmann et de Navier–Stokes*, C. R. Math. Acad. Sci. Paris **299** (1984), 291–294.
- [8] C. Bardos, F. Golse and C.D. Levermore, *Sur les limites asymptotiques de la théorie cinétique conduisant à la dynamique des fluides incompressibles*, C. R. Math. Acad. Sci. Paris **309** (1989), 727–732.
- [9] C. Bardos, F. Golse and C.D. Levermore, *Fluid dynamic limits of the Boltzmann equation I*, J. Statist. Phys. **63** (1991), 323–344.
- [10] C. Bardos, F. Golse and C.D. Levermore, *Fluid dynamic limits of kinetic equations II: Convergence proofs for the Boltzmann equation*, Comm. Pure Appl. Math. **46** (1993), 667–753.
- [11] C. Bardos, F. Golse and C.D. Levermore, *The acoustic limit for the Boltzmann equation*, Arch. Ration. Mech. Anal. **153** (2000), 177–204.
- [12] C. Bardos, F. Golse, B. Perthame and R. Sentis, *The nonaccretive radiative transfer equations: Existence of solutions and Rosseland approximation*, J. Funct. Anal. **77** (1988), 434–460.
- [13] C. Bardos and C.D. Levermore, *Kinetic equations and an incompressible limit that recovers viscous heating*, Preprint.
- [14] C. Bardos and S. Ukai, *The classical incompressible Navier–Stokes limit of the Boltzmann equation*, Math. Models Methods Appl. Sci. **1** (1991), 235–257.
- [15] J.T. Beale, T. Kato and A.J. Majda, *Remarks on the breakdown of smooth solutions for the 3D Euler equations*, Comm. Math. Phys. **94** (1984), 61–66.
- [16] L. Boltzmann, *Lectures on Gas Theory*, S.G. Brush, transl. ed., University of California Press, Berkeley–Los Angeles, CA (1964).
- [17] F. Bouchut and L. Desvillettes, *A proof of the smoothing properties of the positive part of Boltzmann's kernel*, Rev. Mat. Iberoamericana **14** (1998), 47–61.
- [18] F. Bouchut, F. Golse and M. Pulvirenti, *Kinetic Equations and Asymptotic Theory*, L. Desvillettes and B. Perthame, eds, Ser. Appl. Math., Vol. 4, Gauthier-Villars, Paris (2000).
- [19] A. Bressan, *Hyperbolic Systems of Conservation Laws. The One-Dimensional Cauchy Problem*, Oxford University Press, Oxford (2000).
- [20] A. Bressan, *Hyperbolic systems of conservation laws in one space dimension*, Proc. Internat. Congress of Mathematicians, Vol. I, Higher Ed. Press, Beijing (2002), 159–178.
- [21] L. Caffarelli, R.V. Kohn and L. Nirenberg, *On the regularity of the solutions of Navier–Stokes equations*, Comm. Pure Appl. Math. **35** (1982), 771–831.
- [22] R.E. Caflisch, *The fluid dynamic limit of the nonlinear Boltzmann equation*, Comm. Pure Appl. Math. **33** (1980), 651–666.
- [23] R.E. Caflisch, *The Boltzmann equation with a soft potential. I. Linear, spatially-homogeneous*, Comm. Math. Phys. **74** (1980), 71–95.
- [24] R.E. Caflisch and B. Nicolaenko, *Shock profile solutions of the Boltzmann equation*, Comm. Math. Phys. **86** (1982), 161–194.
- [25] C. Cercignani, *Bifurcation problems in fluid mechanics*, Meccanica – J. Italian Assoc. Theoret. Appl. Mech. **5** (1970), 7–16.

- 1 [26] **C. Cercignani**, *Half-space problems in the kinetic theory of gases*, Trends in Applications of Pure Mathematics to Mechanics, E. Kröner and K. Kirchgässner, eds, Lecture Notes in Phys., Vol. 249, Springer-Verlag, Berlin (1986), 35–50. 1
- 2 [27] **C. Cercignani**, *The Boltzmann Equation and Its Applications*, Springer-Verlag, New York (1988). 2
- 3 [28] **C. Cercignani**, **R. Illner** and **M. Pulvirenti**, *The Mathematical Theory of Dilute Gases*, Springer-Verlag, 3
- 4 New York (1994). 4
- 5 [29] **J.-Y. Chemin**, *Perfect Incompressible Fluids*, Oxford University Press, New York (1998). 5
- 6 [30] **P. Constantin** and **C. Foias**, *Navier–Stokes Equations*, University of Chicago Press, Chicago, IL (1988). 6
- 7 [31] **F. Coron**, **F. Golse** and **C. Sulem**, *A classification of well-posed kinetic layer problems*, Comm. Pure Appl. 7
- 8 Math. **41** (1988), 409–435. 8
- 9 [32] **C. Dafermos**, *Hyperbolic Conservation Laws in Continuum Physics*, Springer-Verlag, Berlin–Heidelberg 9
- 10 (2000). 10
- 11 [33] **A. DeMasi** and **E. Presutti**, *Mathematical Methods for Hydrodynamic Limits*, Lecture Notes in Math., 11
- 12 Vol. 1501, Springer-Verlag, Berlin–Heidelberg (1991). 12
- 13 [34] **L. Desvillettes** and **F. Golse**, *A remark concerning the Chapman–Enskog asymptotics*, Advances in Kinetic 13
- 14 Theory and Computing, B. Perthame, ed., World Scientific, River Edge, NJ (1994), 191–209. 14
- 15 [35] **L. Desvillettes** and **C. Villani**, *On the trend to global equilibrium for spatially inhomogeneous kinetic 15*
- 16 systems: *The Boltzmann equation*, Preprint. 16
- 17 [36] **R.J. DiPerna** and **P.-L. Lions**, *Global weak solutions of the Vlasov–Maxwell system*, Comm. Pure Appl. 17
- 18 Math. **42** (1989), 729–757. 18
- 19 [37] **R.J. DiPerna** and **P.-L. Lions**, *On the Cauchy problem for the Boltzmann equation: Global existence and 19*
- 20 weak stability results, Ann. of Math. **130** (1990), 321–366. 20
- 21 [38] **R.J. DiPerna**, **P.-L. Lions** and **Y. Meyer**, *L^p regularity of velocity averages*, Ann. Inst. H. Poincaré Anal. 21
- 22 Non Linéaire **8** (1991), 271–287. 22
- 23 [39] **H.S. Dumas**, **L. Dumas** and **F. Golse**, *Remarks on the notion of mean free path for a periodic array of 23*
- 24 spherical obstacles, J. Statist. Phys. **87** (1997), 943–950. 24
- 25 [40] **R.S. Ellis** and **M.A. Pinsky**, *The first and second fluid approximations to the linearized Boltzmann equa- 25*
- 26 tion, J. Math. Pures Appl. **54** (1975), 125–156. 26
- 27 [41] **R. Esposito** and **M. Pulvirenti**, *From particles to fluids*, Preprint. 27
- 28 [42] **E. Feireisl**, *Dynamics of Viscous Compressible Fluids*, Oxford University Press, Oxford (2004). 28
- 29 [43] **K.O. Friedrichs** and **P.D. Lax**, *System of conservation equations with a convex extension*, Proc. Natl. Acad. 29
- 30 Sci. USA **68** (1971), 1686–1688. 30
- 31 [44] **J. Fritz**, *An Introduction to the Theory of Hydrodynamic Limits*, The University of Tokyo, Tokyo (2001). 31
- 32 [45] **G. Gallavotti**, *Foundations of Fluid Dynamics*, Springer-Verlag, Berlin (2002). 32
- 33 [46] **L. Garding**, *Cauchy’s problem for hyperbolic equations*, Lecture Notes, University of Chicago (1957). 33
- 34 [47] **P. Gérard**, *Moyennisation et régularité deux-microlocale*, Ann. Sci. École Norm. Sup. (4) **23** (1990), 34
- 35 89–121. 35
- 36 [48] **P. Gérard**, *Microlocal defect measures*, Comm. Partial Differential Equations **16** (1991), 1761–1794. 36
- 37 [49] **P. Gérard** and **F. Golse**, *Averaging regularity results for PDEs under transversality assumptions*, Comm. 37
- 38 Pure Appl. Math. **45** (1992), 1–26. 38
- 39 [50] **R.T. Glassey**, *The Cauchy Problem in Kinetic Theory*, SIAM, Philadelphia, PA (1996). 39
- 40 [51] **J. Glimm**, *Solutions in the large for nonlinear hyperbolic systems of equations*, Comm. Pure Appl. Math. 40
- 41 **18** (1965), 697–715. 41
- 42 [52] **S.K. Godunov**, *An interesting class of quasilinear systems*, Dokl. Akad. Nauk SSSR **139** (1961), 521–523. 42
- 43 [53] **F. Golse** and **C.D. Levermore**, *The Stokes–Fourier and acoustic limits for the Boltzmann equation*, Comm. 43
- 44 Pure Appl. Math. **55** (2002), 336–393. 44
- 45 [54] **F. Golse**, **C.D. Levermore** and **L. Saint-Raymond**, *La méthode de l’entropie relative pour les limites hy- 45*
- drodynamiques de modèles cinétiques, Séminaire: Equations aux Dérivées Partielles, 1999–2000, Exposé 40
- No. XIX, Ecole Polytechnique, Palaiseau (2000). 41
- [55] **F. Golse**, **P.-L. Lions**, **B. Perthame** and **R. Sentis**, *Regularity of the moments of the solution of a transport 42*
- equation, J. Funct. Anal. **76** (1988), 110–125. 43
- [56] **F. Golse**, **B. Perthame** and **R. Sentis**, *Un résultat de compacité pour les équations de transport et applica- 44*
- tion au calcul de la limite de la valeur propre principale de l’opérateur de transport, C. R. Math. Acad. 44
- Sci. Paris **301** (1985), 341–344. 45

- 1 [57] **F. Golse, B. Perthame and C. Sulem**, *On a boundary layer problem for the nonlinear Boltzmann equation*, 1
Arch. Ration. Mech. Anal. **103** (1988), 81–96. 2
- 3 [58] **F. Golse and F. Poupaud**, *Stationary solutions of the linearized Boltzmann equation in a half-space*, Math. 3
Methods Appl. Sci. **11** (1989), 483–502. 4
- 5 [59] **F. Golse and F. Poupaud**, *Limite fluide des équations de Boltzmann des semi-conducteurs pour une statis- 5
tique de Fermi–Dirac*, Asymptot. Anal. **6** (1992), 135–160 (in French). 6
- 7 [60] **F. Golse and L. Saint-Raymond**, *Velocity averaging in L^1 for the transport equation*, C. R. Math. Acad. 6
Sci. Paris **334** (2002), 557–562. 7
- 8 [61] **F. Golse and L. Saint-Raymond**, *The Navier–Stokes limit of the Boltzmann equation for bounded collision 8
kernels*, Invent. Math. **155** (2004), 81–161. 9
- 10 [62] **F. Golse and L. Saint-Raymond**, *The Navier–Stokes limit of the Boltzmann equation for hard potentials*, in 10
preparation. 11
- 12 [63] **H. Grad**, *Principles of the kinetic theory of gases*, Handbook der Physik, Bd. XII, S. Flügge, ed., Springer- 11
Verlag, Berlin (1958), 205–294. 12
- 13 [64] **H. Grad**, *Asymptotic theory of the Boltzmann equation II*, Rarefied Gas Dynamics, Vol. I, J.A. Laurmann, 13
ed., Academic Press, New York (1963), 26–59. 14
- 15 [65] **J.-P. Guiraud**, *Problème aux limites intérieur pour l’équation de Boltzmann linéaire*, J. Mécanique **9** 15
(1970), 443–490 (in French). 16
- 17 [66] **J.-P. Guiraud**, *Problème aux limites intérieures pour l’équation de Boltzmann*, Actes du Congrès Interna- 16
tional des Mathématiciens, Nice, 1970, Vol. 3, Gauthier-Villars, Paris (1971), 115–122 (in French). 17
- 18 [67] **J.-P. Guiraud**, *Problème aux limites intérieur pour l’équation de Boltzmann en régime stationnaire, faible- 18
ment non linéaire*, J. Mécanique **11** (1972), 183–231 (in French). 19
- 20 [68] **D. Hilbert**, *Mathematical problems*, Internat. Congress of Mathematicians, Paris, 1900; English transl.: 20
Bull. Amer. Math. Soc. **37** (2000), 407–436. 21
- 22 [69] **D. Hilbert**, *Begründung der kinetischen Gastheorie*, Math. Ann. **72** (1912), 562–577. 21
- 23 [70] **R. Illner and M. Shinbrot**, *The Boltzmann equation: Global existence for a rare gas in an infinite vacuum*, 22
Comm. Math. Phys. **95** (1984), 217–226. 23
- 24 [71] **T. Kato**, *Nonstationary flows of viscous and ideal fluids in \mathbb{R}^3* , J. Funct. Anal. **9** (1972), 296–305. 24
- 25 [72] **T. Kato**, *The Cauchy problem for quasi-linear symmetric hyperbolic systems*, Arch. Ration. Mech. Anal. 25
58 (1975), 181–205. 26
- 27 [73] **M. Lachowicz**, *On the initial layer and the existence theorem for the nonlinear Boltzmann equation*, Math. 27
Methods Appl. Sci. **9** (1987), 342–366. 28
- 29 [74] **O.A. Ladyzhenskaya**, *The Mathematical Theory of Viscous Incompressible Flows*, Gordon and Breach, 28
New York–London–Paris (1969). 29
- 30 [75] **L. Landau and E. Lifshitz**, *Course of Theoretical Physics, Vol. 6, Fluid Mechanics*, Pergamon Press, Oxford 30
(1987). 31
- 32 [76] **L. Landau, E. Lifshitz and L. Pitayevski**, *Course of Theoretical Physics, Vol. 10, Physical Kinetics*, Perg- 32
amon Press, Oxford (1981). 33
- 34 [77] **O.E. Lanford**, *Time evolution of large classical systems*, Dynamical Systems, Theory and Applications, 33
Lecture Notes in Phys., Vol. 38, Springer-Verlag, Berlin (1975), 1–111. 34
- 35 [78] **P.D. Lax**, *Hyperbolic systems of conservation laws and the mathematical theory of shock waves*, SIAM 35
Regional Conf. Lect. No. 11, Philadelphia, PA (1973). 36
- 37 [79] **J. Leray**, *Essai sur les mouvements plans d’un liquide visqueux que limitent des parois*, J. Math. Pures 37
Appl. **13** (1934), 331–418. 38
- 38 [80] **J. Leray**, *Essai sur le mouvement d’un liquide visqueux emplissant l’espace*, Acta Math. **63** (1934), 38
193–248. 39
- 40 [81] **J. Leray**, *Hyperbolic Differential Equations*, Lecture Notes, The Institute for Advanced Study, Princeton, 40
NJ (1953). 41
- 42 [82] **C.D. Levermore**, *Gas dynamics beyond Navier–Stokes*, in preparation. 42
- 43 [83] **P.-L. Lions**, *Conditions at infinity for Boltzmann’s equation*, Comm. Partial Differential Equations **19** 43
(1994), 335–367. 44
- 44 [84] **P.-L. Lions**, *Compactness in Boltzmann’s equation via Fourier integral operators and applications I*, 44
J. Math. Kyoto Univ. **34** (1994), 391–427. 45

- 1 [85] **P.-L. Lions**, *Compactness in Boltzmann's equation via Fourier integral operators and applications II*, 1
 2 *J. Math. Kyoto Univ.* **34** (1994), 429–461. 2
- 3 [86] **P.-L. Lions**, *Mathematical Topics in Fluid Mechanics, Vol. 1, Incompressible Models*, Clarendon Press/
 4 Oxford University Press, New York (1996). 4
- 5 [87] **P.-L. Lions and N. Masmoudi**, *From Boltzmann equation to the Navier–Stokes and Euler equations I*, *Arch.*
 6 *Ration. Mech. Anal.* **158** (2001), 173–193. 5
- 7 [88] **P.-L. Lions and N. Masmoudi**, *From Boltzmann equation to the Navier–Stokes and Euler equations II*,
 8 *Arch. Ration. Mech. Anal.* **158** (2001), 195–211. 6
- 9 [89] **T.-P. Liu**, *Solutions in the large for the equations of nonisentropic gas dynamics*, *Indiana Univ. Math. J.* **26**
 10 (1977), 147–177. 9
- 11 [90] **T.-P. Liu and S.-H. Yu**, *Boltzmann equation: Micro–macro decompositions and positivity of shock profiles*,
 12 *Comm. Math. Phys.* **246** (2004), 133–179. 10
- 13 [91] **A. Majda**, *Compressible Fluid Flow and Systems of Conservation Laws in Several Space Variables*,
 14 Springer-Verlag, New York (1984). 11
- 15 [92] **C. Marchioro and M. Pulvirenti**, *Mathematical Theory of Incompressible Nonviscous Fluids*, Springer-
 16 Verlag, New York (1994). 12
- 17 [93] **N. Maslova**, *The Kramers problem in the kinetic theory of gases*, *Zh. Vychisl. Mat. Mat. Fiz.* **22** (1982),
 18 700–709 (in Russian). 13
- 19 [94] **N. Maslova**, *Boltzmann equation*, *Dynamical Systems II*, Ya. Sinai, ed., Springer-Verlag, Berlin (1989). 14
- 20 [95] **N. Maslova**, *Nonlinear Evolution Equations. Kinetic Approach*, World Scientific, River Edge, NJ (1993). 15
- 21 [96] **N. Masmoudi and L. Saint-Raymond**, *From the Boltzmann equation to the Stokes–Fourier system in a*
 22 *bounded domain*, *Comm. Pure Appl. Math.* **56** (2003), 1263–1293. 16
- 23 [97] **S. Mischler**, *Kinetic equation with Maxwell boundary conditions*, Preprint. 17
- 24 [98] **C.B. Morrey**, *On the derivation of the equations of hydrodynamics from statistical mechanics*, *Comm.*
 25 *Pure Appl. Math.* **8** (1955), 279–326. 18
- 26 [99] **B. Nicolaenko**, *Shock wave solutions of the Boltzmann equation as a nonlinear bifurcation problem from*
 27 *the essential spectrum*, *Théories cinétiques classiques et relativistes*, Centre Natl. Recherche Sci., Paris
 28 (1975), 127–150. 19
- 29 [100] **B. Nicolaenko**, *A general class of nonlinear bifurcation problems from a point in the essential spectrum.*
 30 *Application to shock wave solutions of kinetic equations*, *Applications of Bifurcation Theory*, Academic
 31 Press, New York (1977), 333–357. 20
- 32 [101] **B. Nicolaenko and J.K. Thurber**, *Weak shock and bifurcating solutions of the non-linear Boltzmann equa-*
 33 *tion*, *J. Mécanique* **14** (1975), 305–338. 21
- 34 [102] **T. Nishida**, *A note on a theorem of Nirenberg*, *J. Differential Geom.* **12** (1977), 629–633. 22
- 35 [103] **T. Nishida**, *Fluid dynamical limit of the nonlinear Boltzmann equation to the level of the compressible*
 36 *Euler equation*, *Comm. Math. Phys.* **61** (1978), 119–148. 23
- 37 [104] **S. Olla, S.R.S. Varadhan and H.-T. Yau**, *Hydrodynamical limit for a Hamiltonian system with weak noise*,
 38 *Comm. Math. Phys.* **155** (1993), 523–560. 24
- 39 [105] **B. Perthame**, *Introduction to the collision models in Boltzmann's theory*, *Modeling of Collisions*,
 40 P.-A. Raviart, ed., Masson, Paris (1997). 25
- 41 [106] **B. Perthame and P. Souganidis**, *A limiting case for velocity averaging*, *Ann. Sci. École Norm. Sup. (4)* **31**
 42 (1998), 591–598. 26
- 43 [107] **L. Saint-Raymond**, *Discrete time Navier–Stokes limit for the BGK Boltzmann equation*, *Comm. Partial*
 44 *Differential Equation* **27** (2002), 149–184. 27
- 45 [108] **L. Saint-Raymond**, *Du modèle BGK de l'équation de Boltzmann aux équations d'Euler des fluides incom-*
 46 *pressibles*, *Bull. Sci. Math.* **126** (2002), 493–506. 28
- [109] **L. Saint-Raymond**, *From the BGK model to the Navier–Stokes equations*, *Ann. Sci. École Norm. Sup. (4)*
36 (2003), 271–317. 29
- [110] **L. Saint-Raymond**, *Convergence of solutions to the Boltzmann equation in the incompressible Euler limit*,
Arch. Ration. Mech. Anal. **166** (2003), 47–80. 30
- [111] **L.A. Santalò**, *Sobre la distribución probable de corpúsculos en un cuerpo, deducida de la distribución en*
sus secciones y problemas analogos, *Rev. Un. Mat. Argentina* **9** (1943), 145–164. 31
- [112] **V. Scheffer**, *Hausdorff measure and the Navier–Stokes equations*, *Comm. Math. Phys.* **55** (1977), 97–112. 32

- 1 [113] **J. Serrin**, *On the interior regularity of weak solutions of the Navier–Stokes equations*, Arch. Ration. Mech. 1
2 Anal. **9** (1962), 187–195. 2
- 3 [114] **T. Sideris**, *Formation of Singularities in 3D compressible fluids*, Comm. Math. Phys. **101** (1985), 475–485. 3
- 4 [115] **Y. Sone**, *Kinetic Theory and Fluid Dynamics*, Birkhäuser, Boston (2002). 4
- 5 [116] **S. Ukai**, *On the existence of global solutions of a mixed problem for the nonlinear Boltzmann equation*, 5
6 Proc. Japan Acad. **50** (1974), 179–184. 5
- 7 [117] **S. Ukai and K. Asano**, *Steady solutions of the Boltzmann equation for a gas flow past an obstacle. I. Existence*, 6
7 Arch. Ration. Mech. Anal. **84** (1983), 249–291. 7
- 8 [118] **S. Ukai and K. Asano**, *The Euler limit and initial layer of the nonlinear Boltzmann equation*, Hokkaido 8
9 Math. J. **12** (1983), 311–332. 9
- 10 [119] **S. Ukai, T. Yang and S.-H. Yu**, *Nonlinear boundary layers of the Boltzmann equation. I. Existence*, Comm. 10
11 Math. Phys. **236** (2003), 373–393. 10
- 12 [120] **S.R.S. Varadhan**, *Entropy methods in hydrodynamic scaling*, Proc. Internat. Congress of Mathematicians, 11
12 Zürich, 1994, Vol. 1, Birkhäuser, Basel (1995), 196–208. 12
- 13 [121] **C. Villani**, *A review of mathematical topics in collisional kinetic theory*, Handbook of Mathematical Fluid 13
14 Dynamics, Vol. I, North-Holland, Amsterdam (2002), 71–305. 14
- 15 [122] **H.T. Yau**, *Relative entropy and hydrodynamics of Ginzburg–Landau models*, Lett. Math. Phys. **22** (1991), 15
16 63–80. 15
- 17 [123] **V.I. Yudovich**, *Non-stationary flow of an ideal incompressible liquid*, Zh. Vychisl. Mat. Mat. Fiz. **3** (1963), 16
18 1032–1066 (in Russian). 17
- 19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45