
RIEMANN-HILBERT CORRESPONDENCE FOR MIXED TWISTOR \mathcal{D} -MODULES

by

Teresa Monteiro Fernandes & Claude Sabbah

Abstract. We introduce the notion of regularity for a relative holonomic \mathcal{D} -module in the sense of [16]. We prove that the solution functor from the bounded derived category of regular relative holonomic modules to that of relative constructible complexes is essentially surjective by constructing a right quasi-inverse functor. When restricted to relative \mathcal{D} -modules underlying a regular mixed twistor \mathcal{D} -module, this functor satisfies the left quasi-inverse property.

Contents

Introduction.....	1
1. Some complementary results to [16].....	4
2. Regular holonomic $\mathcal{D}_{X \times S/S}$ -modules.....	9
3. Relative tempered cohomology functors.....	14
4. Proof of the main results.....	27
Appendix. Locally constant sheaves of $p^{-1}\mathcal{O}_S$ -modules.....	33
References.....	36

Introduction

Let X and S be complex manifolds and let p be the projection $X \times S \rightarrow S$. In [16], we have considered a restricted notion of holomorphic family parametrized by S of holonomic \mathcal{D}_X -modules. These are coherent modules on the sheaf $\mathcal{D}_{X \times S/S}$ of relative differential operators whose characteristic variety, in the product $(T^*X) \times S$, is contained in $\Lambda \times S$ for some Lagrangean conic closed subset Λ of T^*X . This notion is restrictive in the sense that Λ does not vary with respect to S . We have also introduced the derived category of sheaves of $p^{-1}\mathcal{O}_S$ -modules with \mathbb{C} -constructible cohomology, also called S - \mathbb{C} -constructible complexes, together with the corresponding notion of perversity, and we have proved that the de Rham functor DR and its dual, the solution functor Sol, on the bounded derived category of $\mathcal{D}_{X \times S/S}$ -modules with holonomic cohomology take values in the derived category of S - \mathbb{C} -constructible complexes.

2010 Mathematics Subject Classification. 14F10, 32C38, 32S40, 32S60, 35Nxx, 58J10.

Key words and phrases. Holonomic relative D -module, regularity, relative constructible sheaf, relative perverse sheaf, mixed twistor D -module.

The research of TMF was supported by Fundação para a Ciência e Tecnologia, PEst OE/MAT/UI0209/2011. The research of CS was supported by the grant ANR-13-IS01-0001-01 of the Agence nationale de la recherche.

Many properties in the relative setting can be obtained from those in the “absolute case” (i.e., when S is reduced to a point), by specializing the parameter and by considering analogous properties for the restricted objects by the functors $Li_{s_o}^*$ when s_o varies in S . As a consequence, *strictness*, that is, $p^{-1}\mathcal{O}_S$ -flatness (or absence of $p^{-1}\mathcal{O}_S$ -torsion if $\dim S = 1$), plays an important role at various places. On the other hand, for an S - \mathbb{C} -constructible perverse complex F , the dual S - \mathbb{C} -constructible complex $\mathbf{D}F$ need not be perverse, and the subcategory of S - \mathbb{C} -constructible complexes F such that F and $\mathbf{D}F$ are perverse is specially interesting. Both notions (strictness and perversity of F and $\mathbf{D}F$) are related.

Proposition 1. *Assume that F and $\mathbf{D}F$ are perverse. Let $(X)_\alpha$ be a stratification of X adapted to F . Then, for any open strata X_α , $\mathcal{H}^{-d_X}i_\alpha^{-1}F$ is a locally free $p^{-1}\mathcal{O}_S$ -module of finite rank ($d_X := \dim X$).*

*Conversely, let Y be an hypersurface of X and let F be a locally free $p^{-1}\mathcal{O}_S$ -module of finite rank on $(X \setminus Y) \times S$. Then $j_!F[d_X]$ and its dual $Rj_*F[d_X]$ are perverse ($j : (X \setminus Y) \times S \hookrightarrow X \times S$).*

Moreover, when F is the de Rham complex or the solution complex of a holonomic $\mathcal{D}_{X \times S/S}$ -module, we have the following improvement of [16, Th. 1.2].

Proposition 2. *Let \mathcal{M} belong to $D_{\text{hol}}^b(\mathcal{D}_{X \times S/S})$. Then the following conditions are equivalent.*

- (1) \mathcal{M} is concentrated in degree 0 and $\mathcal{H}^0(\mathcal{M})$ is strict.
- (2) $\mathbf{D}\mathcal{M}$ is concentrated in degree 0 and $\mathcal{H}^0(\mathbf{D}\mathcal{M})$ is strict.
- (3) $\text{Sol}(\mathcal{M})[d_X]$ and $\text{DR}(\mathcal{M})[d_X]$ are perverse ($d_X = \dim X$).

Going further, it is natural to define the subcategory of *regular* relative holonomic \mathcal{D} -modules by imposing the regularity condition to each $Li_{s_o}^*\mathcal{M}$ (cf. Section 2.1).

Our main objective in this article is to approach the problem of constructing a quasi-inverse functor to Sol restricted to the category of regular holonomic $\mathcal{D}_{X \times S/S}$ -modules. For this purpose, in analogy with the method of Kashiwara [6], we introduce the functor RH^S , from the derived category $D_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$ of S - \mathbb{C} -constructible complexes to the bounded derived category $D_{\text{rhol}}^b(\mathcal{D}_{X \times S/S})$ of $\mathcal{D}_{X \times S/S}$ -modules with regular holonomic modules. Roughly speaking, it is a relative version of the functor $T\mathcal{H}om(\cdot, \mathcal{O})$ of Kashiwara [6] using the language of ind-sheaves or sheaves on a subanalytic site ([11], [12], [17]). In the locally constant case it coincides with the construction due to Deligne [2].

However, contrary to the absolute case, the behaviour by pull-back is not always controlled, due to the lack of an existence theorem of a Bernstein-Sato polynomial, so it remains conjectural that the derived category $D_{\text{rhol}}^b(\mathcal{D}_{X \times S/S})$ is stable by inverse images. This constitutes a major obstacle to obtain an equivalence of categories as in the absolute case. Our first main result concerns essential surjectivity of $\text{Sol} : D_{\text{rhol}}^b(\mathcal{D}_{X \times S/S}) \mapsto D_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$, when S is a curve.

Theorem 3. *Assume that $\dim S = 1$ and let $F \in D_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$. Then $\text{RH}^S(F) \in D_{\text{rhol}}^b(\mathcal{D}_{X \times S/S})$ and we have a functorial isomorphism ${}^p\text{Sol}(\text{RH}^S(F)) \simeq F$ in $D_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$ (with ${}^p\text{Sol}\mathcal{M} := \text{Sol}\mathcal{M}[d_X]$).*

As a consequence, we obtain:

Corollary 4. *Assume that $\dim S = 1$ and let $F \in D_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$ be such that F and DF are perverse. Set $\mathcal{M} := \mathrm{RH}^S(F) \in D_{\mathrm{rhol}}^b(\mathcal{D}_{X \times S/S})$. Then \mathcal{M} is concentrated in degree 0 and $\mathcal{H}^0 \mathcal{M}$ is strict.*

The category $\mathrm{MTM}(X)$ of mixed twistor \mathcal{D} -modules on the complex manifold X , together with the corresponding functors (pullback, pushforward by a projective map) was introduced by T. Mochizuki in [14]. We shall restrict our interest to the category $\mathrm{MTM}^{\mathrm{reg}}(X)$ of *regular mixed twistor \mathcal{D} -modules*. We then set $S = \mathbb{C}^*$. Any object of $\mathrm{MTM}^{\mathrm{reg}}(X)$, when restricted to $X \times S$, consists of a pair of W -filtered $\mathcal{D}_{X \times S/S}$ -modules and a sesquilinear pairing between them taking values in the sheaf of distributions which are analytic with respect to S (cf. [19] and [13] for the case of pure regular twistor \mathcal{D} -modules). For the sake of simplicity, we shall say that \mathcal{M} *underlies a regular mixed twistor \mathcal{D} -module* if it is one of the $\mathcal{D}_{X \times S/S}$ -modules of the pair. This defines a subcategory of $\mathrm{Mod}_{\mathrm{rhol}}(\mathcal{D}_{X \times S/S})$ (morphisms are similarly induced by morphisms in $\mathrm{MTM}^{\mathrm{reg}}$), which is not full however, but is stable by relative inverse images and relative proper direct images [14]. These properties are essential to prove our main application.

Theorem 5. *Assume that \mathcal{M} underlies a regular mixed twistor \mathcal{D} -module. Then there exists a canonical isomorphism*

$$(*) \quad \mathcal{M} \simeq \mathrm{RH}^S(\mathrm{PSol}(\mathcal{M}))$$

which is functorial with respect to morphisms in $\mathrm{Mod}(\mathcal{D}_{X \times S/S})$ between objects \mathcal{M}, \mathcal{N} of $\mathrm{MTM}^{\mathrm{reg}}(X)$. Moreover, we have a natural isomorphism

$$(**) \quad \mathrm{Hom}_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{N}) \simeq \mathrm{Hom}_{p^{-1}\mathcal{O}_S}(\mathrm{PSol} \mathcal{N}, \mathrm{PSol} \mathcal{M}).$$

We do not know how to characterize the essential image of $\mathcal{D}_{X \times S/S}$ -modules underlying a regular mixed twistor \mathcal{D} -module by the functor PSol , although we know that, for such a module \mathcal{M} , $\mathrm{PSol}(\mathcal{M})$ and its dual $\mathrm{PDR}(\mathcal{M})$ are perverse in $D_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$.

This paper is organized as follows. In Section 1, we prove the complements to [16], that is, Propositions 1 and 2. We establish in Theorem 2.6 and Corollary 2.8 the generalization of Deligne's results on the extension of a relative holomorphic connexion on the complementary of a normal crossing divisor. The construction of RH^S , explained in Section 3, is based on the notion of relative tempered distributions and holomorphic functions introduced in [15], which form subanalytic sheaves in the relative subanalytic site. Another constraint (vanishing of cohomologies) bounds our construction to the case $\dim S = 1$, which is not inconvenient for the application since mixed twistor \mathcal{D} -modules satisfy this condition. In the locally free case, the above extension is also obtained using the functor RH^S as proved in Lemma 4.2. Moreover, in this case, it provides an equivalence of categories (Theorem 2.11). We obtain Theorem 3 as a consequence of Lemma 4.2, and Theorem 5 is proved in Section 4.2 by reducing to [6, Cor. 8.6]. In the appendix we collect various results which are essential for the remaining part of the paper.

Acknowledgements. This work has benefited from discussions with Andrea d'Agnolo, Masaki Kashiwara, Yves Laurent and Luca Prelli, whom we warmly thank. The proof of Lemma 2.12 was kindly provided to us by Daniel Barlet.

1. Some complementary results to [16]

1.1. Notation and preliminary results. Throughout this work X and S will denote complex manifolds and $p_X : X \times S \rightarrow S$ will denote the projection. We will set $d_X := \dim X$, $d_S := \dim S$, and for any complex space Z , we will set similarly $d_Z = \dim Z$. We will often write p instead of p_X when there is no risk of ambiguity. We say that a $p^{-1}\mathcal{O}_S$ -module is *strict* if it is $p^{-1}\mathcal{O}_S$ -flat. Given $s_o \in S$, we denote by $Li_{s_o}^*(\cdot)$ the derived functor on $D^b(p^{-1}\mathcal{O}_S)$ of

$$F \longrightarrow F \otimes_{p^{-1}\mathcal{O}_S} p^{-1}(\mathcal{O}_S/\mathfrak{m}_{s_o}),$$

where \mathfrak{m}_{s_o} denotes the maximal ideal of holomorphic functions on S vanishing at s_o . The following results are straightforward.

Lemma 1.1. *Let $N \in D^{\geq 0}(p^{-1}\mathcal{O}_S)$ and let $s_o \in S$. Then $Li_{s_o}^*(N) \in D^{\geq -d_S}(X)$. If moreover, for any k , $\mathcal{H}^k(N)$ is strict then $Li_{s_o}^*(N) \in D^{\geq 0}(X)$.*

Lemma 1.2. *For any locally closed subset Z of $X \times S$, for any $F \in D^b(p^{-1}\mathcal{O}_S)$ and for any $s_o \in S$, we have $R\Gamma_Z(Li_{s_o}^*(F)) \simeq Li_{s_o}^*(R\Gamma_Z(F))$.*

We shall also need the following result which is contained in the proof of [16, Prop. 2.2]:

Proposition 1.3. *Let F belong to $D^b(p^{-1}\mathcal{O}_S)$ and assume that for every $(x_o, s_o) \in X \times S$ and for every j , $\mathcal{H}^j(F)_{(x_o, s_o)}$ is finitely generated over \mathcal{O}_{S, s_o} . Assume that, for a given j , $\mathcal{H}^j(Li_{s_o}^*(F)) = 0$ for any $s_o \in S$. Then $\mathcal{H}^j(F) = 0$. In particular, if, for a given integer k and for every s_o , $Li_{s_o}^*(F) \in D^{\geq k}(X)$ (respectively $Li_{s_o}^*(F) \in D^{\leq k}(X)$), then $F \in D^{\geq k}(X \times S)$ (respectively $F \in D^{\leq k}(X \times S)$).*

1.2. S - \mathbb{C} -constructibility and perversity. We refer to the appendix for the notion of S -locally constant sheaf on $X \times S$. We have defined in [16] the categories of S - \mathbb{R} -constructible sheaves (resp. S - \mathbb{C} -constructible sheaves) and the corresponding derived categories $D_{\mathbb{R}\text{-c}}^b(p^{-1}\mathcal{O}_S)$ (resp. $D_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$). For an object of $D^b(p^{-1}\mathcal{O}_S)$, the condition that it is an object F of $D_{\mathbb{R}\text{-c}}^b(p^{-1}\mathcal{O}_S)$ is a local property on X , since it is characterized by the property that the microsupport of F is contained in $\Lambda \times (T^*S)$ for some closed \mathbb{R}_+^* -conic Lagrangean subanalytic subset Λ of T^*X . Similarly, the condition that it is an object of $D_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$ is local, since it consists in adding that the microsupport is \mathbb{C}^\times -conic (cf. [16, Prop. 2.5 & Def. 2.19]).

The category $D_{\mathbb{R}\text{-c}}^b(p^{-1}\mathcal{O}_S)$ is endowed with a natural t-structures, which however is not preserved by duality, as seen by considering a skyscraper $p^{-1}\mathcal{O}_S$ -module on $X \times S$. We refer to loc. cit. for details.

Lemma 1.4. *For a given object $F \in \text{PD}_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$, F and DF are perverse if, and only if, for all $s_o \in S$, $Li_{s_o}^*(F)$ is perverse regarded as an object of $D_{\mathbb{C}\text{-c}}^b(\mathbb{C}_X)$.*

Proof. If $F \in \text{PD}_{\mathbb{C}\text{-c}}^{\leq 0}(p^{-1}\mathcal{O}_S)$ then $Li_{s_o}^*(F) \in \text{PD}_{\mathbb{C}\text{-c}}^{\leq 0}(\mathbb{C}_X)$ and the converse holds by Proposition 1.3. The assertion then follows by [16, Prop. 2.28]. q.e.d.

Proof of Proposition 1. For the first statement we note that, according to the assumption and the definition of t-structure, $i_\alpha^{-1}F$ is concentrated in degree $-d_X$ and $i_\alpha^{-1}\mathcal{H}^{-d_X}F$ is a $p^{-1}\mathcal{O}_S$ -coherent module. On the other hand, according to Lemma 1.4,

for any $s_o \in S$, $Li_{s_o}^* F$ is perverse, hence it is concentrated in degrees $\geq -d_X$. Recall that a coherent \mathcal{O}_S -module F_{x_o} is locally free if and only if $Li_{s_o}^* F_{x_o}$ is concentrated in degree zero for every $s_o \in S$. It follows that $i_\alpha^{-1} \mathcal{H}^{-d_X} F$ is locally free.

Conversely, since $Li_{s_o}^*$ commutes with $j_!$, Lemma 1.4 implies that $j_! F[d_X]$ and its dual are perverse. On the other hand, we have

$$\mathbf{D}(j_! F[d_X]) \simeq Rj_* R\mathcal{H}om_{p^{-1}\mathcal{O}_S}(F, p^{-1}\mathcal{O}_S)[d_X].$$

Since F is locally free, $\mathbf{D}'(F) := R\mathcal{H}om_{p^{-1}\mathcal{O}_S}(F, p^{-1}\mathcal{O}_S)$ is concentrated in degree zero and $\mathcal{H}^0 \mathbf{D}'(F)$ is locally free. Thus the statement follows by biduality. \square e.d.

1.3. Coherent $\mathcal{D}_{X \times S/S}$ -modules. Let $i : Z \hookrightarrow X$ be the inclusion of a closed submanifold in X . The following adaptation of Kashiwara's result (cf. e.g. [7, §4.8]) is straightforward.

Theorem 1.5 (Kashiwara's equivalence). *The pushforward functor ${}_{\mathbf{D}}i_*$ induces an equivalence between the category of coherent $\mathcal{D}_{Z \times S/S}$ -modules and that of coherent $\mathcal{D}_{X \times S/S}$ -modules supported on $Z \times S$. A quasi-inverse functor is $\mathcal{H}^{-\text{codim } Z} {}_{\mathbf{D}}i^*$, and $\mathcal{H}^j {}_{\mathbf{D}}i^* = 0$ for $j \neq -\text{codim } Z$ on objects of the latter category.*

The behaviour of coherence by pushforward with respect to the parameter space is obtained in the following proposition. Let $\pi : S \rightarrow S'$ be a morphism of complex manifolds. Let \mathcal{M} be a coherent $\mathcal{D}_{X \times S/S}$ -module which is π -good, that is, by definition, such that for any point $(x, s') \in X \times S'$ there exists a neighbourhood $U \times V'$ of (x, s') such that $\mathcal{M}_{U \times \pi^{-1}(V')}$ has a good filtration $F_\bullet \mathcal{M}$. The proof of the following proposition is similar to that given in [7, §4.7].

Proposition 1.6. *Assume that π is proper and that \mathcal{M} is π -good. Then $R\pi_* \mathcal{M} \in \mathbf{D}_{\text{coh}}^b(\mathcal{D}_{X \times S'/S'})$. Moreover, if $\text{Char } \mathcal{M} \subset \Lambda \times S$ with $\Lambda \subset T^*X$, then for each $k \in \mathbb{N}$, $\text{Char } R^k \pi_* \mathcal{M} \subset \Lambda \times S'$. In particular, if moreover \mathcal{M} is holonomic, then $R\pi_* \mathcal{M} \in \mathbf{D}_{\text{hol}}^b(\mathcal{D}_{X \times S'/S'})$.*

Proposition 1.7 ([20, Prop. 5.10, Th. 5.15]). *Let $f : X \rightarrow Y$ be a proper morphism of complex manifolds. Then there exists a functorial isomorphism ${}_{\mathbf{D}}f_! \mathbf{D}(\mathcal{M}) \rightarrow \mathbf{D}({}_{\mathbf{D}}f_! \mathcal{M})$ in $\mathbf{D}^b(\mathcal{D}_{X \times S/S})^{\text{op}}$, compatible with the composition with respect to f .*

As a consequence, using the projection formula for sheaves and replacing \mathcal{M} by $\mathbf{D}'(\mathcal{M})$ we recover the relative version of [7, Th. 4.33].

Corollary 1.8 (Adjunction formula). *Let $f : X \rightarrow Y$ be a proper morphism of complex manifolds. For $\mathcal{M} \in \mathbf{D}^b(\mathcal{D}_{X \times S/S})$ and $\mathcal{N} \in \mathbf{D}^b(\mathcal{D}_{Y \times S/S})$, there exists a canonical morphism $Rf_* R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, {}_{\mathbf{D}}f^* \mathcal{N})[d_X] \rightarrow R\mathcal{H}om_{\mathcal{D}_{Y \times S/S}}({}_{\mathbf{D}}f_! \mathcal{M}, \mathcal{N})[d_Y]$ which is an isomorphism.*

1.4. Holonomic $\mathcal{D}_{X \times S/S}$ -modules. The notion of holonomic $\mathcal{D}_{X \times S/S}$ -module has been recalled in the introduction. We refer to [16] for details on some of their properties. For such a $\mathcal{D}_{X \times S/S}$ -module, we will set ${}^{\text{p}}\text{DR } \mathcal{M} := \text{DR } \mathcal{M}[d_X]$ and ${}^{\text{p}}\text{Sol } \mathcal{M} = \text{Sol } \mathcal{M}[d_X]$.

Proposition 1.9. *Let \mathcal{M} be a holonomic $\mathcal{D}_{X \times S/S}$ -module.*

- (1) *Assume that $Li_{s_o}^* \mathcal{M} = 0$ for each $s_o \in S$. Then $\mathcal{M} = 0$.*

(2) Let \mathcal{I}_{s_o} be an ideal sheaf contained in the maximal ideal sheaf \mathfrak{m}_{s_o} consisting of germs of functions vanishing at s_o . Assume that $\mathcal{I}_{s_o}\mathcal{M}_{x^o, s_o} = \mathcal{M}_{x^o, s_o}$. Then $\mathcal{M}_{x^o, s_o} = 0$.

Proof. Assume $\mathcal{M} \neq 0$ and let $\Lambda \subset T^*X$ be a closed conic complex Lagrangian variety such that $\text{Char } \mathcal{M} \subset \Lambda \times S$. Let $(X_\alpha)_\alpha$ be a μ -stratification of X compatible with Λ . We will argue by induction on $\max_\beta \dim X_\beta$, where X_β runs among the strata included in the projection of Λ in X .

Assume first that some stratum X_β is open in X and let $x^o \in X_\beta$, so that $\text{Char } \mathcal{M} \subset (T_X^*X) \times S$ in the neighbourhood of x^o . It follows that \mathcal{M} is $\mathcal{O}_{X \times S}$ -coherent in the neighbourhood of $x^o \times S$, and Assumption 1.9(1) resp. (2) implies, according to Nakayama, that $\mathcal{M}|_{x^o \times S} = 0$ resp. $\mathcal{M}_{x^o, s_o} = 0$.

We are thus reduced to the case where no stratum X_β is open. Choose then a maximal stratum X_β . By applying Kashiwara's equivalence 1.5, which commutes with the \mathcal{O}_S -action, in the neighbourhood of any point of X_β we are reduced to the previous case. By induction on the dimension of the maximal strata, we conclude that Λ can be chosen empty, hence $\mathcal{M} = 0$. q.e.d.

Corollary 1.10. *Let \mathcal{M} be an object of $D_{\text{hol}}^b(\mathcal{D}_{X \times S/S})$. Assume $Li_{s_o}^* \mathcal{M} = 0$ for each $s_o \in S$. Then $\mathcal{M} = 0$.*

Proof. We first prove that if $Li_\Sigma^* \mathcal{M} = 0$ for every codimension-one germ of submanifold Σ , then $\mathcal{M} = 0$. Let σ be a local equation of Σ . Then the assumption is that the cone $C_\Sigma(\mathcal{M})$ of $\sigma : \mathcal{M} \rightarrow \mathcal{M}$ is isomorphic to zero. From the long exact sequence

$$\cdots \longrightarrow \mathcal{H}^j \mathcal{M} \xrightarrow{\sigma} \mathcal{H}^j \mathcal{M} \longrightarrow \mathcal{H}^j C_\Sigma(\mathcal{M}) = 0 \longrightarrow \cdots$$

we conclude as in Proposition 1.9 that $\mathcal{H}^j \mathcal{M} = 0$. We argue now by induction on d_S . In the case $d_S = 1$, every point has codimension one, so there is nothing more to prove. In general, for every germ of hypersurface Σ and every $s_o \in \Sigma$, we have $Li_{s_o}^* Li_\Sigma^* \mathcal{M} \simeq Li_{s_o}^* \mathcal{M} = 0$, so $Li_\Sigma^* \mathcal{M} = 0$ by induction, and the first part of the proof gives the desired assertion. q.e.d.

Corollary 1.11. *Let \mathcal{M} be an object of $D_{\text{hol}}^b(\mathcal{D}_{X \times S/S})$. Assume that $\mathcal{H}^j Li_{s_o}^* \mathcal{M} = 0$ for all $j \neq 0$ and all $s_o \in S$. Then $\mathcal{H}^j \mathcal{M} = 0$ for all $j \neq 0$ and $Li_{s_o}^* \mathcal{H}^0 \mathcal{M} = \mathcal{H}^0 Li_{s_o}^* \mathcal{M}$ for all $s_o \in S$.*

Proof. We prove it by induction on d_S . Let s be part of a local coordinate system centered at some $s_o \in S$ and denote by $i_{S'} : S' = \{s = 0\} \hookrightarrow S$ the inclusion. Then $Li_{S'}^* \mathcal{M}$ is an object of $D_{\text{hol}}^b(\mathcal{D}_{X \times S'/S'})$ and by induction it is concentrated in degree zero. Considering the long exact sequence

$$\cdots \mathcal{H}^j \mathcal{M} \xrightarrow{s} \mathcal{H}^j \mathcal{M} \longrightarrow \mathcal{H}^j Li_{S'}^* \mathcal{M} \longrightarrow \cdots$$

one obtains that $s : \mathcal{H}^j \mathcal{M} \rightarrow \mathcal{H}^j \mathcal{M}$ is onto for $j \neq 0$. According Proposition 1.9(2), we have $\mathcal{H}^j \mathcal{M} = 0$ for $j \neq 0$. The remaining statement is clear. q.e.d.

Corollary 1.12. *Let \mathcal{M} be a strict holonomic $\mathcal{D}_{X \times S/S}$ -module. Then $\mathcal{H}^j \mathbf{D} \mathcal{M} = 0$ for $j \neq 0$ and $\mathcal{H}^0 \mathbf{D} \mathcal{M}$ is a strict holonomic $\mathcal{D}_{X \times S/S}$ -module.*

Proof. By the strictness property, $\mathcal{H}^j Li_{s_o}^* \mathcal{M} = 0$ for $j \neq 0$, hence $Li_{s_o}^* \mathcal{M} = \mathcal{H}^0 Li_{s_o}^* \mathcal{M}$ is a holonomic \mathcal{D}_X -module, so $\mathcal{H}^j DLi_{s_o}^* \mathcal{M} = 0$ for $j \neq 0$ and $\mathcal{H}^0 DLi_{s_o}^* \mathcal{M}$ is \mathcal{D}_X -holonomic. Recall also (cf. [16, Prop. 3.1]) that $Li_{s_o}^* \mathcal{D}\mathcal{M} \simeq DLi_{s_o}^* \mathcal{M}$ for any $s_o \in S$. Therefore, $\mathcal{H}^j Li_{s_o}^* \mathcal{D}\mathcal{M} = 0$ for $j \neq 0$. According to [20, Prop. 2.5], $\mathcal{D}\mathcal{M}$ has holonomic cohomology. As a consequence, according to Corollary 1.11, $\mathcal{H}^j \mathcal{D}\mathcal{M} = 0$ for $j \neq 0$, $\mathcal{H}^0 \mathcal{D}\mathcal{M}$ is $\mathcal{D}_{X \times S/S}$ -holonomic, and $Li_{s_o}^* \mathcal{H}^0 \mathcal{D}\mathcal{M}$ has cohomology in degree zero at most for any $s_o \in S$, since

$$Li_{s_o}^* \mathcal{H}^0 \mathcal{D}\mathcal{M} \simeq \mathcal{H}^0 Li_{s_o}^* \mathcal{D}\mathcal{M} \simeq \mathcal{H}^0 DLi_{s_o}^* \mathcal{M}$$

and $Li_{s_o}^* \mathcal{M}$ is a holonomic \mathcal{D}_X -module. The conclusion follows from Lemma 1.13 below. q.e.d.

Lemma 1.13. *Let \mathcal{M} be a coherent $\mathcal{D}_{X \times S/S}$ -module. Then \mathcal{M} is strict if and only if $\mathcal{H}^j Li_{s_o}^* \mathcal{M} = 0$ for each $s_o \in S$ and each $j \neq 0$.*

Proof. The “only if” part is clear. The “if” part is clear if $d_S = 1$, since strictness is then equivalent to the absence of \mathcal{O}_S -torsion. In general, assume \mathcal{M} is not strict. Recall (cf. [18, Cor. A.0.2]) that there exists then a morphism $\pi : S_1 \rightarrow S$ from a smooth curve S_1 to S such that $\pi^* \mathcal{M}$ has \mathcal{O}_{S_1} -torsion. Let s_1 be a local coordinate on S_1 such that $\mathcal{K} := \ker[s_1 : \pi^* \mathcal{M} \rightarrow \pi^* \mathcal{M}] \neq 0$. Let i denote the composition $\{s_1 = 0\} \hookrightarrow S^1 \hookrightarrow S$. The exact sequence

$$\dots \longrightarrow \mathcal{H}^{-1} Li^* \mathcal{M} \longrightarrow \mathcal{H}^0 L\pi^* \mathcal{M} \xrightarrow{s_1} \mathcal{H}^0 L\pi^* \mathcal{M} \longrightarrow \mathcal{H}^0 Li^* \mathcal{M} \longrightarrow 0$$

show that $\mathcal{H}^{-1} Li^* \mathcal{M}$ surjects onto \mathcal{K} , hence is nonzero. q.e.d.

Corollary 1.14. *Let \mathcal{M} be a strict holonomic $\mathcal{D}_{X \times S/S}$ -module and set $F = {}^p\text{D}\mathcal{R}\mathcal{M}$ or ${}^p\text{Sol}\mathcal{M}$. Then F and $\mathcal{D}F$ are S -perverse.*

Proof. According to Corollary 1.12, this follows from Th. 1.2, Prop. 3.10, Th. 3.11 and Prop. 2.28 in [16]. q.e.d.

Proof of Proposition 2.

(1) \Leftrightarrow (2) follows from Corollary 1.12.

(1) \Rightarrow (3) follows from Corollary 1.14.

(1) \Leftarrow (3): According to Lemma 1.4, for any $s_o \in S$, $Li_{s_o}^* {}^p\text{Sol}(\mathcal{M})$ is perverse, therefore, for any $s_o \in S$, $Li_{s_o}^*(\mathcal{M})$ is concentrated in degree 0 and $\mathcal{H}^0 Li_{s_o}^*(\mathcal{M})$ is holonomic. The result then follows by Corollary 1.11 and Lemma 1.13. q.e.d.

We now can make precise the behaviour of the functors ${}^p\text{D}\mathcal{R}$ and ${}^p\text{Sol}$. According to Lemma 1.1, given $\mathcal{M} \in \mathcal{D}^{\geq 0}(\mathcal{D}_{X \times S/S})$ and $s_o \in S$, we have $Li_{s_o}^*(\mathcal{M}) \in \mathcal{D}^{\geq -d_S}(\mathcal{D}_X)$. If moreover, $\mathcal{H}^k(\mathcal{M})$ is $p^{-1}\mathcal{O}_S$ -flat for every k , then $Li_{s_o}^*(\mathcal{M}) \in \mathcal{D}^{\geq 0}(\mathcal{D}_X)$.

Proposition 1.15. *The functor ${}^p\text{D}\mathcal{R}$ has the following behaviour when considering the standard t -structure on $\mathcal{D}_{\text{hol}}^b(\mathcal{D}_{X \times S/S})$ and the t -structure given on $\mathcal{D}_{\text{c-c}}^b(p^{-1}\mathcal{O}_S)$:*

(1) Let $\mathcal{M} \in \mathcal{D}_{\text{hol}}^{\leq 0}(\mathcal{D}_{X \times S/S})$. Then ${}^p\text{D}\mathcal{R}\mathcal{M} \in {}^p\mathcal{D}_{\text{c-c}}^{\leq 0}(p^{-1}\mathcal{O}_S)$.

(2) Let $\mathcal{M} \in \mathcal{D}_{\text{hol}}^{\geq 0}(\mathcal{D}_{X \times S/S})$. Then ${}^p\text{D}\mathcal{R}\mathcal{M} \in {}^p\mathcal{D}_{\text{c-c}}^{\geq -d_S}(p^{-1}\mathcal{O}_S)$. Moreover, if for any k , $\mathcal{H}^k(\mathcal{M})$ is $p^{-1}\mathcal{O}_S$ -flat then ${}^p\text{D}\mathcal{R}\mathcal{M} \in {}^p\mathcal{D}_{\text{c-c}}^{\geq 0}(p^{-1}\mathcal{O}_S)$.

Proof.

(1) The assumption entails that, for any $s_o \in S$, $Li_{s_o}^*(\mathcal{M}) \in \mathbf{D}_{\text{hol}}^{\leq 0}(\mathcal{D}_X)$. Therefore, $\text{pDR}(Li_{s_o}^*(\mathcal{M})) \simeq Li_{s_o}^*(\text{pDR } \mathcal{M}) \in \text{pD}_{\mathbb{C}\text{-c}}^{\leq 0}(X)$. As a consequence,

$$Li_{s_o}^*(\text{pDR } \mathcal{M}|_{X_\alpha \times S}) \simeq \text{pDR}(Li_{s_o}^*(\mathcal{M}))|_{X_\alpha} \in \mathbf{D}^{\leq -\dim X_\alpha}(X_\alpha).$$

The statement then follows by Proposition 1.3.

(2) We have to prove that $R\Gamma_{X_\alpha \times S}(\text{pDR } \mathcal{M}) \in \mathbf{D}^{\geq -\dim X_\alpha - d_S}(X \times S)$. By Lemma 1.2, we have, for any $s_o \in S$,

$$Li_{s_o}^* R\Gamma_{X_\alpha \times S}(\text{pDR } \mathcal{M}) \simeq R\Gamma_{X_\alpha \times \{s\}}(Li_{s_o}^*(\text{pDR } \mathcal{M}))$$

On the other hand $Li_{s_o}^*(\mathcal{M}) \in \mathbf{D}_{\text{hol}}^{\geq -d_S}(\mathcal{D}_X)$ so $\text{pDR } Li_{s_o}^*(\mathcal{M}) \in \text{pD}_{\mathbb{C}\text{-c}}^{\geq -d_S}(X)$. Therefore

$$R\Gamma_{X_\alpha}(Li_{s_o}^*(F)) \in \mathbf{D}^{\geq -\dim X_\alpha - d_S}(X)$$

and the statement follows again by Proposition 1.3. The same argument implies the second part of the statement since, by Lemma 1.1, when $\mathcal{H}^k(\mathcal{M})$ is strict, for each k , $Li_{s_o}^*(\mathcal{M}) \in \mathbf{D}_{\text{hol}}^{\geq 0}(\mathcal{D}_X)$ for any s . q.e.d.

By the same arguments of Proposition 1.15, we obtain:

Proposition 1.16. *The functor pSol satisfies the following:*

- (1) Let $\mathcal{M} \in \mathbf{D}_{\text{hol}}^{\leq 0}(\mathcal{D}_{X \times S/S})$. Then $\text{pSol } \mathcal{M} \in \text{pD}_{\mathbb{C}\text{-c}}^{\geq 0}(p^{-1}\mathcal{O}_S)$.
- (2) Let $\mathcal{M} \in \mathbf{D}_{\text{hol}}^{\geq 0}(\mathcal{D}_{X \times S/S})$. Then $\text{pSol } \mathcal{M} \in \text{pD}_{\mathbb{C}\text{-c}}^{\leq d_S}(p^{-1}\mathcal{O}_S)$. Moreover, if for any k , $\mathcal{H}^k(\mathcal{M})$ is strict then $\text{pSol } \mathcal{M} \in \text{pD}_{\mathbb{C}\text{-c}}^{\leq 0}(p^{-1}\mathcal{O}_S)$.

Simple examples show that the final statement in Proposition 1.15 does not hold in general in the non strict case.

Theorem 1.17 (Proper pushforward). *Let $f : X \rightarrow Y$ be a proper morphism of complex manifolds and let \mathcal{M} belong to $\mathbf{D}_{\text{hol}}^b(\mathcal{D}_{X \times S/S})$ and has f -good cohomology. Then*

- (a) the pushforward ${}_D f_* \mathcal{M} := Rf_*(\mathcal{D}_{Y \leftarrow X/S} \otimes_{\mathcal{D}_{X \times S/S}}^L \mathcal{M})$ belongs to $\mathbf{D}_{\text{hol}}^b(\mathcal{D}_{Y \times S/S})$,
- (b) the de Rham complex satisfies $\text{pDR } {}_D f_* \mathcal{M} \simeq Rf_* \text{pDR } \mathcal{M}$ functorially in \mathcal{M} ,
- (c) the solution complex satisfies $\text{pSol } {}_D f_* \mathcal{M} \simeq Rf_* \text{pSol } \mathcal{M}$ functorially in \mathcal{M} .

Proof.

(a) The coherence and the holonomicity of the cohomology groups of ${}_D f_* \mathcal{M}$ follow from [20, Th. 4.2 and Cor. 4.3].

(b) The proof for \mathcal{D}_X -modules applies in a straightforward way and does not use holonomicity nor coherence of \mathcal{M} and neither properness of f (cf. e.g. [4, Th. 4.2.5]).

(c) We have

$$\text{pSol } {}_D f_* \mathcal{M} \stackrel{(1)}{\simeq} \text{pDR } {}_D f_* \mathcal{M} \stackrel{(2)}{\simeq} \text{pDR } {}_D f_* \mathbf{D} \mathcal{M} \stackrel{(3)}{\simeq} Rf_* \text{pDR } \mathbf{D} \mathcal{M} \stackrel{(4)}{\simeq} Rf_* \text{pSol } \mathcal{M},$$

where (1) and (4) follow from [16, (5)], (2) is given by [20, Th. 5.15] and (3) follows from (b) above. q.e.d.

2. Regular holonomic $\mathcal{D}_{X \times S/S}$ -modules

2.1. Regularity

Definition 2.1 (Regular holonomic $\mathcal{D}_{X \times S/S}$ -module). Let \mathcal{M} be a holonomic $\mathcal{D}_{X \times S/S}$ -module. We say that \mathcal{M} is regular holonomic if, for any $s_o \in S$, $Li_{s_o}^* \mathcal{M}$ belongs to $D_{\text{rhol}}^b(\mathcal{D}_X)$.

Note that, given an exact sequence in $\text{Mod}_{\text{hol}}(\mathcal{D}_{X \times S/S})$,

$$0 \longrightarrow \mathcal{N} \longrightarrow \mathcal{M} \longrightarrow \mathcal{L} \longrightarrow 0$$

if two of its terms are regular holonomic, then the third one is also regular holonomic.

As usual $D_{\text{rhol}}^b(\mathcal{D}_{X \times S/S})$ denotes the bounded derived category of complexes with regular holonomic cohomology groups. We have

Proposition 2.2. *Given a distinguished triangle in $D_{\text{rhol}}^b(\mathcal{D}_{X \times S/S})$,*

$$\mathcal{N} \longrightarrow \mathcal{M} \longrightarrow \mathcal{L} \xrightarrow{+1}$$

if two of its terms belong to $D_{\text{rhol}}^b(\mathcal{D}_{X \times S/S})$ the same holds for the third.

Proposition 2.3. *Assume that \mathcal{M} is an object of $D_{\text{rhol}}^b(\mathcal{D}_{X \times S/S})$. Then so is $\mathbf{D}\mathcal{M}$.*

Proof. Since the functors \mathbf{D} and $Li_{s_o}^*$ commute, the result follows by the definition and the fact that $D_{\text{rhol}}^b(\mathcal{D}_X)$ is stable by duality. q.e.d.

Corollary 2.4 (of Theorem 1.17). *Let $f : X \rightarrow Y$ be a proper morphism of complex manifolds. Let \mathcal{M} belong to $D_{\text{rhol}}^b(\mathcal{D}_{X \times S/S})$. Then $Rf_*(\mathcal{D}_{Y \leftarrow X/S} \otimes_{\mathcal{D}_{X \times S/S}}^L \mathcal{M})$ belongs to $D_{\text{rhol}}^b(\mathcal{D}_{Y \times S/S})$.*

Proof. The regularity follows from the commutativity of $Li_{s_o}^*$ with Rf_* and \otimes , for any $s_o \in S$. q.e.d.

2.2. Deligne extension of an S -locally constant sheaf. Let D be a normal crossing divisor in X and let $j : X^* := X \setminus D \hookrightarrow X$ denote the inclusion. Let F be a coherent S -locally constant sheaf on $X^* \times S$ and let $(E, \nabla) = (\mathcal{O}_{X^* \times S} \otimes_{p^{-1}\mathcal{O}_S} F, d_X)$ be the associated coherent $\mathcal{O}_{X^* \times S}$ -module with flat relative connection (cf. Remark A.10). In particular, E is naturally endowed with the structure of a left $\mathcal{D}_{X^* \times S/S}$ -module and j_*E with that of a $\mathcal{D}_{X \times S/S}$ -module. There exists a coherent \mathcal{O}_S -module G such that, locally on $X^* \times S$, $F \simeq p^{-1}G$ (cf. Proposition A.2). More precisely, let U be any contractible open set of X^* ; then $F|_{U \times S} \simeq p_U^{-1}G$ (cf. Proposition A.12).

Let $\varpi : \tilde{X} \rightarrow X$ denote the real blowing up of X along the components of D . Denote by $\tilde{j} : X^* \hookrightarrow \tilde{X}$ the inclusion, so that $j = \varpi \circ \tilde{j}$. Let $x^o \in D$, $\tilde{x}^o \in \varpi^{-1}(x^o)$ and let $s_o \in S$. Choose local coordinates $(x_1, \dots, x_\ell, x'_{\ell+1}, \dots, x'_n)$ at x^o such that $D = \{x_1 \cdots x_\ell = 0\}$ and consider the associated polar coordinates $(\rho, \theta, \mathbf{x}') := (\rho_1, \theta_1, \dots, \rho_\ell, \theta_\ell, x'_{\ell+1}, \dots, x'_n)$ so that \tilde{x}^o has coordinates $\rho^o = 0$, θ^o , $\mathbf{x}'^o = 0$. For $\varepsilon > 0$, we set

$$\tilde{U}_\varepsilon := \{\|\rho\| < \varepsilon, \|\mathbf{x}'\| < \varepsilon, \|\theta - \theta^o\| < \varepsilon\}, \quad \tilde{U}_\varepsilon^* := \tilde{U}_\varepsilon \setminus \{\rho_1 \cdots \rho_\ell = 0\}.$$

On the other hand, for $s_o \in S$, we denote by V some open neighbourhood of s_o in S . Note that since \tilde{U}_ε^* is contractible, we have $F|_{\tilde{U}_\varepsilon^* \times S} \simeq p_{\tilde{U}_\varepsilon^*}^{-1}G$ locally on S (cf. Proposition A.12), and thus $(E, \nabla)|_{\tilde{U}_\varepsilon^* \times S} \simeq (\mathcal{O}_{\tilde{U}_\varepsilon^* \times S} \otimes_{p^{-1}\mathcal{O}_S} p^{-1}G, d_X)$.

Definition 2.5 (Moderate growth).

(a) A germ of section $\tilde{v} \in (\tilde{j}_*E)_{(\tilde{x}^o, s_o)}$ is said to have *moderate growth* if for some (or any) system of generator of G_{s_o} , some $\varepsilon > 0$ and some V so that \tilde{v} is defined on $\tilde{U}_\varepsilon \times V$, its coefficients on the chosen generators of $1 \otimes G_{s_o}$ (these are sections of $\mathcal{O}(\tilde{U}_\varepsilon^* \times V)$ by means of the isomorphism above) are bounded by $C\rho^{-N}$, for some $C, N > 0$.

(b) A germ of section $v \in (j_*E)_{(x^o, s_o)}$ is said to have *moderate growth* if for each \tilde{x}^o in $\varpi^{-1}(x^o)$, the corresponding germ in $(\tilde{j}_*E)_{(\tilde{x}^o, s_o)}$ has moderate growth.

Theorem 2.6. *The subsheaf \tilde{E} of j_*E consisting of local sections having moderate growth is stable by ∇ and is $\mathcal{O}_{X \times S}(*D)$ -coherent.*

Proof. The problem is local on $X \times S$. We thus assume that $X \times S$ is a small neighbourhood of (x^o, s_o) as above. In such a neighbourhood, giving the local system is equivalent to giving $T_1, \dots, T_\ell \in \text{Aut}(G)$ which pairwise commute, according to Proposition A.9. According to [21, Cor. 2.3.10 & (3.45)], in the neighbourhood of s_o there exists for each k a logarithm of T_k , hence there exists $A_k \in \text{End}(G)$ such that $T_k = \exp(-2\pi i A_k)$, and the formula (2.11) of [21] can be used to show that there exist $A_1, \dots, A_\ell \in \text{End}(G)$ which pairwise commute. Set $\hat{E}_1 := \mathcal{O}_{X \times S}(*D) \otimes_{p^{-1}\mathcal{O}_S} G$, equipped with the connection ∇ such that $\nabla_{x_k \partial_{x_k}} : \hat{E}_1 \rightarrow \hat{E}_1$ is given by A_k if $k = 1, \dots, \ell$, and zero otherwise. Set $E_1 = \hat{E}_1|_{X^* \times S}$. Then the monodromy representation of ∇ on E_1 is given by T_1, \dots, T_ℓ , from which one deduces an isomorphism $(E, \nabla) \simeq (E_1, \nabla)$, according to Proposition A.9 and Remark A.10. It is then enough to show that $\tilde{E}_1 = \hat{E}_1$, where the former is as in the statement of the theorem, since we clearly have $\tilde{E} \simeq \tilde{E}_1$.

Let us fix local generators $\mathbf{g} := (g_i)$ of G and let us still denote by A_k a matrix of the endomorphism A_k with respect to (g_i) . Any local section v of E_1 can be expressed as $v = (1 \otimes \mathbf{g}) \cdot f$ for some vector f of local holomorphic functions. Let us set $x^A := x_1^{A_1} \cdots x_\ell^{A_\ell}$. A family of ∇ -horizontal generators is then given by $(1 \otimes \mathbf{g}) \cdot x^{-A}$, showing that the generators $1 \otimes g_i$ of \hat{E}_1 have moderate growth, hence are local sections of \tilde{E}_1 . Therefore, sections of j_*E_1 have moderate growth if and only if their coefficients over the generators $1 \otimes g_i$ have moderate growth. Since these coefficients are sections of $\mathcal{O}_{X^* \times S}$, they must be meromorphic. Hence $\tilde{E}_1 = \hat{E}_1$. q.e.d.

Remark 2.7. The functors j_* resp. \tilde{j}_* are exact functors from the category of coherent $\mathcal{O}_{X^* \times S}$ -modules with integrable relative connection to that of $\mathcal{O}_{X \times S}$ -resp. $\tilde{j}_* \mathcal{O}_{X^* \times S}$ -modules with integrable relative connection, since any point of D resp. $\varpi^{-1}(D)$ has a fundamental system of neighborhoods whose intersection with $X \setminus D$ resp. $\tilde{X} \setminus \varpi^{-1}(D)$ is Stein. Similarly, the correspondence $E \mapsto \tilde{E}$ is an exact functor from the category of coherent $\mathcal{O}_{X^* \times S}$ -modules with integrable relative connection to that of $\varpi^{-1} \mathcal{O}_{X \times S}$ -modules with integrable relative connection. Indeed, given a morphism $\varphi : (E, \nabla) \rightarrow (E', \nabla)$, it is clear that the morphism $\tilde{j}_* \varphi$

sends \tilde{E} to \tilde{E}' . The only point to check is right exactness, that is, that the induced morphism $\tilde{\varphi}$ is onto as soon as φ is onto. Keeping the notation of the proof of Theorem 2.6, we can start with E_1, E'_1 . The surjectivity of φ is equivalent to that of the morphism between the corresponding relative local systems, and restricting to x_o , to the induced morphism $G \rightarrow G'$. As a consequence, the morphism $\hat{\varphi}$ is onto, and therefore so is $\tilde{\varphi}$, according to the identification proved in the theorem.

Corollary 2.8. *Assume moreover that $d_S = 1$. Then \tilde{E} is $\mathcal{D}_{X \times S/S}$ -holonomic and regular with characteristic variety contained in $\Lambda \times S$, where Λ is the union of the conormal spaces of the natural stratification of (X, D) . Moreover, if F is $p^{-1}\mathcal{O}_S$ -locally free, then \tilde{E} is strict.*

Proof. We keep the notation of the proof of Theorem 2.6. Since the statement is local on $X \times S$, we can work with E_1 . We fix $s_o \in S$ and we take a local coordinate s centered at s_o .

Step one: assume G is \mathcal{O}_S -locally free. In this case, we are reduced to proving that \hat{E}_1 is strict holonomic with characteristic variety contained in $\Lambda \times S$, since its restriction to any s_o is a \mathcal{D}_X -module of Deligne type, hence is regular holonomic. Note that the strictness of \hat{E}_1 is obvious. We regard G_{s_o} as an $\mathcal{O}_{S, s_o}[A_1, \dots, A_\ell]$ -module which is \mathcal{O}_{S, s_o} -free. Let $\rho : (S', s'_o) \rightarrow (S, s_o)$ be the finite ramification of order N . Then G_{s_o} is identified with the invariant part of the pull-back $G'_{s'_o}$ of G_{s_o} by the Galois group \mathbb{Z}^N . Similarly, \hat{E}_1 is identified with the invariant part of $\rho_*\rho^*\hat{E}_1$ by the Galois group. The assertions of the corollary hold then for \hat{E}_1 if they hold for $\hat{E}'_1 := \rho^*\hat{E}_1$, according to Proposition 1.6.

Lemma 2.9. *There exists a finite ramification $\rho : (S', s'_o) \rightarrow (S, s_o)$ such that the pull-back $G'_{s'_o}$ of G_{s_o} has a finite filtration by $\mathcal{O}_{S', s'_o}[A_1, \dots, A_\ell]$ -submodules whose successive quotients have rank one over \mathcal{O}_{S', s'_o} .*

By the lemma and the previous remarks, we are reduced to the case where G has rank one over \mathcal{O}_S , so $A_k(s)$ is a holomorphic function $\lambda_k(s)$ and $\hat{E}_1 = \mathcal{O}_{X \times S}(*D)$ endowed with the $\mathcal{D}_{X \times S/S}$ -action defined by $\partial_{x_k} \cdot 1 = \lambda_k(s)/x_k$ if $k = 1, \dots, \ell$, and $\partial_{x_k} \cdot 1 = 0$ otherwise. This is the external product over \mathcal{O}_S of the $\mathcal{D}_{\mathbb{C} \times S/S}$ -modules $\mathcal{O}_{\mathbb{C} \times S}(*0)$ endowed with the connection $x_k\partial_{x_k} - \lambda_k(s)$ ($k = 1, \dots, \ell$) or with the connection ∂_{x_k} ($k > \ell$). It is therefore enough to show the corollary in the case $d_X = 1$. The latter case being obvious, we are reduced to proving the $\mathcal{D}_{\mathbb{C} \times S/S}$ -coherence of $(\mathcal{O}_{\mathbb{C} \times S}(*0), x\partial_x - \lambda(s))$ and to showing that the characteristic variety is contained in $(T_{\mathbb{C}}^*\mathbb{C}) \times S \cup (T_0^*\mathbb{C}) \times S$.

Up to isomorphism, we can assume that, if $\lambda(0) \in \mathbb{N}$, then $\lambda(0) = 0$. We then denote by μ its order of vanishing, i.e., $\lambda(s) = s^\mu u(s)$ with $u(0) \neq 0$. Then one has the relation

$$\partial_x^m \cdot \frac{1}{x} = \frac{\prod_{j=1}^m (\lambda(s) - j)}{x^{m+1}},$$

in which the numerator is thus invertible in \mathcal{O}_{S, s_o} . It follows that $1/x$ is a $\mathcal{D}_{\mathbb{C} \times S/S}$ -generator of this module. We thus have a surjective morphism

$$\mathcal{D}_{\mathbb{C} \times S/S} / \mathcal{D}_{\mathbb{C} \times S/S}(x\partial_x - (\lambda(s) - 1)) \longrightarrow (\mathcal{O}_{\mathbb{C} \times S}(*0), x\partial_x - \lambda(s))$$

sending the class of 1 to $1/x$. It obviously becomes an isomorphism after tensoring with $\mathcal{O}_{\mathbb{C} \times S}(*0)$. Since the left-hand side has no \mathcal{O}_S -torsion (cf. [16, Ex. 3.12]), we conclude that the above morphism is an isomorphism. Now, the assertions of the corollary are clear for $\mathcal{D}_{\mathbb{C} \times S/S} / \mathcal{D}_{\mathbb{C} \times S/S}(x\partial_x - (\lambda(s) - 1))$.

Step two. We now relax the assumption of local freeness on G . If $\mathcal{O}_{S, s_0}^N \rightarrow G_{s_0}$ is onto, then the kernel is torsion free, hence \mathcal{O}_{S, s_0} -free, and we have an exact sequence

$$0 \rightarrow G'' \rightarrow G' \rightarrow G \rightarrow 0$$

where G', G'' are \mathcal{O}_{S, s_0} -free of finite rank. By flatness of $\mathcal{O}_{X \times S}(*D)$ over \mathcal{O}_S , we have an exact sequence

$$0 \rightarrow \widehat{E}_1'' \rightarrow \widehat{E}_1' \rightarrow \widehat{E}_1 \rightarrow 0,$$

from which we deduce, according to Step one, that \widehat{E}_1 is holonomic with characteristic variety contained in $\Lambda \times S$. Moreover, the cohomology of $Li_{s_0}^* \widehat{E}_1$ appears as the kernel and cokernel of the morphism $i_{s_0}^* \widehat{E}_1'' \rightarrow i_{s_0}^* \widehat{E}_1'$, hence is also regular holonomic. \square e.d.

Proof of the lemma. Let us first work with the $\mathcal{O}_{S, s_0}(*0)$ -vector space $G_{s_0}(*0) := \mathcal{O}_{S, s_0}(*0) \otimes_{\mathcal{O}_{S, s_0}} G_{s_0}$. There exists a finite ramification

$$\rho : (S', s'_0) \rightarrow (S, s_0)$$

such that each equation $\det(t\text{Id} - A_k(s')) = 0$ has all its solutions in the field $\mathcal{O}_{S', s'_0}(*0)$. These solutions, being algebraic over \mathcal{O}_{S', s'_0} , belong to this ring. We can then assume from the beginning that all eigenvalues belong to \mathcal{O}_{S, s_0} . Then $G_{s_0}(*0)$ decomposes as an $\mathcal{O}_{S, s_0}(*0)[A_1, \dots, A_\ell]$ -module with respect to the multi-eigenvalues $\lambda(s) = (\lambda_1(s), \dots, \lambda_\ell(s))$ as $G_{s_0}(*0) = \bigoplus_{\lambda} G_{s_0}(*0)_{\lambda}$, and $A_k(s) - \lambda_k(s)$ is nilpotent on $G_{s_0}(*0)_{\lambda}$. If we choose a total order on the set of λ 's, we can define a filtration

$$G_{s_0}(*0)_{\leq \lambda} := \bigoplus_{\lambda' \leq \lambda} G_{s_0}(*0)_{\lambda'}.$$

It induces a filtration $G_{s_0, \leq \lambda} := G_{s_0}(*0)_{\leq \lambda} \cap G_{s_0}$, where the intersection is taken in $G_{s_0}(*0)$, and every successive quotient $G_{s_0, \leq \lambda} / G_{s_0, < \lambda}$ is an $\mathcal{O}_{S, s_0}[A_1, \dots, A_\ell]$ -submodule of $G_{s_0}(*0)_{\lambda}$, hence is \mathcal{O}_{S, s_0} -locally free and $A_k(s) - \lambda_k(s)$ is nilpotent on it for every $k = 1, \dots, \ell$.

We can therefore assume from the beginning that every $A_k(s)$ is nilpotent on G_{s_0} . We now argue by induction on ℓ . Consider the kernel filtration $G_{s_0}(*0)_j := \ker A_1^j$. This is a filtration by $\mathcal{O}_{S, s_0}(*0)[A_1, \dots, A_\ell]$ -submodules and A_1 acts by zero on the quotient module $G_{s_0}(*0)_j / G_{s_0}(*0)_{j-1}$ for every j . As above, we can induce this filtration on G_{s_0} by setting $G_{s_0, j} := G_{s_0}(*0)_j \cap G_{s_0}$ and $G_{s_0, j} / G_{s_0, j-1}$ is contained in $G_{s_0}(*0)_j / G_{s_0}(*0)_{j-1}$, hence has no \mathcal{O}_{S, s_0} -torsion, i.e., is \mathcal{O}_{S, s_0} -free. By induction on ℓ , we find a filtration whose successive quotients are free \mathcal{O}_{S, s_0} -modules and on which every A_k acts by zero, so that every rank-one \mathcal{O}_{S, s_0} -submodule is also trivially an $\mathcal{O}_{S, s_0}[A_1, \dots, A_\ell]$ -submodule, and the lemma is proved. \square e.d.

Definition 2.10. A coherent $\mathcal{D}_{X \times S/S}$ -module \mathcal{L} is said of D-type with singularities along D if it satisfies the following conditions:

- (1) $\text{Char } \mathcal{L} \subset \pi^{-1}(D) \times S \cup T_X^* X \times S$,
- (2) \mathcal{L} is regular holonomic and strict,
- (3) $\mathcal{L} \simeq \mathcal{L}(*D)$.

Proposition 2.11. *The category of holonomic systems \mathcal{L} of D -type along D is equivalent to the category of locally free $p_{X \setminus D}^{-1} \mathcal{O}_S$ -modules of finite type F on $(X \setminus D) \times S$ under the correspondences $\mathcal{L} \mapsto F = \mathcal{H}^0 \text{DR } \mathcal{L}|_{(X \setminus D) \times S}$ and $F \mapsto \mathcal{L} = \tilde{E}$.*

Before entering the proof of Proposition 2.11 we need the following description of relative moderate growth.

Lemma 2.12. *Assume that F is a locally free $p_{X \setminus D}^{-1} \mathcal{O}_S$ -modules of finite type. Then a section v of $j_* (F \otimes_{p^{-1} \mathcal{O}_S} \mathcal{O}_{(X \setminus D) \times S})$ is a section of \tilde{E} if and only if, for each $s_o \in S$, $v(\cdot, s_o)$ has moderate growth as a section of the \mathbb{C} -local system $\tilde{E}|_{\{s=s_o\}}$ on $X \setminus D$. In particular, $i_{s_o}^* \tilde{E} = \widehat{i_{s_o}^* E}$.*

Proof (provided by Daniel Barlet).

Case (1). Let us assume that F is S -constant. We may assume that the rank of F is 1. The statement being local, we may take local coordinates in a neighbourhood of $(x_o, s_o) \in D \times S$ and assume that we are given a holomorphic function $v(x, s)$ in $(U \setminus D) \times V$, where U is an open ball centered in x_o and V is an open ball centered in s_o , such that, for any $s \in V$, $v(x, s)$ is meromorphic with poles along D . To prove that v is meromorphic with poles along $D \times V$ it is sufficient, by Hartogs, to assume D non singular, hence defined by a coordinate $t = 0$. Writing the Laurent's expansion of v

$$v(x, s) = \sum_{i \in \mathbb{Z}} v_i(x, s) t^i,$$

where the v_i are holomorphic in $U \times V$, we introduce, for $m \geq 0$, the increasing sequence of closed analytic sets

$$X_m = \{(x, s) \mid v_k(x, s) = 0, \forall k \leq -m\}.$$

By the assumption, $U \times V = \bigcup_m X_m$ hence there must exist m_0 such that $X_{m_0} = U \times V$, which proves the claim.

Case (2). Let us assume that $D = \{x_1 \cdots x_\ell = 0\}$. For a general S -local system F locally free of rank d , let G and

$$(T_i(s), A_i(s))_{i=1, \dots, \ell}, \quad s \in V$$

be given by Proposition A.9 and Theorem 2.6, such that $T_i(s) = \exp(-2i\pi A_i(s))$, $i = 1, \dots, \ell$. Let (v_1, \dots, v_d) be a section of $F \otimes_{p_{X \setminus D}^{-1} \mathcal{O}_S} \mathcal{O}_{(X \setminus D) \times S}$ defined in $U' \times V$, for an open convex subset U' of $U \setminus D$ (where we keep the notation of Case (1)). Then, according to Case (1),

$$x_1^{A_1(s)} \cdots x_\ell^{A_\ell(s)} \begin{pmatrix} v_1(x, s) \\ \vdots \\ v_d(x, s) \end{pmatrix} = \begin{pmatrix} u_1(x, s) \\ \vdots \\ u_d(x, s) \end{pmatrix},$$

where each u_i is a meromorphic function with poles along D . Since the action of the matrix $x_1^{A_1(s)} \cdots x_\ell^{A_\ell(s)}$ does not affect the growth along D the statement follows. q.e.d.

Proof of Proposition 2.11. According to Theorem 2.6, if F is a $p^{-1}\mathcal{O}_S$ -local system locally free on $(X \setminus D) \times S$, \tilde{E} is a coherent $\mathcal{D}_{X \times S/S}$ -module of D-type. Conversely, suppose that \mathcal{L} is a coherent $\mathcal{D}_{X \times S/S}$ -module of D-type along D . The strictness assumption entails that $F := \mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{O}_{X \times S}, \mathcal{L})|_{(X \setminus D) \times S}$ is locally free of finite type. Let $\psi : \mathcal{L} \rightarrow j_*(F \otimes_{p|_{(X \setminus D) \times S} \mathcal{O}_S} \mathcal{O}_{(X \setminus D) \times S})$ be the natural morphism of $\mathcal{D}_{X \times S/S}$ -modules. Since ψ is an isomorphism on $(X \setminus D) \times S$ and $\mathcal{L} = \mathcal{L}(*D)$, it is injective. By definition, for each section u of $\psi(\mathcal{L})$ and each fixed $s_o \in S$, $u(\cdot, s_o)$ is a section of a regular holonomic \mathcal{D}_X -module, hence has moderate growth in the sense of [8, p. 862]. According to Lemma 2.12, u is a local section of \tilde{E} , hence $\psi(\mathcal{L}) \subset \tilde{E}$. The quotient sheaf $\tilde{E}/\psi(\mathcal{L})$ also satisfies $\tilde{E}/\psi(\mathcal{L})(*D) = \tilde{E}/\psi(\mathcal{L})$, and is zero on $(X \setminus D) \times S$. Therefore $\psi(\mathcal{L}) = \tilde{E}$. q.e.d.

3. Relative tempered cohomology functors

We shall keep the notations of Section 1 but X and S may also be real analytic manifolds.

3.1. Relative subanalytic site. We recall below the main constructions and results contained in [15] and obtain complementary results to be used in the sequel. We refer to [11] as a foundational paper and to [12] for a detailed exposition on the general theory of sheaves on sites.

Let X and S be real analytic manifolds. On $X \times S$ it is natural to consider the family \mathcal{T} consisting of finite unions of open relatively compact subsets and the family \mathcal{T}' of finite unions of open relatively compact sets of the form $U \times V$ making $X \times S$ both a \mathcal{T} - and a \mathcal{T}' -space in the sense of [3] and [10]. The associated sites $(X \times S)_{\mathcal{T}}$ and $(X \times S)_{\mathcal{T}'}$ are nothing more than, respectively, $(X \times S)_{\text{sa}}$ and the product of sites $X_{\text{sa}} \times S_{\text{sa}}$.

We shall denote by ρ , without reference to $X \times S$ unless otherwise specified, the natural functor of sites $\rho : X \times S \rightarrow (X \times S)_{\text{sa}}$ associated to the inclusion $\text{Op}((X \times S)_{\text{sa}}) \subset \text{Op}(X \times S)$. Accordingly, we shall consider the associated functors ρ_* , ρ^{-1} , $\rho!$ introduced in [12] and studied in [17].

We shall also denote by $\rho' : X \times S \rightarrow (X \times S)_{\mathcal{T}'}$ the natural functor of sites. Following [3] we have functors ρ'_* and $\rho'_!$ from $\text{Mod}(\mathbb{C}_{X \times S})$ to $\text{Mod}(\mathbb{C}_{X_{\text{sa}} \times S_{\text{sa}}})$.

Note that W is a \mathcal{T}' -open subset or, equivalently, $W \in \text{Op}(X_{\text{sa}} \times S_{\text{sa}})$, if W is a locally finite union of relatively compact subanalytic open subsets W of the form $U \times V$, $U \in \text{Op}(X_{\text{sa}})$, $V \in \text{Op}(S_{\text{sa}})$. We denote by $\eta : (X \times S)_{\text{sa}} \rightarrow X_{\text{sa}} \times S_{\text{sa}}$ the natural functor of sites associated to the inclusion $\text{Op}(X_{\text{sa}} \times S_{\text{sa}}) \hookrightarrow \text{Op}((X \times S)_{\text{sa}})$.

Remark 3.1. As well-known consequences of the properties of \mathcal{T} -spaces (cf. [11], see also [17]) we recall:

- ρ'^{-1} and $\rho'_!$ are exact and commute with tensor products.
- If $f : X \rightarrow Y$ is a morphism, ρ'^{-1} commutes with f^{-1} and ρ'_* commutes with f_* .
- $\rho'^{-1} \circ \rho'_* = \rho'^{-1} \circ \rho'_! = \text{Id}$.

- Adjunctions:

$$\begin{aligned}\rho'_* \mathcal{H}om(\rho'^{-1}(\cdot), \cdot) &\simeq \mathcal{H}om(\cdot, \rho'_*(\cdot)), \\ \rho'^{-1} \mathcal{H}om(\rho'_!(\cdot), \cdot) &\simeq \mathcal{H}om(\cdot, \rho'^{-1}(\cdot)).\end{aligned}$$

- ρ'_* commutes with $\mathcal{H}om$ and $R\mathcal{H}om$.
- Let f be an analytic map $X \rightarrow Y$. Still denoting by f the morphism $f \times \text{Id}_S : X \times S \rightarrow Y \times S$ or the associated morphism of sites, $X_{\text{sa}} \times S_{\text{sa}} \rightarrow Y_{\text{sa}} \times S_{\text{sa}}$, according to [12, 17.5], we have

- a left exact functor of relative direct image

$$f_* : \text{Mod}(\mathbb{C}_{X_{\text{sa}} \times S_{\text{sa}}}) \longrightarrow \text{Mod}(\mathbb{C}_{Y_{\text{sa}} \times S_{\text{sa}}}),$$

- an exact functor of relative inverse image

$$f^{-1} : \text{Mod}(\mathbb{C}_{Y_{\text{sa}} \times S_{\text{sa}}}) \longrightarrow \text{Mod}(\mathbb{C}_{X_{\text{sa}} \times S_{\text{sa}}}),$$

and (f^{-1}, f_*) is a pair of adjoint functors.

- ρ'^{-1} commutes with f^{-1} and ρ'_* commutes with f_* .

For example, the fourth item follows from adjunction and from the second item:

$$\mathcal{H}om(\rho'_*(\cdot), \rho'_*(\cdot)) \simeq \rho'_* \mathcal{H}om(\rho'^{-1} \circ \rho'_*(\cdot), \cdot) \simeq \rho'_* \mathcal{H}om(\cdot, \cdot).$$

For the commutation with $R\mathcal{H}om$ one uses injective resolutions plus the property that ρ'_* transform injective objects into quasi-injective objects which are $\mathcal{H}om(\rho'_*(F), \cdot)$ -acyclic for any F .

If \mathcal{R} is a sheaf of rings on $X_{\text{sa}} \times S_{\text{sa}}$, these properties remain true in $\text{Mod}(\mathcal{R})$. According to [12, Th.18.1.6 & Prop.18.5.4], $\text{Mod}(\mathcal{R})$ is a Grothendieck category so it admits enough injectives and enough flat objects. Hence the derived functors appearing in the sequel are well-defined.

3.2. Complements on S-constructible sheaves. In the sequel we shall assume that $d_S = 1$. The following result shows that $\text{Mod}_{\mathbb{R}\text{-c}}(p^{-1}\mathcal{O}_S)$ is a ρ_* as well as a ρ'_* -acyclic category.

Proposition 3.2. *Let $F \in \text{Mod}_{\mathbb{R}\text{-c}}(p^{-1}\mathcal{O}_S)$. Then $\mathcal{H}^k \rho_*(F) = \mathcal{H}^j \rho'_*(F) = 0$ for $k > 0$. In particular ρ'_* is exact on $\text{Mod}_{\mathbb{R}\text{-c}}(p^{-1}\mathcal{O}_S)$.*

Proof. Let U and V be open subanalytic relatively compact sets respectively in X and in S . Since dimension of S is 1, we may assume that V is Stein. Similarly to the proof of [17, Lem.2.1.1], it is sufficient to prove that, for each $k \neq 0$, there exists a finite covering $\{U_j \times V_j\}_{j \in I}$, $U_j \times V_j \in \mathcal{F}'$, of $U \times V$, such that $H^k(U_j \times V_j, F) = 0$.

Let $X = \bigcup_{\alpha} X_{\alpha}$ be a Whitney stratification adapted to F . By [9, Prop.8.2.5 (Triangulation Theorem)], there exist a simplicial complex $\mathbf{K} = (K, \Delta)$ and a homeomorphism $i : |\mathbf{K}| \simeq X$ such that, for any simplex $\sigma \in \Delta$, there exists α such that $i(|\sigma|) \subset X_{\alpha}$ and $i(|\sigma|)$ is a subanalytic manifold of X . Moreover, we may assume that U is a finite union of the images by i of open subsets $U(\sigma)$ of $|\mathbf{K}|$, with $U(\sigma) = \bigcup_{\tau \in \Delta, \tau \supset \sigma} |\tau|$. We shall see that we may take for $U_j \times V_j$ the open sets $i(U(\sigma)) \times V$. Therefore, still denoting by i the homeomorphism $|\mathbf{K}| \times S \rightarrow X \times S$, it is enough to prove that for any $\sigma \in \Delta$ and any $x \in |\sigma|$, we have:

$$(i) \quad H^0(U(\sigma) \times V; i^*F) \simeq H^0(V, F_{\{x\} \times S}),$$

(ii) $H^k(U(\sigma) \times V; i^*F) = H^k(V, F|_{\{x\} \times S}) = 0$, for $j \neq 0$.

The proof of (i) and (ii) proceeds mimicking the proof of [9, Prop. 8.1.4], using Proposition A.7, the fact that the $F|_{\{x\} \times S}$ is \mathcal{O}_S -coherent and that V is Stein. q.e.d.

Remark 3.3. By construction the isomorphisms (i) commute with the restrictions to open subsets $V' \subset V$ in S .

Recall that, given an abelian category \mathcal{C} , $\mathbf{K}^b(\mathcal{C})$ denotes the category of complexes in \mathcal{C} having bounded cohomology, the morphisms being defined up to homotopy. For a locally closed set of $X \times S$, \mathbb{C}_Z denotes both the constant sheaf on Z and its extension by zero as a sheaf on $X \times S$ (cf. [9, Prop. 2.5.4]).

Proposition 3.4. *Let $F \in \text{Mod}_{\mathbb{R}\text{-c}}(p^{-1}\mathcal{O}_S)$. Then F is quasi-isomorphic to a complex*

$$0 \longrightarrow \bigoplus_{i_\alpha \in I_\alpha} p^{-1}\mathcal{O}_S \otimes \mathbb{C}_{U_{\alpha, i_\alpha} \times V_{\alpha, i_\alpha}} \longrightarrow \cdots \longrightarrow \bigoplus_{i_\beta \in I_\beta} p^{-1}\mathcal{O}_S \otimes \mathbb{C}_{U_{\beta, i_\beta} \times V_{\beta, i_\beta}} \longrightarrow 0,$$

where $\{U_{i_j, i_j}\}_{j, i_j}$ are locally finite families of relatively compact open subanalytic subsets of X and $\{V_{i_j, i_j}\}_{j, i_j}$ are locally finite families of relatively compact open subanalytic subsets of S .

Proof. We shall adapt the outline of the proof of [1, Prop. A.2]. Let $X = \bigcup_\alpha X_\alpha$ be a Whitney stratification adapted to F . We keep the notation of Proposition 3.2 and its proof when using [9, Prop. 8.2.5].

For each integer i , let $\Delta_i \subset \Delta$ denote the subset of simplices of dimension $\leq i$ and set $\mathbf{K}_i = (K, \Delta_i)$. We shall prove by induction on i that there exists a morphism $\phi_i : G_i \rightarrow F$ in $\mathbf{K}^b(p^{-1}\mathcal{O}_S)$ such that:

- (a) The G_i^k are finite direct sums of $p^{-1}\mathcal{O}_S \otimes \mathbb{C}_{h(U_\sigma) \times V_{i, \sigma}}$ for some $\sigma \in \Delta_i$ and subanalytic open set $V_{i, \sigma}$ of S ,
- (b) The family $(h(U(|\sigma|)) \times V_{i, \sigma})_{\sigma \in \Delta_i}$ is a locally finite covering of $h(|\mathbf{K}_i|) \times S$,
- (c) One has

$$\phi_i|_{|\mathbf{K}_0| \times S} : G_i|_{|\mathbf{K}_0| \times S} \longrightarrow F|_{|\mathbf{K}_0| \times S}$$

is a quasi-isomorphism.

Case $i = 0$. Let $x \in h(|\mathbf{K}_0|)$, i.e., $x = h(\sigma)$ for some $\sigma \in \mathbf{K}_0$ and let $s_\sigma \in S$. We have that $F|_{\{x\} \times S} \simeq G_{0, \sigma}$ for some $G_{0, \sigma} \in \mathbf{D}_{\text{coh}}^b(\mathcal{O}_S)$; we then choose a subanalytic open set $V_\sigma \subset S$ such that $s \in V_\sigma$ and that $G_{0, \sigma}|_{V_\sigma}$ admits a bounded locally free $\mathcal{O}_S|_{V_\sigma}$ resolution $R_{0, \sigma} \rightarrow G_{0, \sigma}|_{V_\sigma}$. Since $\dim S = 1$, we may assume that V_σ is Stein.

Clearly, the family $(h(U(\sigma)) \times V_\sigma)_{\sigma \in \Delta_0}$ is a locally finite covering of $h(|\mathbf{K}_0|) \times S$. By (i) of Proposition 3.2 we have isomorphisms of \mathbb{C} -vector spaces

$$\Gamma(h(U(\sigma)) \times V'; F) \simeq \Gamma(V'; F|_{\{x\} \times V_\sigma}).$$

In view of the freeness of $R_{\sigma, 0}$, of the fact that V' is Stein and of isomorphisms (i) and (ii) of Proposition 3.2, we conclude quasi-isomorphisms in $\mathbf{K}^b(\mathbb{C})$

(iii) $\Gamma(V'; R_{\sigma, 0}) \rightarrow \Gamma(h(U(\sigma)) \times V'; F)$,

which commute with restrictions. On the other hand we have isomorphisms

(iv) $\phi_{V'} : \Gamma(h(U(\sigma)) \times V'; F) \simeq \Gamma(V'; p_* \mathcal{H}om_{\mathbf{K}^b(\mathbb{C}_{X \times S})}(\mathbb{C}_{h(U(\sigma)) \times V_\sigma}, F))$,

which also commute with restrictions to open subsets V'' of V' .

Combining (iii) and (iv) we get a quasi-isomorphism in $\mathbf{K}(\text{Mod}(\mathcal{O}_{V_\sigma}))$

$$R_{\sigma,0} \longrightarrow p_* \mathcal{H}om_{\mathbf{K}^b(\mathbb{C}_{X \times S})|_{V_\sigma}}(\mathbb{C}_{h(U(\sigma)) \times V_\sigma}, F)|_{V_\sigma}.$$

By adjunction, we get a morphism in $\mathbf{K}^b(\text{Mod}(p^{-1}\mathcal{O}_{V_\sigma}))$:

$$p|_{V_\sigma}^{-1} R_{\sigma,0} \longrightarrow \mathcal{H}om_{\mathbf{K}^b(\mathbb{C}_{X \times S})}(\mathbb{C}_{h(U(\sigma)) \times V_\sigma}, F)|_{X \times V_\sigma}$$

which, by the functorial properties of $\mathcal{H}om$ and \otimes , induces a morphism

$$(3.5) \quad \phi_{\sigma,0} : p^{-1} R_{\sigma,0} \otimes \mathbb{C}_{h(U(\sigma)) \times V_\sigma} \longrightarrow F.$$

By construction $\phi_0 := \bigoplus_{\sigma \in \Lambda_0} \phi_{\sigma,0}$ gives the desired morphism.

General case. Let us assume that ϕ_i is constructed and let us consider the distinguished triangle in $\mathbf{K}^b(\text{Mod}(p^{-1}\mathcal{O}_S))$:

$$H_i \xrightarrow{v_i} G_i \xrightarrow{\phi_i} F \xrightarrow{+1},$$

where $H_i|_{h(|\mathbf{K}_i|) \times S}$ is quasi-isomorphic to 0. Therefore

$$\bigoplus_{\sigma \in \Delta_{i+1} \setminus \Delta_i} H_i|_{h(|\sigma|) \times S} \longrightarrow H_i|_{h(|\mathbf{K}_{i+1}|) \times S}$$

is a quasi-isomorphism.

Likewise the case $i = 0$, let us choose, for each $\sigma \in \Delta_{i+1} \setminus \Delta_i$ and each $s_o \in S$, an open subanalytic relatively compact open subset $V_{\sigma,i+1}$ in S containing s , a complex $R_{\sigma,i+1}$ of free $\mathcal{O}_{V_{\sigma,i+1}}$ -modules quasi-isomorphic to $F|_{\{x\} \times V_{\sigma,i+1}}$, for arbitrary $x \in |\sigma|$, and a morphism

$$R_{\sigma,i+1} \longrightarrow p_* \mathcal{H}om(\mathbb{C}_{h(U(\sigma)) \times V_{\sigma,i+1}}, H_i)|_{V_{\sigma,i+1}}.$$

The family obtained as union of $(h(U_\sigma) \times V_{\sigma,i})_{\sigma \in \Delta_i}$ and $(h(U_\sigma) \times V_{\sigma,i+1})_{\sigma \in \Delta_{i+1} \setminus \Delta_i}$ clearly satisfies (b) with respect to $h(|\mathbf{K}_{i+1}|) \times S$.

As above we deduce a morphism

$$\phi'_{i+1} : G'_{i+1} := \bigoplus_{\sigma \in \Delta_{i+1} \setminus \Delta_i} p^{-1} R_{\sigma,i+1} \otimes \mathbb{C}_{h(U(\sigma)) \times V_{\sigma,i+1}} \longrightarrow H_i$$

such that, for $(x, s) \in h(|\mathbf{K}_{i+1}| \setminus |\mathbf{K}_i|) \times S$, the $\phi'_{i+1}(x,s)$ are quasi-isomorphisms. For $(x, s) \in h(|\mathbf{K}_i|) \times S$, the condition on H_i entails that $\phi'_{i+1}(x,s)$ is trivially a quasi-isomorphism. Therefore, $\phi'_{i+1}|_{h(|\mathbf{K}_{i+1}|) \times S}$ is a quasi-isomorphism.

Let G_{i+1} and H_{i+1} be defined by the distinguished triangles

$$G'_{i+1} \xrightarrow{\phi'_{i+1}} H_i \longrightarrow H_{i+1} \xrightarrow{+1} \quad \text{and} \quad G'_{i+1} \xrightarrow{v_i \circ \phi'_{i+1}} G_i \longrightarrow G_{i+1} \xrightarrow{+1}.$$

By construction and the induction hypothesis, G_{i+1} satisfies (a). The octahedral axiom applied to the preceding triangles induces a morphism $\phi_{i+1} : G_{i+1} \rightarrow F$ and hence a distinguished triangle

$$H_{i+1} \longrightarrow G_{i+1} \xrightarrow{\phi_{i+1}} F \xrightarrow{+1}.$$

Since by its construction $H_{i+1}|_{h(|\mathbf{K}_{i+1}|) \times S}$ is quasi-isomorphic to zero, ϕ_{i+1} satisfies (c) as desired. q.e.d.

Let $q : X \times S \rightarrow X$ denote the projection on the first factor.

Corollary 3.6. *Let $F \in D_{\mathbb{R}\text{-c}}^b(p_X^{-1}\mathcal{O}_S)$, $F' \in D_{\mathbb{R}\text{-c}}^b(\mathbb{C}_X)$ and let $s_o \in S$. Then*

- (a) $\rho'_*F \otimes \rho'_*q^{-1}F' \simeq \rho'_*(F \otimes q^{-1}F')$.
- (b) *The natural morphism*

$$\rho'_*p^{-1}(\mathcal{O}_S/\mathfrak{m}_{s_o}) \otimes_{\rho'_*p^{-1}\mathcal{O}_S}^L \rho'_*F \longrightarrow \rho'_*(p^{-1}(\mathcal{O}_S/\mathfrak{m}_{s_o}) \otimes_{p^{-1}\mathcal{O}_S}^L F)$$

is an isomorphism.

Proof.

(a) According to Proposition 3.4, which provides a $p^{-1}(\mathcal{O}_S)$ -flat resolution of F , we may assume that $F = \mathbb{C}_{U \times V} \otimes p^{-1}\mathcal{O}_S$. Similarly, as proved in [6], we may assume $F' = \mathbb{C}_{U'}$ for some open relatively compact subset $U' \subset X$. Therefore $F \otimes q^{-1}F' = \mathbb{C}_{(U \cap U') \times S} \otimes p^{-1}(\mathcal{O}_S \otimes \mathbb{C}_{V'})$. So, on one hand, according to [15, Lem. 3.6(2)] we have

$$\rho'_*(F \otimes q^{-1}F') = \rho'_*\mathbb{C}_{(U \cap U') \times S} \otimes \rho'_*p^{-1}(\mathcal{O}_S \otimes \mathbb{C}_{V'})$$

On the other hand we have, for the same reason,

$$\begin{aligned} \rho'_*(F) \otimes \rho'_*(q^{-1}F') &= \rho'_*(\mathbb{C}_{U \times S} \otimes p^{-1}(\mathcal{O}_S \otimes \mathbb{C}_{V'})) \otimes \rho'_*(\mathbb{C}_{U' \times S}) \\ &= \rho'_*(\mathbb{C}_{U \times S}) \otimes \rho'_*p^{-1}(\mathcal{O}_S \otimes \mathbb{C}_{V'}) \otimes \rho'_*(\mathbb{C}_{U' \times S}) \end{aligned}$$

and the result follows from the equality $\rho'_*(\mathbb{C}_{(U \cap U') \times S}) = \rho'_*(\mathbb{C}_{U \times S}) \otimes \rho'_*(\mathbb{C}_{U' \times S})$ (cf. [3, Th. 2.2.6(2)]).

(b) According to Proposition 3.2, ρ'_* is exact on $D_{\mathbb{R}\text{-c}}^b(p_X^{-1}\mathcal{O}_S)$ and, as above, we may assume that $F = \mathbb{C}_{U \times V} \otimes p^{-1}\mathcal{O}_S$. Up to shrinking V (possible by the construction of the family $\{V_{i_j, i_j}\}_{j, i_j}$ mentioned in Proposition 3.4), we can also assume that there is a holomorphic coordinate s vanishing at s_o defined on V . It remains to observe that the left term in (b) is realized by the complex $\rho'_*F \xrightarrow{s-s_o} \rho'_*F$ and the right term by $\rho'_*(F \xrightarrow{s-s_o} F)$. They are thus isomorphic by the exactness of ρ'_* . q.e.d.

3.3. Relative subanalytic sheaves. Let G be a sheaf on $(X \times S)_{\mathcal{T}'}$. One defines the (separated) presheaf $\eta^{\leftarrow}G$ on $(X \times S)_{\text{sa}}$ by setting, for $W \in \text{Op}((X \times S)_{\text{sa}})$,

$$\eta^{\leftarrow}G(W) = \varinjlim_{W \subset W'} G(W')$$

with $W' \in \text{Op}((X \times S)_{\mathcal{T}'})$. Let $\eta^{-1}G$ be the associated sheaf.

Let F be a subanalytic sheaf on $(X \times S)_{\text{sa}}$. We shall denote by $F^{S, \#}$ the sheaf on $X_{\text{sa}} \times S_{\text{sa}}$ associated to the presheaf

$$\begin{aligned} \text{Op}(X_{\text{sa}} \times S_{\text{sa}}) &\longrightarrow \text{Mod}(\mathbb{C}) \\ U \times V &\longmapsto \Gamma(X \times V; \rho^{-1}\Gamma_{U \times S}F) \simeq \text{Hom}(\mathbb{C}_U \boxtimes \rho_!\mathbb{C}_V, F) \\ &\simeq \varprojlim_{\substack{W \in V \\ W \in \text{Op}^c(S_{\text{sa}})}} \Gamma(U \times W; F). \end{aligned}$$

With the notations above, for a morphism $f : X \rightarrow Y$ of analytic manifolds, we have

$$\begin{aligned} f^{-1}(F^{S, \#}) &\simeq ((f \times \text{Id}_S)^{-1}F)^{S, \#}, \\ f_*(F^{S, \#}) &\simeq ((f \times \text{Id}_S)_*F)^{S, \#}, \end{aligned}$$

for any $F \in \text{Mod}(\mathbb{K}_{(X \times S)_{\text{sa}}})$. We set

$$(3.7) \quad F^S := \eta^{-1}F^{S, \#}$$

and call it the relative sheaf associated to F . It is a sheaf on $(X \times S)_{\text{sa}}$ and $(\cdot)^S$ defines a left exact functor on $\text{Mod}(\mathbb{C}_{(X \times S)_{\text{sa}}})$. We will denote by $(\cdot)^{RS, \#}$ and $(\cdot)^{RS} \simeq \eta^{-1} \circ (\cdot)^{RS, \#}$ the associated right derived functors.

Recall that by [15, Lem. 3.4], we have an isomorphism $\text{Id} \simeq R\eta_* \eta^{-1}$.

Definition 3.8. We define $\mathfrak{D}\mathfrak{b}_{X \times S}^{t, S}$ as the relative sheaf associated to $\mathfrak{D}\mathfrak{b}_{X \times S}^t$.

By [15, Props. 5.1(2), 5.2(i) and 5.3(2)], $\mathfrak{D}\mathfrak{b}_Y^{t, S}$ has the following properties:

- (i) $\begin{cases} \Gamma(U \times V; \mathfrak{D}\mathfrak{b}_{X \times S}^{t, S}) = \Gamma(X \times V; \rho^{-1} \Gamma_{U \times S} \mathfrak{D}\mathfrak{b}_{X \times S}^t) \\ \simeq \Gamma(X \times V; T\mathcal{H}om(\mathbb{C}_{U \times S}, \mathfrak{D}\mathfrak{b}_{X \times S}^t)), \end{cases}$
- (ii) $\begin{cases} \rho^{-1} R\mathcal{H}om(G \boxtimes H, \mathfrak{D}\mathfrak{b}_{X \times S}^{t, S}) \simeq \rho^{-1} R\mathcal{H}om(G \boxtimes \rho_! H, \mathfrak{D}\mathfrak{b}_{X \times S}^{t, S}) \\ \simeq R\mathcal{H}om(\mathbb{C}_X \boxtimes H, T\mathcal{H}om(G \boxtimes \mathbb{C}_S, \mathfrak{D}\mathfrak{b}_{X \times S}^t)), \\ \text{for any } G \in \mathbb{D}_{\mathbb{R}\text{-c}}^b(X), H \in \mathbb{D}_{\mathbb{R}\text{-c}}^b(S). \end{cases}$

(iii) $\mathfrak{D}\mathfrak{b}_Y^{t, S}$ is $\Gamma(U \times V; \cdot)$ -acyclic for each $U \in \text{Op}(X_{\text{sa}})$, $V \in \text{Op}(S_{\text{sa}})$. In particular, $\mathfrak{D}\mathfrak{b}_{X \times S}^t$ is $(\cdot)^{S, \#}$ -acyclic and hence $(\cdot)^S$ -acyclic.

As a consequence,

$$(3.9) \quad \rho^{-1} \mathfrak{D}\mathfrak{b}_{X \times S}^{t, S} \simeq \rho'^{-1} \mathfrak{D}\mathfrak{b}_{X \times S}^{t, S, \#} \simeq \mathfrak{D}\mathfrak{b}_{X \times S}.$$

Let us now assume that X and S are complex analytic manifolds and denote as usual by $\overline{X} \times \overline{S}$ the complex conjugate manifold. By [15, Lem. 5.4 and Lem. 5.5], there is a natural action of $\eta^{-1} \rho_* p^{-1} \mathcal{O}_S$, $\rho_! \mathcal{D}_{X \times S}$ and of $\rho_! \mathcal{D}_{\overline{X} \times \overline{S}}$ on $\mathfrak{D}\mathfrak{b}_{X \times S}^{t, S}$ and the same argument holds for ρ' instead of ρ and $\mathfrak{D}\mathfrak{b}_{X \times S}^{t, S, \#}$ instead of $\mathfrak{D}\mathfrak{b}_{X \times S}^{t, S}$. We then define $\mathcal{O}_{X \times S}^{t, S}$ as the derived relative sheaf associated to $\mathcal{O}_{X \times S}^t$, that is

$$\mathcal{O}_{X \times S}^{t, S} \simeq (R\mathcal{H}om_{\rho_! \mathcal{D}_{\overline{X} \times \overline{S}}}(\rho_! \mathcal{O}_{\overline{X} \times \overline{S}}, \mathfrak{D}\mathfrak{b}_{X \times S}^t))^{RS}.$$

More precisely, setting $\mathcal{O}_{X \times S}^{t, S, \#} := (\mathcal{O}_{X \times S}^t)^{RS, \#}$, then $\mathcal{O}_{X \times S}^{t, S} \simeq \eta^{-1} \mathcal{O}_{X \times S}^{t, S, \#}$.

According to the $(\cdot)^{S, \#}$ and the $(\cdot)^S$ -acyclicity of $\mathfrak{D}\mathfrak{b}_{X \times S}^t$ (cf. [15]) we have:

$$\mathcal{O}_{X \times S}^{t, S} \simeq R\mathcal{H}om_{\rho_! \mathcal{D}_{\overline{X} \times \overline{S}}}(\rho_! \mathcal{O}_{\overline{X} \times \overline{S}}, \mathfrak{D}\mathfrak{b}_{X \times S}^{t, S}),$$

and

$$\mathcal{O}_{X \times S}^{t, S, \#} \simeq R\mathcal{H}om_{\rho_! \mathcal{D}_{\overline{X} \times \overline{S}}}(\rho_! \mathcal{O}_{\overline{X} \times \overline{S}}, \mathfrak{D}\mathfrak{b}_{X \times S}^{t, S, \#}).$$

Moreover, by Propositions 4.1 and 5.7 of loc. cit., for $G = \mathbb{C}_U$ and $H = \mathbb{C}_V$:

$$(3.10) \quad \begin{aligned} \rho'^{-1} R\mathcal{H}om(\mathbb{C}_{U \times V}, \mathcal{O}_{X \times S}^{t, S, \#}) &\simeq \rho^{-1} R\mathcal{H}om(\mathbb{C}_{U \times V}, \mathcal{O}_{X \times S}^{t, S}) \\ &\simeq \rho^{-1} R\mathcal{H}om(\mathbb{C}_U \boxtimes \rho_! \mathbb{C}_V, \mathcal{O}_{X \times S}^t) \\ &\simeq R\mathcal{H}om(\mathbb{C}_{X \times V}, T\mathcal{H}om(\mathbb{C}_{U \times S}, \mathcal{O}_{X \times S}^t)), \\ &\text{for any } U \in \text{Op}(X_{\text{sa}}) \text{ and } V \in \text{Op}(S_{\text{sa}}). \end{aligned}$$

Since $\mathcal{O}_{X \times S} \simeq \rho'^{-1}(\mathcal{O}_{X \times S}^{t, S, \#})$, by adjunction we get a morphism in $\mathbb{D}^b(\rho_! \mathcal{D}_{X \times S/S})$

$$(3.11) \quad \mathcal{O}_{X \times S}^{t, S, \#} \longrightarrow R\rho'_* \mathcal{O}_{X \times S}$$

Lemma 3.12. *The morphism (3.11) induces an isomorphism in $\mathbb{D}^b(\rho'_* p^{-1} \mathcal{O}_S)$:*

$$\begin{aligned} R\mathcal{H}om_{\rho_! \mathcal{D}_{X \times S/S}}(\rho_! \mathcal{O}_{X \times S}, \mathcal{O}_{X \times S}^{t, S, \#}) \\ \xrightarrow{\sim} R\mathcal{H}om_{\rho_! \mathcal{D}_{X \times S/S}}(\rho_! \mathcal{O}_{X \times S}, R\rho'_* \mathcal{O}_{X \times S}) \simeq \rho'_* p^{-1} \mathcal{O}_S. \end{aligned}$$

Proof. We start by proving the first isomorphism. Since the family of open subanalytic sets of the form $U \times V$ generate the open coverings of $X_{\text{sa}} \times S_{\text{sa}}$, it is sufficient to prove that, for any open subanalytic relatively compact sets U in X and V in S , morphism (3.11) induces an isomorphism, functorial in $U \times V$

$$\begin{aligned} R\Gamma(U \times V, R\mathcal{H}om_{\rho'_! \mathcal{D}_{X \times S/S}}(\rho'_! \mathcal{O}_{X \times S}, \mathcal{O}_{X \times S}^{t,S,\#})) \\ \longrightarrow R\Gamma(U \times V; R\mathcal{H}om_{\rho'_! \mathcal{D}_{X \times S/S}}(\rho'_! \mathcal{O}_{X \times S}, R\rho'_* \mathcal{O}_{X \times S})) \end{aligned}$$

We have a chain of isomorphisms

$$\begin{aligned} R\Gamma(U \times V, R\mathcal{H}om_{\rho'_! \mathcal{D}_{X \times S/S}}(\rho'_! \mathcal{O}_{X \times S}, \mathcal{O}_{X \times S}^{t,S,\#})) \\ \simeq \text{RHom}(\rho'_* \mathbb{C}_{U \times V}, R\mathcal{H}om_{\rho'_! \mathcal{D}_{X \times S/S}}(\rho'_! \mathcal{O}_{X \times S}, \mathcal{O}_{X \times S}^{t,S,\#})) \\ \simeq \text{RHom}_{\rho'_! \mathcal{D}_{X \times S/S}}(\rho'_! \mathcal{O}_{X \times S}, R\mathcal{H}om(\mathbb{C}_{U \times V}, \mathcal{O}_{X \times S}^{t,S,\#})) \\ \simeq \text{RHom}_{\mathcal{D}_{X \times S/S}}(\mathcal{O}_{X \times S}, \rho'^{-1} R\mathcal{H}om(\rho'_* \mathbb{C}_{U \times V}, \mathcal{O}_{X \times S}^{t,S,\#})) \quad (\text{by adjunction}) \\ \simeq \text{RHom}_{\mathcal{D}_{X \times S/S}}(\mathcal{O}_{X \times S}, R\mathcal{H}om(\mathbb{C}_{X \times V}, T\mathcal{H}om(\mathbb{C}_{U \times S}, \mathcal{O}_{X \times S}))) \\ \simeq \text{RHom}(\mathbb{C}_{X \times V}, R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{O}_{X \times S}, T\mathcal{H}om(\mathbb{C}_{U \times S}, \mathcal{O}_{X \times S}))), \quad (\text{by (3.10)}). \end{aligned}$$

Similarly we have the chain of isomorphisms:

$$\begin{aligned} R\Gamma(U \times V; R\mathcal{H}om_{\rho'_! \mathcal{D}_{X \times S/S}}(\rho'_! \mathcal{O}_{X \times S}, R\rho'_* \mathcal{O}_{X \times S})) \\ \simeq \text{RHom}(\rho'_* \mathbb{C}_{U \times V}, R\mathcal{H}om_{\rho'_! \mathcal{D}_{X \times S/S}}(\rho'_! \mathcal{O}_{X \times S}, R\rho'_* \mathcal{O}_{X \times S})) \\ \simeq \text{RHom}_{\rho'_! \mathcal{D}_{X \times S/S}}(\rho'_! \mathcal{O}_{X \times S}, R\mathcal{H}om(\rho'_* \mathbb{C}_{U \times V}, R\rho'_* \mathcal{O}_{X \times S})) \\ \simeq \text{RHom}_{\mathcal{D}_{X \times S/S}}(\mathcal{O}_{X \times S}, \rho'^{-1} R\mathcal{H}om(\rho'_* \mathbb{C}_{U \times V}, R\rho'_* \mathcal{O}_{X \times S})) \\ \simeq \text{RHom}_{\mathcal{D}_{X \times S/S}}(\mathcal{O}_{X \times S}, R\mathcal{H}om(\mathbb{C}_{X \times V}, R\mathcal{H}om(\mathbb{C}_{U \times S}, \mathcal{O}_{X \times S}))) \\ \simeq \text{RHom}(\mathbb{C}_{X \times V}, R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{O}_{X \times S}, R\mathcal{H}om(\mathbb{C}_{U \times S}, \mathcal{O}_{X \times S}))). \end{aligned}$$

The isomorphisms of each chain are compatible with (3.11) because they come from natural equivalences of functors. We have thus reduced the proof to showing that the morphism

$$(i) \quad R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{O}_{X \times S}, T\mathcal{H}om(\mathbb{C}_{U \times S}, \mathcal{O}_{X \times S})) \\ \longrightarrow R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{O}_{X \times S}, R\mathcal{H}om(\mathbb{C}_{U \times S}, \mathcal{O}_{X \times S}))$$

is an isomorphism in $\mathbf{D}^b(p^{-1}\mathcal{O}_S)$, functorial in U .

Note that $\mathcal{D}_{X \times S} \otimes_{\mathcal{D}_{X \times S/S}} \mathcal{O}_{X \times S}$ is nothing but the transfer module $\mathcal{D}_{X \times S \rightarrow S}$ associated to p . In local coordinates x in X , it is realized by

$$\mathcal{D}_{X \times S} / \mathcal{D}_{X \times S} \partial_{x_1} + \mathcal{D}_{X \times S} \partial_{x_2} + \cdots + \mathcal{D}_{X \times S} \partial_{x_n}.$$

Thus, for any subanalytic open set U in X , we have

$$\begin{aligned} R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{O}_{X \times S}, R\mathcal{H}om(\mathbb{C}_{U \times S}, \mathcal{O}_{X \times S})) \\ \simeq R\mathcal{H}om_{\mathcal{D}_{X \times S}}(\mathcal{D}_{X \times S \rightarrow S}, T\mathcal{H}om(\mathbb{C}_{U \times S}, \mathcal{O}_{X \times S})). \end{aligned}$$

Hence, the proof of (i) reduces to prove that the morphism

$$(ii) \quad R\mathcal{H}om_{\mathcal{D}_{X \times S}}(\mathcal{D}_{X \times S \rightarrow S}, T\mathcal{H}om(\mathbb{C}_{U \times S}, \mathcal{D}\mathbf{b}_{X \times S})) \\ \longrightarrow R\mathcal{H}om_{\mathcal{D}_{X \times S}}(\mathcal{D}_{X \times S \rightarrow S}, R\mathcal{H}om(\mathbb{C}_{U \times S}, \mathcal{D}\mathbf{b}_{X \times S})) \\ \simeq R\mathcal{H}om_{\mathcal{D}_{X \times S}}(\mathcal{D}_{X \times S \rightarrow S}, \Gamma_{U \times S}(\mathcal{D}\mathbf{b}_{X \times S}))$$

is an isomorphism, functorial in U . Since each derivation D_{x_i} defines an epimorphism on the submodule $\ker \partial_{x_1} \cap \cdots \cap \ker \partial_{x_{i-1}}$ of $\Gamma_{U \times S}(\mathfrak{D}_{X \times S}) / T\mathcal{H}om(\mathbb{C}_{X \times S}, \mathfrak{D}_{X \times S})$, in view of the local Koszul resolution of $\mathcal{D}_{X \times S \rightarrow S}$, we get that

$$R\mathcal{H}om_{\mathcal{D}_{X \times S}}(\mathcal{D}_{X \times S \rightarrow S}, \Gamma_{U \times S}(\mathfrak{D}_{X \times S}) / T\mathcal{H}om(\mathbb{C}_{X \times S}, \mathfrak{D}_{X \times S}))$$

is concentrated in degree 0 and its \mathcal{H}^0 is isomorphic to $\ker \partial_{x_1} \cap \cdots \cap \ker \partial_{x_n} = 0$ locally. Therefore, after applying the functor $R\mathcal{H}om_{\mathcal{D}_{\bar{X} \times \bar{S}}}(\mathcal{O}_{\bar{X} \times \bar{S}}, \cdot)$, we deduce a natural isomorphism

$$\begin{aligned} R\mathcal{H}om_{\mathcal{D}_{X \times S}}(\mathcal{D}_{X \times S \rightarrow S}, T\mathcal{H}om(\mathbb{C}_{U \times S}, \mathcal{O}_{X \times S})) \\ \simeq R\mathcal{H}om_{\mathcal{D}_{X \times S}}(\mathcal{D}_{X \times S \rightarrow S}, R\mathcal{H}om(\mathbb{C}_{U \times S}, \mathcal{O}_{X \times S})). \end{aligned}$$

The functoriality on U is obvious so this ends the proof of (ii) hence the first isomorphism is proved.

To show the second isomorphism, we have, for any U, V as above,

$$\begin{aligned} \mathrm{RHom}(\mathbb{C}_{X \times V}, R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{O}_{X \times S}, R\mathcal{H}om(\mathbb{C}_{U \times S}, \mathcal{O}_{X \times S}))) \\ \simeq \mathrm{RHom}(\mathbb{C}_{U \times V}, R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{O}_{X \times S}, \mathcal{O}_{X \times S})) \\ \simeq R\Gamma(U \times V; R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{O}_{X \times S}, \mathcal{O}_{X \times S})). \end{aligned}$$

Since

$$R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{O}_{X \times S}, \mathcal{O}_{X \times S}) \simeq p_X^{-1} \mathcal{O}_S,$$

the last expression is isomorphic to

$$R\Gamma(U \times V, p_X^{-1} \mathcal{O}_S) = R\Gamma(U \times V, \rho'_* p_X^{-1} \mathcal{O}_S),$$

as desired. q.e.d.

3.4. The functors TH^S and RH^S . Recall that $\rho'^{-1} \rho'_! = \mathrm{Id}$ so $\rho'^{-1} \rho'_! \mathcal{D}_{X \times S/S} = \mathcal{D}_{X \times S/S}$. On $\mathrm{D}_{\mathbb{R}\text{-c}}^b(p^{-1} \mathcal{O}_S)$ we define the (derived) functors:

- TH^S given by the assignment

$$F \longmapsto \mathrm{TH}^S(F) := \rho'^{-1} R\mathcal{H}om_{\rho'_* p^{-1} \mathcal{O}_S}(\rho'_* F, \mathfrak{D}_{X \times S}^{t, S, \sharp}),$$

- RH^S given by the assignment

$$F \longmapsto \mathrm{RH}^S(F) := \rho'^{-1} R\mathcal{H}om_{\rho'_* p^{-1} \mathcal{O}_S}(\rho'_* F, \mathcal{O}_{X \times S}^{t, S, \sharp})[d_X],$$

for $F \in \mathrm{D}_{\mathbb{R}\text{-c}}^b(p^{-1} \mathcal{O}_S)$.

Clearly $\rho^{-1} = \rho'^{-1} \eta_*$ but $\rho_* \neq \eta^{-1} \rho'_*$ hence, if in the preceding definitions we replace ρ' by ρ , we obtain a different notion. Note also that by the adjunction formula for ρ' we have

$$(3.13) \quad \mathrm{RH}^S(F) \simeq R\mathcal{H}om_{\mathcal{D}_{\bar{X} \times \bar{S}}}(\mathcal{O}_{\bar{X} \times \bar{S}}, \mathrm{TH}^S(F))[d_X].$$

We have a functorial isomorphism in $\mathrm{D}^b(\mathcal{D}_{X \times S/S})$:

$$R\mathcal{H}om_{p^{-1} \mathcal{O}_S}(F, \mathcal{O}_{X \times S}) \simeq \rho'^{-1} R\mathcal{H}om_{\rho'_* p^{-1} \mathcal{O}_S}(\rho'_* F, R\rho'_* \mathcal{O}_{X \times S})$$

Combining this isomorphism with (3.11) we obtain a functorial morphism

$$(3.14) \quad \mathrm{RH}^S(F)[-d_X] \longrightarrow R\mathcal{H}om_{p^{-1} \mathcal{O}_S}(F, \mathcal{O}_{X \times S}),$$

and therefore, for any object \mathcal{M} of $D^b(\mathcal{D}_{X \times S/S})$, a bi-functorial morphism

$$(3.15) \quad R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathrm{RH}^S(F)[-d_X]) \longrightarrow R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, R\mathcal{H}om_{p^{-1}\mathcal{O}_S}(F, \mathcal{O}_{X \times S})).$$

Lemma 3.16. *Let $F \in D_{\mathbb{R}\text{-c}}^b(p^{-1}\mathcal{O}_S)$. Then the natural morphism*

$$R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{O}_{X \times S}, \mathrm{RH}^S(F)[-d_X]) \longrightarrow R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{O}_{X \times S}, R\mathcal{H}om_{p^{-1}\mathcal{O}_S}(F, \mathcal{O}_{X \times S}))$$

is an isomorphism. In particular ${}^p\mathrm{DR}(\mathrm{RH}^S(F)) \simeq DF$.

Proof. We have a chain of functorial isomorphisms in $D^b(p^{-1}\mathcal{O}_S)$

$$\begin{aligned} R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{O}_{X \times S}, \mathrm{RH}^S(F)[-d_X]) &\simeq \rho'^{-1} R\mathcal{H}om_{\rho'_! \mathcal{D}_{X \times S/S}}(\rho'_! \mathcal{O}_{X \times S}, R\mathcal{H}om_{\rho'_* p^{-1}\mathcal{O}_S}(R\rho'_* F, \mathcal{O}_{X \times S}^{t, S, \#})) \\ &\simeq \rho'^{-1} R\mathcal{H}om_{\rho'_* p^{-1}\mathcal{O}_S}(R\rho'_* F, R\mathcal{H}om_{\rho'_! \mathcal{D}_{X \times S/S}}(\rho'_! \mathcal{O}_{X \times S}, \mathcal{O}_{X \times S}^{t, S, \#})). \end{aligned}$$

Similarly, the other side, we have a chain of functorial isomorphisms

$$\begin{aligned} R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{O}_{X \times S}, R\mathcal{H}om_{p^{-1}\mathcal{O}_S}(F, \mathcal{O}_{X \times S})) &\simeq \rho'^{-1} R\mathcal{H}om_{\rho'_! \mathcal{D}_{X \times S/S}}(\rho'_! \mathcal{O}_{X \times S}, R\mathcal{H}om_{\rho'_* p^{-1}\mathcal{O}_S}(R\rho'_* F, R\rho'_* \mathcal{O}_{X \times S})) \\ &\simeq \rho'^{-1} R\mathcal{H}om_{\rho'_* p^{-1}\mathcal{O}_S}(R\rho'_* F, R\mathcal{H}om_{\rho'_! \mathcal{D}_{X \times S/S}}(\rho'_! \mathcal{O}_{X \times S}, R\rho'_* \mathcal{O}_{X \times S})). \end{aligned}$$

The first part of the statement then follows by Lemma 3.12. Let us prove the last assertion. We have, functorially in F , a chain of isomorphisms

$$\begin{aligned} R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{O}_{X \times S}, R\mathcal{H}om_{p_X^{-1}\mathcal{O}_S}(F, \mathcal{O}_{X \times S})) &\simeq R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{O}_{X \times S} \otimes_{p^{-1}\mathcal{O}_S} F, R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{O}_{X \times S}, \mathcal{O}_{X \times S})) \\ &\simeq R\mathcal{H}om_{p_X^{-1}\mathcal{O}_S}(F, p^{-1}\mathcal{O}_S) \simeq D'(F). \quad \text{q.e.d.} \end{aligned}$$

Example 3.17. For a given hypersurface Y of X (possibly with singularities) let us denote by j the open inclusion $X \setminus Y \hookrightarrow X$ as well as the associated map $(X \setminus Y) \times S \hookrightarrow X \times S$.

(1) Assume $F \simeq p^{-1}\mathcal{O}_S^\ell|_{(X \setminus Y) \times S}$ for some $\ell \in \mathbb{N}$. Then $T\mathcal{H}om(\mathbb{C}_{(X \setminus Y) \times S}, \mathcal{O}_{X \times S})$ is a regular holonomic $\mathcal{D}_{X \times S}$ -module endowed with a natural structure of $\mathcal{D}_{X \times S/S}$ -module, as proved in [6], hence it is regular holonomic as a $\mathcal{D}_{X \times S/S}$ -module. More precisely, given locally an equation $f = 0$ defining Y , $T\mathcal{H}om(\mathbb{C}_{(X \setminus Y) \times S}, \mathcal{O}_{X \times S})$ is the localized of $\mathcal{O}_{X \times S}$ with respect to f . By Corollary 3.6(a) we have

$$\begin{aligned} \mathrm{RH}^S(j!F)[-d_X] &\simeq \rho'^{-1} R\mathcal{H}om_{\rho'_* p^{-1}\mathcal{O}_S}(\rho'_* p^{-1}\mathcal{O}_S^\ell \otimes \rho'_* \mathbb{C}_{(X \setminus Y) \times S}, \mathcal{O}_{X \times S}^{t, S, \#}) \\ &\simeq \rho'^{-1} R\mathcal{H}om_{\rho'_* p^{-1}\mathcal{O}_S}(\rho'_* p^{-1}\mathcal{O}_S^\ell, R\mathcal{H}om(\rho'_* \mathbb{C}_{(X \setminus Y) \times S}, \mathcal{O}_{X \times S}^{t, S, \#})) \\ &\simeq \rho^{-1} R\mathcal{H}om(\rho_* \mathbb{C}_{(X \setminus Y) \times S}, \mathcal{O}_{X \times S}^{t, S})^\ell, \end{aligned}$$

and by (3.10) we get

$$\mathrm{RH}^S(j!F)[-d_X] \simeq T\mathcal{H}om(\mathbb{C}_{(X \setminus Y) \times S}, \mathcal{O}_{X \times S})^\ell.$$

(2) Let us assume that $F \simeq p_X^{-1}G$ with G coherent over \mathcal{O}_S , that is, F is an S -constant local system on $(X \setminus Y) \times S$.

According to Case (1), by considering a local free resolution \mathcal{O}_S^\bullet of G on a sufficiently small open subset V of S , we obtain that $\mathrm{RH}^S(j_!F)[-d_X]|_{p^{-1}V}$ is quasi-isomorphic to a complex for which the terms are finite direct sums of $T\mathcal{H}om(\mathbb{C}_{(X \setminus Y) \times S}, \mathcal{O}_{X \times S})$ and the differentials are given by the right multiplication by matrices with entries in $p^{-1}\mathcal{O}_S$, hence $\mathcal{D}_{X \times S/S}$ -linear morphisms. Therefore the cohomology groups are regular holonomic $\mathcal{D}_{X \times S/S}$ -modules.

(3) We assume $X = \mathbb{C}$. For any $F \in \mathrm{Mod}_{\mathbb{C}\text{-c}}(p^{-1}\mathcal{O}_S)$ such that $(\mathbb{C}^*, \{0\})$ is an adapted stratification we have $F \otimes \mathbb{C}_{\{0\} \times S} \simeq p^{-1}G \otimes \mathbb{C}_{\{0\} \times S}$ for some coherent \mathcal{O}_S -module G . Then, by Corollary 3.6(a),

$$\begin{aligned} \mathrm{RH}^S(F) &\simeq R\mathcal{H}om_{p^{-1}\mathcal{O}_S}(p^{-1}G, T\mathcal{H}om(\mathbb{C}_{\{0\} \times S}, \mathcal{O}_{X \times S}))[1] \\ &\simeq R\mathcal{H}om_{p^{-1}\mathcal{O}_S}(p^{-1}G, B_{\{0\} \times S|X \times S}). \end{aligned}$$

3.5. Extension in the case of an open subanalytic set. Let U denote an open subanalytic subset of X , let $U_{X_{\mathrm{sa}}}$ be the subanalytic site induced by X_{sa} on U (cf. [17, Rem. 1.1.1]), and let $j : U_{X_{\mathrm{sa}}} \times S_{\mathrm{sa}} \rightarrow X_{\mathrm{sa}} \times S_{\mathrm{sa}}$ denote the open embedding of subanalytic sites. Recall that the authorized open coverings in $U_{X_{\mathrm{sa}}}$ are those obtained as intersections of coverings of X_{sa} with U . We keep the notation j for the morphism $j \times \mathrm{Id}_S : U \times S \rightarrow X \times S$ and ρ' for the morphism of sites $U \times S \rightarrow U_{X_{\mathrm{sa}}} \times S_{\mathrm{sa}}$. One easily checks from the notion of morphism of sites (see [12, Chap. 16] for details) that the items in Remark 3.1 still hold in this framework with $f = j$.

Lemma 3.18. *Let F be a $p^{-1}\mathcal{O}_S$ -coherent S -locally constant sheaf on $U \times S$. Then, for any $\rho'_*p_{U \times S}^{-1}\mathcal{O}_S$ -module \mathcal{L} , the natural morphism*

$$(3.18^*) \quad \rho'_* \mathbf{D}'F \otimes_{\rho'_*p^{-1}\mathcal{O}_S}^L \mathcal{L} \longrightarrow R\mathcal{H}om_{\rho'_*p^{-1}\mathcal{O}_S}(\rho'_*F, \mathcal{L})$$

is an isomorphism.

Proof. The construction of (3.18*) is similar to the case of usual sheaves:

(1) For any $\mathcal{F}, \mathcal{G}, \mathcal{H} \in \mathrm{Mod}(\rho'_*p^{-1}\mathcal{O}_S)$, we have a natural morphism

$$\mathcal{H}om_{\rho'_*p^{-1}\mathcal{O}_S}(\mathcal{F}, \mathcal{G}) \otimes_{\rho'_*p^{-1}\mathcal{O}_S} \mathcal{H} \longrightarrow \mathcal{H}om_{\rho'_*p^{-1}\mathcal{O}_S}(\mathcal{F}, \mathcal{G} \otimes_{\rho'_*p^{-1}\mathcal{O}_S} \mathcal{H}).$$

We deduce a natural morphism

$$(3.19) \quad R\mathcal{H}om_{\rho'_*p^{-1}\mathcal{O}_S}(\mathcal{F}, \mathcal{G}) \otimes_{\rho'_*p^{-1}\mathcal{O}_S}^L \mathcal{H} \longrightarrow R\mathcal{H}om_{\rho'_*p^{-1}\mathcal{O}_S}(\mathcal{F}, \mathcal{G} \otimes_{\rho'_*p^{-1}\mathcal{O}_S}^L \mathcal{H})$$

by considering a flat resolution of \mathcal{H} and an injective resolution of \mathcal{G} .

(2) We have $\mathbf{D}'F := R\mathcal{H}om_{p^{-1}\mathcal{O}_S}(F, p^{-1}\mathcal{O}_S)$. Recall that ρ'_* commutes with $\mathcal{H}om$ and $R\mathcal{H}om$ hence

$$\rho'_* \mathbf{D}'F \simeq R\mathcal{H}om_{\rho'_*p^{-1}\mathcal{O}_S}(\rho'_*F, \rho'_*p^{-1}\mathcal{O}_S).$$

Therefore, the desired morphism (3.18*) is obtained from (3.19) with

$$\mathcal{F} = \rho'_*F, \mathcal{G} = \rho'_*p^{-1}\mathcal{O}_S, \mathcal{H} = \mathcal{L}.$$

To prove that it is an isomorphism it is sufficient to consider a locally finite covering of $U \times S$ by open subsets of the form $U' \times V$, $U' \in \mathrm{Op}(U_{X_{\mathrm{sa}}})$, $V \in \mathrm{Op}(S_{\mathrm{sa}})$,

such that $F|_{U' \times V}$ admits a free $p^{-1}\mathcal{O}_S$ -resolution F^\bullet of finite length. It follows that ρ'_*F^\bullet is a $\mathcal{H}om_{\rho'_*p^{-1}\mathcal{O}_S}(\cdot, \mathcal{G})$ -injective resolution of $\rho'_*F|_{U' \times V}$, for any $\mathcal{G} \in \text{Mod}(\rho'_*p^{-1}\mathcal{O}_S|_{U' \times V})$. Hence we are reduced to the case $F = p^{-1}\mathcal{O}_S$, which is clear. q.e.d.

Corollary 3.20. *Let F be a $p^{-1}\mathcal{O}_S$ -coherent S -locally constant sheaf on $U \times S$. Then we have a natural isomorphism in $\text{Mod}_{\text{coh}}(\mathcal{D}_{U \times S/S})$:*

$$\text{RH}^S(\mathbf{D}'F)[-n] \simeq F \otimes_{p^{-1}\mathcal{O}_S} \mathcal{O}_{U \times S}$$

Proof. We have

$$\text{RH}^S(\mathbf{D}'F)[-n] = \rho'^{-1} R\mathcal{H}om_{\rho'_*p^{-1}\mathcal{O}_S}(\rho'_*\mathbf{D}'F, \mathcal{O}_{U \times S}^{t,S,\#})$$

Then we apply (3.18*) with $\mathcal{L} = \mathcal{O}_{U \times S}^{t,S,\#}$, recalling that ρ'^{-1} commutes with tensor products and that $\rho'^{-1}\mathcal{O}_{U \times S}^{t,S,\#} \simeq \mathcal{O}_{U \times S}$. q.e.d.

Lemma 3.21. *With the preceding notations, let $F \in \mathbf{D}_{\mathbb{R}\text{-c}}^b(p_U^{-1}\mathcal{O}_S)$. Then there are natural isomorphisms in $\mathbf{D}^b(\mathcal{D}_{X \times S/S})$*

$$(3.21*) \quad \text{TH}^S(j_!F) \simeq \rho'^{-1} Rj_* R\mathcal{H}om_{\rho'_*p_U^{-1}\mathcal{O}_S}(\rho'_*F, j^{-1}\mathfrak{D}_{X \times S}^{t,S,\#}),$$

$$(3.21**) \quad \text{RH}^S(j_!F)[-d_X] \simeq \rho'^{-1} Rj_* R\mathcal{H}om_{\rho'_*p_U^{-1}\mathcal{O}_S}(\rho'_*F, j^{-1}\mathcal{O}_{X \times S}^{t,S,\#}).$$

Moreover, if $U = X \setminus D$, where D is a normal crossing divisor, and F is $p^{-1}\mathcal{O}_S$ -locally free of finite rank,

$$\text{RH}^S(j_!F)[-d_X]$$

is concentrated in degree zero.

Proof. The proofs of (3.21*) and (3.21**) are similar so we only prove (3.21**). Let us start by noting that, from [11, (2.4.4), Prop.2.4.4], one deduces an isomorphism of functors on $\mathbf{D}^b(\mathbb{C}_{X_{sa} \times S_{sa}})$

$$R\mathcal{H}om(\rho'_*j_!\mathbb{C}_{U \times S}, (\cdot)) \longrightarrow Rj_* R\mathcal{H}om(\rho'_*\mathbb{C}_{U \times S}, j^{-1}(\cdot)) \simeq Rj_* j^{-1}(\cdot)$$

using the following facts:

- $\rho'_*j_!\mathbb{C}_{U \times S} = (j_!\mathbb{C}_{U \times S})_{X_{sa} \times S_{sa}}$ since $X_{sa} \times S_{sa}$ is a \mathcal{T}' -space as explained in the beginning of this section.
- One derives isomorphism (2.4.4) of loc. cit. using injective resolutions since j^{-1} transforms injective objects into injective objects.

We have:

$$\begin{aligned} \text{RH}^S(j_!F) &\simeq \rho'^{-1} R\mathcal{H}om_{\rho'_*p_X^{-1}\mathcal{O}_S}(\rho'_*j_!F, \mathcal{O}_{X \times S}^{t,S,\#}) \\ &\simeq \rho'^{-1} R\mathcal{H}om_{\rho'_*p_X^{-1}\mathcal{O}_S}(\rho'_*(Rj_*F \otimes j_!\mathbb{C}_{U \times S}), \mathcal{O}_{X \times S}^{t,S,\#}) \\ &\stackrel{(1)}{\simeq} \rho'^{-1} R\mathcal{H}om_{\rho'_*p_X^{-1}\mathcal{O}_S}(\rho'_*Rj_*F \otimes \rho'_*j_!\mathbb{C}_{U \times S}, \mathcal{O}_{X \times S}^{t,S,\#}) \\ &\simeq \rho'^{-1} R\mathcal{H}om_{\rho'_*p_X^{-1}\mathcal{O}_S}(\rho'_*Rj_*F, R\mathcal{H}om(\rho'_*j_!\mathbb{C}_{U \times S}, \mathcal{O}_{X \times S}^{t,S,\#})) \\ &\stackrel{(2)}{\simeq} \rho'^{-1} R\mathcal{H}om_{\rho'_*p_X^{-1}\mathcal{O}_S}(Rj_*\rho'_*F, Rj_*j^{-1}\mathcal{O}_{X \times S}^{t,S,\#}) \\ &\stackrel{(3)}{\simeq} \rho'^{-1} Rj_* R\mathcal{H}om_{\rho'_*p_U^{-1}\mathcal{O}_S}(\rho'_*F, j^{-1}\mathcal{O}_{X \times S}^{t,S,\#}). \end{aligned}$$

The isomorphism (1) follows from Corollary 3.6(a), (2) follows from the commutation of ρ'_* and j_* and the preceding remark, and (3) follows by adjunction.

Let us now prove the second part of the statement. We use a similar argument as in [6, Lem. 7.4]. We assume that F is S -locally free of rank ℓ and let $d_X = n$. Note that, according to Example 3.17(1), the result is true if F is constant. Note also that the commutation of ρ'^{-1} with j^{-1} together with Corollary 3.20 imply that $\mathrm{RH}^S(j_!F)[-d_X]|_{(X \setminus Y) \times S}$ is concentrated in degree zero.

By [15, Prop. 5.2i)] and the exactness of $j^{-1}, j^{-1} \mathfrak{D}_{X \times S}^{t, S, \#}$ is concentrated in degree zero, hence, by the assumption on F

$$R\mathcal{H}om_{\rho'_* p_U^{-1} \mathcal{O}_S}(\rho'_* F, j^{-1} \mathfrak{D}_{X \times S}^{t, S, \#})$$

is also concentrated in degree zero.

Let us assume that $D = \{x_1 \cdots x_d = 0\}$ where (x_1, \dots, x_n) are local coordinates in a neighbourhood $\mathrm{nb}_X(x_o)$ of $x_o \in D$. Let also s denote a local coordinate in $W \subset S$. Let us identify $\mathrm{nb}_D(x_o)$ with an open subset of \mathbb{C}^n and W to an open subset in \mathbb{C} . Let U' denote an open ball in $\mathrm{nb}_D(x_o)$ and V denote an open ball in W . Let G and

$$(T_i(s), A_i(s))_{i=1, \dots, d}, \quad s \in V$$

be given by Proposition A.9 and Theorem 2.6 with respect to F , such that $T_i(s) = \exp(-2i\pi A_i(s)), i = 1, \dots, d$.

Let ϕ be a section of $\mathcal{H}om_{\rho'_* p_U^{-1} \mathcal{O}_S}(\rho'_* F, j^{-1} \mathfrak{D}_{X \times S}^{t, S, \#})$ defined on $(U' \setminus D) \times V$. Note that, on any open set of the form $\gamma \times V$ with γ open subanalytic simply connected in $U' \setminus D$,

$$\exp(A_1(s) \log x_1) \cdots \exp(A_d(s) \log x_d)$$

is a matrix with holomorphic entries which are tempered in $X \times V$ (cf. [15, Ex. 5.1]). Then, the $\rho'_* p^{-1} \mathcal{O}_S$ linearity of ϕ implies that

$$\exp(A_1(s) \log x_1) \cdots \exp(A_d(s) \log x_d) \phi$$

is a well-defined section of $\mathcal{H}om_{\rho'_* p_U^{-1} \mathcal{O}_S}(\rho'_* G, j^{-1} \mathfrak{D}_{X \times S}^{t, S, \#})$ on $(U' \setminus D) \times V$. Since the open sets of the form $(U' \setminus D) \times V$ form a basis of the topology in $U_{X_{\mathrm{sa}}} \times S_{\mathrm{sa}}$, this means that the multiplication by $\prod_i \exp(A_i(s) \log x_i)$ defines an isomorphism

$$(3.22) \quad \mathcal{H}om_{\rho'_* p_U^{-1} \mathcal{O}_S}(\rho'_* F, j^{-1} \mathfrak{D}_{X \times S}^{t, S, \#}) \simeq \mathcal{H}om_{\rho'_* p_U^{-1} \mathcal{O}_S}(\rho'_* p^{-1} G, j^{-1} \mathfrak{D}_{X \times S}^{t, S, \#})$$

where $p^{-1}G$ is S -constant and free. Therefore the righthand side of (3.22) is isomorphic to $(j^{-1} \mathfrak{D}_{X \times S}^{t, S, \#})^\ell$. By the $\Gamma(U' \times V', \bullet)$ -acyclicity of $\mathfrak{D}_{X \times S}^{t, S, \#}$ for arbitrary relatively compact subanalytic sets U' and V' in X and S respectively (cf. [15, Prop. 5.2(i)]), we have $Rj_* j^{-1} \mathfrak{D}_{X \times S}^{t, S, \#} \simeq j_* j^{-1} \mathfrak{D}_{X \times S}^{t, S, \#}$, that is, it is concentrated in degree zero, hence $\mathrm{TH}^S(j_!F)$ is concentrated in degree zero. Moreover, by the construction, the isomorphism (3.22) preserves the actions of $j^{-1} \rho'_! \mathcal{O}_{X \times S}$ -modules and of $j^{-1} \rho'_! \mathcal{D}_{\overline{X \times S}}$ -modules. Thus, by (3.13), (3.22) induces an isomorphism of $\rho'^{-1} j_* j^{-1} \rho'_! \mathcal{O}_{X \times S}$ -modules, hence of $\rho'^{-1} \rho'_! \mathcal{O}_{X \times S} \simeq \mathcal{O}_{X \times S}$ -modules

$$\mathrm{RH}^S(j_!F)[-n] \simeq \mathrm{RH}^S(j_! D' p^{-1} G)[-n].$$

So we may assume from the beginning that F is constant and the result follows. q.e.d.

3.6. Functorial properties. Let $f : Y \rightarrow X$ be a morphism of real or complex analytic manifolds. We shall still denote by f the associated morphism $f \times \text{Id}_S : Y \times S \rightarrow X \times S$. We shall study the associated derived functor ${}_D f_*$. We begin with the relative version of [6, Th. 4.1]:

Theorem 3.23. *Let $f : Y \rightarrow X$ be a morphism of real analytic manifolds, let $F \in \mathbb{D}_{\mathbb{R}\text{-c}}^b(p_Y^{-1}\mathcal{O}_S)$, and assume that f is proper on $\text{Supp } F$. Then we have a canonical isomorphism in $\mathbb{D}(\mathcal{D}_{X \times S/S})$:*

$${}_D f_* \text{TH}^S(F) \simeq \text{TH}^S(Rf_* F).$$

Proof. We can replace F with a complex as in Proposition 3.4 and we argue by induction on its length, so it is sufficient to assume that F is of the form $\mathbb{C}_{U \times V} \otimes p_Y^{-1}\mathcal{O}_S$, where U (resp. V) is a relatively compact subanalytic open subset of Y (resp. S). In that case, one has $\text{TH}^S(F) \simeq T\mathcal{H}om(\mathbb{C}_{U \times V}, \mathfrak{D}\mathfrak{b}_{Y \times S})$. On the other hand

$$Rf_*(\mathbb{C}_{U \times V} \otimes p_Y^{-1}\mathcal{O}_S) \simeq Rf_*(\mathbb{C}_{U \times V}) \otimes p^{-1}\mathcal{O}_S$$

hence

$$\text{TH}^S(Rf_*(\mathbb{C}_{U \times V} \otimes p_Y^{-1}\mathcal{O}_S)) \simeq T\mathcal{H}om(Rf_*(\mathbb{C}_U) \otimes \mathbb{C}_{X \times V}, \mathfrak{D}\mathfrak{b}_{X \times S}).$$

Therefore the statement follows by the absolute case in [6, Th. 4.1].

q.e.d.

Recalling (3.13) and adapting [6, Lem. 7.2] one obtains:

Theorem 3.24. *Let $f : Y \rightarrow X$ be a morphism of complex analytic manifolds, let $F \in \mathbb{D}_{\mathbb{R}\text{-c}}^b(p_Y^{-1}\mathcal{O}_S)$, and assume that f is proper on $\text{Supp } F$. Then we have a canonical isomorphism in $\mathbb{D}(\mathcal{D}_{X \times S/S})$:*

$${}_D f_* \text{RH}^S(F) \simeq \text{RH}^S(Rf_* F).$$

Proposition 3.25. *For any $F \in \mathbb{D}_{\mathbb{R}\text{-c}}^b(p_Y^{-1}\mathcal{O}_S)$ and any $s_o \in S$, there is a natural morphism*

$$Li_{s_o}^* \text{RH}^S(F)[-d_X] \longrightarrow T\mathcal{H}om(Li_{s_o}^* F, \mathcal{O}_X)$$

which is an isomorphism, where we identify X with $X \times \{s_o\}$ and X_{sa} with $X_{\text{sa}} \times \{s_o\}$.

Proof. Let us construct the morphism. We have

$$\begin{aligned} Li_{s_o}^* \text{RH}^S(F)[-d_X] &= p^{-1}(\mathcal{O}_S/\mathfrak{m}_{s_o}) \overset{L}{\otimes}_{p^{-1}\mathcal{O}_S} \text{RH}^S(F)[-d_X] \\ &\simeq \rho'^{-1} \left(\rho'_*(p^{-1}(\mathcal{O}_S/\mathfrak{m}_{s_o})) \overset{L}{\otimes}_{\rho'_* p^{-1}\mathcal{O}_S} R\mathcal{H}om_{\rho'_* p^{-1}\mathcal{O}_S}(\rho'_* F, \mathcal{O}_{X \times S}^{t,S,\sharp}) \right) \\ &\stackrel{(*)}{\simeq} \rho'^{-1} R\mathcal{H}om_{\rho'_*(p^{-1}(\mathcal{O}_S/\mathfrak{m}_{s_o}))} \left(\rho'_*(p^{-1}(\mathcal{O}_S/\mathfrak{m}_{s_o})) \overset{L}{\otimes}_{p^{-1}\mathcal{O}_S} F, \right. \\ &\quad \left. \rho'_*(p^{-1}(\mathcal{O}_S/\mathfrak{m}_{s_o})) \overset{L}{\otimes}_{\rho'_* p^{-1}\mathcal{O}_S} \mathcal{O}_{X \times S}^{t,S,\sharp} \right) \\ &\simeq \rho'^{-1} R\mathcal{H}om_{\rho'_*(p^{-1}(\mathcal{O}_S/\mathfrak{m}_{s_o}))} \left(\rho'_* Li_{s_o}^* F, \rho'_*(p^{-1}(\mathcal{O}_S/\mathfrak{m}_{s_o})) \overset{L}{\otimes}_{\rho'_* p^{-1}\mathcal{O}_S} \mathcal{O}_{X \times S}^{t,S,\sharp} \right), \end{aligned}$$

where $(*)$ uses Corollary 3.6(b). So it remains to show

$$\rho'_*(p^{-1}(\mathcal{O}_S/\mathfrak{m}_{s_o})) \overset{L}{\otimes}_{\rho'_* p^{-1}\mathcal{O}_S} \mathcal{O}_{X \times S}^{t,S,\sharp} |_{X_{\text{sa}} \times \{s\}} \simeq \mathcal{O}_X^t$$

Since $Li_{s_o}^*$ commutes with $R\mathcal{H}om_{\mathcal{D}_{\overline{X \times S}}}(\mathcal{O}_{\overline{X \times S}}, \bullet)$ it is sufficient to show

$$\rho'_*(p^{-1}(\mathcal{O}_S/\mathfrak{m}_{s_o})) \overset{L}{\otimes}_{\rho'_* p^{-1}\mathcal{O}_S} \mathfrak{D}\mathfrak{b}_{X \times S}^{t,S,\sharp} |_{X_{\text{sa}} \times \{s\}} \simeq \mathfrak{D}\mathfrak{b}_X^t$$

Taking any local coordinate s on S centered at s_o , this amounts to showing

$$\{\Gamma(U \times V; \mathfrak{D}\mathfrak{b}_{X \times S}^{t,S;\#}) \xrightarrow{s} \Gamma(U \times V; \mathfrak{D}\mathfrak{b}_{X \times S}^{t,S;\#})\} \simeq \Gamma(U; \mathfrak{D}\mathfrak{b}_X^t)$$

for any relatively compact open subsets $U \subset X$, $V \subset S$. We note that

- $\Gamma(U \times V; \mathfrak{D}\mathfrak{b}_{X \times S}^{t,S;\#}) = \Gamma(X \times V, T\mathcal{H}om(U \times S, \mathfrak{D}\mathfrak{b}_{X \times S}))$,
- $Li_{s_o}^* T\mathcal{H}om(U \times S, \mathfrak{D}\mathfrak{b}_{X \times S})|_{\{s_o\}} \simeq T\mathcal{H}om(U, \mathfrak{D}\mathfrak{b}_X)$ (cf. [10, Th. 4.5 (4.8)]).

Therefore

$$\Gamma(X \times V; T\mathcal{H}om(U \times S, \mathfrak{D}\mathfrak{b}_{X \times S})) \xrightarrow{s} \Gamma(X \times V; T\mathcal{H}om(U \times S, \mathfrak{D}\mathfrak{b}_{X \times S}))$$

is quasi-isomorphic to

$$\Gamma(X; T\mathcal{H}om(U, \mathfrak{D}\mathfrak{b}_X)) = \Gamma(U; \mathfrak{D}\mathfrak{b}_X^t),$$

which gives the desired result. q.e.d.

4. Proof of the main results

In order to apply the results of §§3.2–3.6, we continue assuming that $d_S = 1$.

Remark 4.1 (The locally constant case). In view of Corollary 3.20 and Remark A.10, Theorem 3 is true if F is an S -locally constant sheaf. Similarly, the isomorphism of Theorem 5 holds for $\mathcal{M} = F \otimes_{p^{-1}\mathcal{O}_S} \mathcal{O}_{X \times S}$. Moreover, we recover Deligne’s Riemann-Hilbert correspondence by means of RH^S as an equivalence between the category of S -local systems on $X \times S$ and the category of coherent $\mathcal{O}_{X \times S}$ -modules endowed with a relative flat connexion.

4.1. Proof of Theorem 3. We first consider the setting of Section 2.2. Given an S -locally constant sheaf F of $p_{X \setminus D}^{-1}\mathcal{O}_S$ -modules, we set $E_F := \mathcal{O}_{(X \setminus D) \times S} \otimes_{p_{X \setminus D}^{-1}\mathcal{O}_S} F$. The $\mathcal{O}_{X \times S}$ -module j_*E_F carries a natural structure of $\mathcal{D}_{X \times S/S}$ -module. According to Corollary 2.8, if F is $p^{-1}\mathcal{O}_S$ -coherent, \widetilde{E}_F is regular holonomic and has a characteristic variety contained in $(\pi^{-1}(D) \times S) \cup (T_X^*X \times S)$, where π is the projection from T^*X to X . Moreover $\widetilde{E}_F \simeq \widetilde{E}_F(*D)$, hence $R\Gamma_{[D \times S]} \widetilde{E}_F = 0$.

Lemma 4.2. *Assume that F is $p^{-1}\mathcal{O}_S$ -locally free of finite rank. Then we have, functorially in F ,*

$$\widetilde{E}_F \simeq \text{RH}^S(\text{PSol } \widetilde{E}_F).$$

Proof. We first rewrite the right-hand side as $\text{RH}^S(j_! \mathbf{D}'F)[-d_X]$. Indeed, since $\mathbf{D}'F = \text{Sol } E_F$ on $X \setminus D$, we have a natural morphism $j_! \mathbf{D}'F \rightarrow \text{Sol } \widetilde{E}_F$. We will prove that it is an isomorphism. Since both complexes are S - \mathbb{C} -constructible, we are left with proving the same property after applying $Li_{s_o}^*$ for any $s_o \in S$, according to [16, Prop. 2.2]. On the other hand, $Li_{s_o}^* \text{Sol } \widetilde{E}_F = \text{Sol } Li_{s_o}^* \widetilde{E}_F$ after [16, Prop. 2.1], and $\text{Sol } Li_{s_o}^* \widetilde{E}_F = \text{Sol } i_{s_o}^* \widetilde{E}_F$ since \widetilde{E}_F is strict (Corollary 2.8). Moreover, $i_{s_o}^* \widetilde{E}_F = \widetilde{i_{s_o}^* E_F}$ (Lemma 2.12). Similarly, one checks that $Li_{s_o}^* j_! \mathbf{D}'F = j_! \mathbf{D}'i_{s_o}^* F$. It is well-known that $j_! \mathbf{D}'i_{s_o}^* F \rightarrow \text{Sol } \widetilde{i_{s_o}^* E_F}$ is an isomorphism, hence the desired assertion.

According to Lemma 3.21, the complex $\mathrm{RH}^S(j_! \mathbf{D}'F)[-d_X]$ is concentrated in degree zero, and according to Lemma 3.18 (with $\mathcal{H} = j^{-1} \mathcal{O}_{X \times S}^{t, S, \sharp}$), (3.21**) and the $p^{-1} \mathcal{O}_S$ -flatness of F we have

$$\begin{aligned} \mathrm{RH}^S(j_! \mathbf{D}'F)[-d_X] &\simeq \rho'^{-1} Rj_* R\mathcal{H}om_{\rho'_* p^{-1} \mathcal{O}_S}(\rho'_* \mathbf{D}'F, j^{-1} \mathcal{O}_{X \times S}^{t, S, \sharp}) \\ &\simeq \rho'^{-1} Rj_*(\rho'_* F \otimes_{\rho'_* p^{-1} \mathcal{O}_S}^L j^{-1} \mathcal{O}_{X \times S}^{t, S, \sharp}) \\ &\simeq \rho'^{-1} Rj_*(\rho'_* F \otimes_{\rho'_* p^{-1} \mathcal{O}_S} j^{-1} \mathcal{O}_{X \times S}^{t, S, \sharp}). \end{aligned}$$

We shall prove that $\rho'^{-1} Rj_*(\rho'_* F \otimes_{\rho'_* p_{X \setminus D}^{-1} \mathcal{O}_S} j^{-1} \mathcal{O}_{X \times S}^{t, S, \sharp})$ coincides with \widetilde{E}_F . Firstly, applying the commutation of ρ'^{-1} with j^{-1} together Corollary 3.20 entails that $\rho'^{-1} Rj_*(\rho'_* F \otimes_{\rho'_* p_{X \setminus D}^{-1} \mathcal{O}_S} j^{-1} \mathcal{O}_{X \times S}^{t, S, \sharp})$ and \widetilde{E}_F coincide on $(X \setminus D) \times S$. Therefore it is enough to prove that, for each $(y, s_o) \in D \times S$, for any $W \in \mathrm{Op}(X)$ running in a basis of neighborhoods of y and for any $V \in \mathrm{Op}(S_{\mathrm{sa}})$ running in a basis of neighborhoods of s_o , we have

$$(a) \quad \varprojlim_{\substack{V' \in \mathrm{Op}^c(S_{\mathrm{sa}}), s_o \in V', V' \in V \\ U \in \mathrm{Op}^c(X_{\mathrm{sa}}), y \in U, U \in W}} \Gamma((U \setminus D) \times V'; \rho'_* F \otimes_{\rho'_* p_{X \setminus D}^{-1} \mathcal{O}_S} j^{-1} \mathcal{O}_{X \times S}^{t, S, \sharp}) = \Gamma((W \setminus D) \times V; \widetilde{E}_F).$$

Recall that, by definition of \otimes , the subanalytic sheaf $\rho'_* F \otimes_{\rho'_* p_{X \setminus D}^{-1} \mathcal{O}_S} j^{-1} \mathcal{O}_{X \times S}^{t, S, \sharp}$ is the sheaf associated to the presheaf defined by the formula:

$$\omega \times \omega' \mapsto \Gamma(\omega \times \omega'; F) \otimes \Gamma(\omega \times \omega'; j^{-1} \mathcal{O}_{X \times S}^{t, S, \sharp}), \quad \omega \in \mathrm{Op}((X \setminus D)_{X_{\mathrm{sa}}}), \omega' \in \mathrm{Op}(S_{\mathrm{sa}}).$$

(b) Therefore a section h in

$$\Gamma((U \setminus D) \times V'; \rho'_* F \otimes_{\rho'_* p_{X \setminus D}^{-1} \mathcal{O}_S} j^{-1} \mathcal{O}_{X \times S}^{t, S, \sharp})$$

is uniquely determined by the data of an open covering of $U \setminus D$ by simply connected Stein open subanalytic sets $(U_\beta)_{\beta \in B}$ and of a family $(h_\beta)_{\beta \in B}$ of vectors of ℓ holomorphic functions, $h_\beta = (h_{i, \beta})_{i=1, \dots, \ell}$, such that, for each $i = 1, \dots, \ell$, $h_{i, \beta}$ is a holomorphic function defined in $U_\beta \times V'$ tempered in $X \times V'$ and such that the h_β have the monodromy of F .

Taking local coordinates (x_1, \dots, x_n) in X , s in S , such that D is given by an equation $x_1 \cdots x_d = 0$ in a neighbourhood of $y = 0 \in D$, we may assume that

$$\begin{aligned} W &= B_\delta(0) = \{(x_1, \dots, x_n), |x_j| < \delta, j = 1, \dots, d\}, \quad V = B_\delta(s_o), \\ U &= B_\varepsilon(0) \text{ and } V' = B_\varepsilon(s_o), \end{aligned}$$

for some $\delta > 0$ sufficiently small and arbitrary ε satisfying $0 < \varepsilon < \delta$.

(1) Let $f \in \Gamma((W \setminus D) \times V; \widetilde{E}_F)$. We can decompose $W \setminus D$ as a union of a finite family $(W_{\alpha, \varepsilon_\alpha})_{\alpha \in A, \varepsilon_\alpha > 0}$ of open convex subsets such that, for each $(\alpha, \varepsilon_\alpha)$ and for each U , $W_{\alpha, \varepsilon_\alpha} \cap U := U_{\alpha, \varepsilon_\alpha}^*$ is a convex open subset (hence Stein) in the conditions of Definition 2.5.

For each $\alpha \in A$ we can choose an isomorphism

$$\psi_{\alpha, \varepsilon_\alpha} : F|_{W_{\alpha, \varepsilon_\alpha} \times V} \simeq p^{-1} \mathcal{O}_S^\ell|_{W_{\alpha, \varepsilon_\alpha} \times V}$$

which induces an isomorphism

$$\phi_{\alpha, \varepsilon_\alpha} : \widetilde{E}_F|_{W_{\alpha, \varepsilon_\alpha} \times V} \simeq \mathcal{O}_{X \times S}^\ell|_{W_{\alpha, \varepsilon_\alpha} \times V}.$$

Then, setting $\phi_{\alpha, \varepsilon_\alpha}(f) := f_{\alpha, \varepsilon_\alpha}$, the family $(f_{\alpha, \varepsilon_\alpha})_{\alpha \in A}$ has the monodromy of F .

Let $f_{i, \alpha, \varepsilon_\alpha}$ denote the i component of $f_{\alpha, \varepsilon_\alpha}$, $i = 1, \dots, \ell$. By construction, each $f_{i, \alpha, \varepsilon_\alpha}$ is holomorphic (hence tempered) at any (x, s) such that $x \in \partial U_{\alpha, \varepsilon_\alpha}^* \setminus D$ and $s \in V$. Hence, by [15, Prop. 5.8], $f_{i, \alpha, \varepsilon_\alpha}$ satisfies the estimation of Definition 2.5 if and only if, for each $i = 1, \dots, \ell$, $\alpha \in A$, $f_{i, \alpha, \varepsilon_\alpha}|_{U_{\alpha, \varepsilon_\alpha}^* \times V'}$ is tempered at $X \times V'$.

Therefore, the family $(f_{\alpha, \varepsilon_\alpha})_{\alpha \in A}$ defines an element of

$$\Gamma((U \setminus D) \times V'; \rho'_* F \otimes_{\rho'_* p_{X \setminus D}^{-1} \mathcal{O}_S} j^{-1} \mathcal{O}_{X \times S}^{t, S, \sharp}).$$

With $\varepsilon \rightarrow \delta$ we obtain f as a section on $(W \setminus D) \times V$ of

$$\rho'^{-1} Rj_*(\rho'_* F \otimes_{\rho'_* p_{X \setminus D}^{-1} \mathcal{O}_S} j^{-1} \mathcal{O}_{X \times S}^{t, S, \sharp}).$$

(2) By the characterization of the elements of

$$\Gamma((U \setminus D) \times V'; \rho'_* F \otimes_{\rho'_* p_{X \setminus D}^{-1} \mathcal{O}_S} j^{-1} \mathcal{O}_{X \times S}^{t, S, \sharp})$$

given in (b), the converse is similar. q.e.d.

Lemma 4.3. *Let F be an S -locally constant coherent $p^{-1} \mathcal{O}_S$ -module on $X \setminus D$. Then $\mathrm{RH}^S(j_! F) \in \mathrm{D}_{\mathrm{rhol}}^b(\mathcal{D}_{X \times S/S})$.*

Proof. One considers the exact sequence of S -local systems

$$0 \longrightarrow F_{\mathrm{tors}} \longrightarrow F \longrightarrow F_{\mathrm{lf}} \longrightarrow 0$$

where F_{tors} denotes the S -local system of $p^{-1} \mathcal{O}_S$ -torsion sections of F and F_{lf} denotes the quotient F/F_{tors} . According to Lemma 4.2 and Theorem 2.6, the result holds for F_{lf} . By the functoriality of RH^S , it will hold for F provided it holds for F_{tors} .

So we now assume that F is a torsion module. By definition, F is $p_{X \setminus D}^{-1} \mathcal{O}_S$ -coherent so the support of F is contained in $p_{X \setminus D}^{-1} S_0$, where S_0 is a discrete subset of S . Let us consider $s_o \in S_0$, $x_o \in D$ and let us prove that $\mathrm{RH}^S(j_! F)$ is regular holonomic in a neighborhood of (x_o, s_o) . If s is a local coordinate vanishing at s_o , we can choose a power N such that $s^N F = 0$. Arguing by induction on N , one easily reduces to the case $N = 1$. In that case, F is isomorphic to $F' \boxtimes \mathcal{O}_S/s \mathcal{O}_S$ for some \mathbb{C} -local system F' on $X \setminus D$. We have

$$\begin{aligned} \mathrm{RH}^S(j_! F) &= \rho'^{-1} R\mathcal{H}om_{\rho'_* p_{X \setminus D}^{-1} \mathcal{O}_S}(\rho'_* j_! F, \mathcal{O}_{X \times S}^{t, S, \sharp}) \\ &\simeq \rho'^{-1} R\mathcal{H}om(\rho'_*(j_!(F' \boxtimes \mathcal{O}_S/s \mathcal{O}_S)), \mathcal{O}_{X \times S}^{t, S, \sharp}). \end{aligned}$$

On the other hand

$$\begin{aligned} j_!(F' \boxtimes \mathcal{O}_S/s \mathcal{O}_S) &\simeq j_!((F' \boxtimes \mathbb{C}_S) \otimes_{p_{X \setminus D}^{-1} \mathcal{O}_S} (\mathcal{O}_S/s \mathcal{O}_S)) \\ &\simeq j_!(F' \boxtimes \mathbb{C}_S) \otimes_{p_X^{-1} \mathcal{O}_S} (\mathcal{O}_S/s \mathcal{O}_S) \\ &\simeq (j_! F' \boxtimes \mathbb{C}_S) \otimes_{p_X^{-1} \mathcal{O}_S} (\mathcal{O}_S/s \mathcal{O}_S), \end{aligned}$$

hence, according to [15, Prop. 4.7(1)], we get

$$\mathrm{RH}^S(j_! F) \simeq R\mathcal{H}om_{p_X^{-1} \mathcal{O}_S}(p_X^{-1} (\mathcal{O}_S/s \mathcal{O}_S), T\mathcal{H}om(j_! F' \boxtimes \mathbb{C}_S, \mathcal{O}_{X \times S})).$$

Since $T\mathcal{H}om(j_! F' \boxtimes \mathbb{C}_S, \mathcal{O}_{X \times S})$ is in $\mathrm{Mod}_{\mathrm{rhol}}(\mathcal{D}_{X \times S/S})$, the result follows. q.e.d.

End of the proof of Theorem 3. We have the tools to follow the outline of Kashiwara's proof in the absolute case [6, §7.3]. We may assume that F is an S - \mathbb{C} -constructible sheaf. Then we argue by induction on the dimension of a closed analytic set Z such that $Z \times S \subset \text{Supp } F$. Let Z_0 be a closed analytic subset of Z such that

- (1) $F|_{(Z \setminus Z_0) \times S}$ is locally constant over $p^{-1}\mathcal{O}_S$.
- (2) $Z \setminus Z_0$ is non singular.

By the induction hypothesis, $\text{RH}^S(F_{Z_0 \times S})$ belongs to $\text{D}_{\text{rhol}}^b(\mathcal{D}_{X \times S/S})$. So we may assume that $F_{Z_0 \times S} = 0$. The question is local on Z . Consider then a projective morphism $\pi : X' \rightarrow X$ such that X' is non singular and $\pi(X') = Z$, $Z'_0 := \pi^{-1}(Z_0)$ is a normal crossing divisor in X' and $\pi : X' \setminus Z'_0 \rightarrow Z \setminus Z_0$ is an isomorphism. Let $F' = \pi^{-1}(F)$. Then we obtain that $F'|_{Z'_0 \times S} = 0$, $F'|_{(X' \setminus Z'_0) \times S}$ is locally constant, and $R\pi_* F' = F$. Now the first part of Theorem 3 follows straightforwardly from Proposition 3.24 and Lemma 4.3.

For the second part, we note that, since $\text{RH}^S(F)$ is holonomic by the first part, Lemma 3.16, together with [16, Cor. 3.9], implies $\mathbf{D}^{\text{pSol}}(\text{RH}^S(F)) \simeq \mathbf{D}F$, and we conclude by bi-duality [16, Prop. 2.23]. As a consequence, pSol is essentially surjective from $\text{D}_{\text{rhol}}^b(\mathcal{D}_{X \times S/S})$ to $\text{D}_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$. q.e.d.

Proof of Corollary 4. Set $\mathcal{M} = \text{RH}^S(F)$. By Proposition 2, it is enough to check that, if F and $\mathbf{D}F$ are perverse, then so are $\text{pSol } \mathcal{M}$ and $\text{pDR } \mathcal{M}$. But $\text{pSol } \mathcal{M} \simeq F$ by Theorem 3, and Lemma 3.16 gives $\text{pDR } \mathcal{M} \simeq \mathbf{D}F$. q.e.d.

4.2. Proof of Theorem 5. Let \mathcal{M}, \mathcal{N} be holonomic $\mathcal{D}_{X \times S/S}$ -modules and let F be an object of $\text{D}_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$. Recall (cf. [9, (2.6.7)]) that we have a natural isomorphism in $\text{D}(p^{-1}\mathcal{O}_S)$:

$$R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, R\mathcal{H}om_{p^{-1}\mathcal{O}_S}(F, \mathcal{O}_{X \times S})) \simeq R\mathcal{H}om_{p^{-1}\mathcal{O}_S}(F, \text{Sol } \mathcal{M}),$$

which is bi-functorial with respect to \mathcal{M}, F . By composing with (3.15), we obtain a bi-functorial morphism

$$R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \text{RH}^S(F)[-d_X]) \longrightarrow R\mathcal{H}om_{p^{-1}\mathcal{O}_S}(F, \text{Sol } \mathcal{M}).$$

Choosing $F = \text{Sol } \mathcal{N}$ finally produces a bi-functorial morphism

$$(4.4) \quad R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \text{RH}^S(\text{pSol } \mathcal{N})) \longrightarrow R\mathcal{H}om_{p^{-1}\mathcal{O}_S}(\text{Sol } \mathcal{N}, \text{Sol } \mathcal{M}).$$

If (4.4) is an isomorphism, then by taking global sections we find a bi-functorial isomorphism

$$\text{Hom}_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \text{RH}^S(\text{pSol } \mathcal{N})) \longrightarrow \text{Hom}_{p^{-1}\mathcal{O}_S}(\text{Sol } \mathcal{N}, \text{Sol } \mathcal{M})$$

and the isomorphism (*) stated in Theorem 5 is obtained as that corresponding to $\text{Id}_{\text{Sol } \mathcal{M}}$ when $\mathcal{N} = \mathcal{M}$, while (**) follows by applying (*) to \mathcal{N} . We consider the following three statements.

- (a) If \mathcal{M}, \mathcal{N} underlie objects of $\text{MTM}^{\text{reg}}(X)$, then the complex

$$R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \text{RH}^S(\text{pSol } \mathcal{N}))$$

is S - \mathbb{C} -constructible.

- (b) If \mathcal{M}, \mathcal{N} underlie objects of $\text{MTM}^{\text{reg}}(X)$, then (4.4) is an isomorphism.
- (c) If \mathcal{M} underlies an object of $\text{MTM}^{\text{reg}}(X)$, then so does $\text{RH}^S(\text{pSol } \mathcal{M})$.

We also denote by $(a)_n$ (resp. $(b)_n$, $(c)_n$) the statement (a) (resp. (b), (c)) for \mathcal{M}, \mathcal{N} with support in X of dimension $\leq n$. The first part of Theorem 5 follows from $(b)_n$ for any $n \geq 0$ by setting $\mathcal{N} = \mathcal{M}$. We will prove $(c)_n$ for any $n \geq 0$, which will be enough, according to the lemma below.

Lemma 4.5. *For any $n \geq 0$, the statements $(a)_n$, $(b)_n$ and $(c)_n$ are equivalent.*

Notice already that $(b)_n \Rightarrow (c)_n$ is obtained exactly as (4.4) \Rightarrow Theorem 5.

Proof of $(a)_n \Rightarrow (b)_n$. Assume that $(a)_n$ holds true. We will prove that so does $(b)_n$ by applying the functor $Li_{s_o}^*$ for any $s_o \in S$, to reduce to [6, Cor. 8.6].

Assume that \mathcal{M}, \mathcal{N} underlie objects of $\text{MTM}^{\text{reg}}(X)$ and have support of dimension $\leq n$. According to $(a)_n$ and to [16, Prop. 2.2], (4.4) is an isomorphism as soon as $Li_{s_o}^*(4.4)$ is an isomorphism for any $s_o \in S$, since we already know that $R\mathcal{H}om_{p^{-1}\mathcal{O}_S}(\text{Sol } \mathcal{N}, \text{Sol } \mathcal{M})$ is S - \mathbb{C} -constructible, as $\text{Sol } \mathcal{M}, \text{Sol } \mathcal{N}$ are so (cf. [16, Th. 3.7]).

On the other hand, arguing as for [16, Prop. 2.1] by using [7, (A.10)] (together with [9, (2.6.7)]), $Li_{s_o}^*(4.4)$ is the morphism

$$R\mathcal{H}om_{\mathcal{D}_X}(Li_{s_o}^* \mathcal{M}, Li_{s_o}^* \text{RH}^S(\text{pSol } \mathcal{N})) \longrightarrow R\mathcal{H}om_{\mathbb{C}}(Li_{s_o}^* \text{Sol } \mathcal{N}, Li_{s_o}^* \text{Sol } \mathcal{M}),$$

and still by [16, Prop. 2.1], we can replace the right-hand side with

$$R\mathcal{H}om_{\mathbb{C}}(\text{Sol } Li_{s_o}^* \mathcal{N}, \text{Sol } Li_{s_o}^* \mathcal{M}).$$

By the strictness of \mathcal{M} underlying an object of $\text{MTM}(X)$, we have $Li_{s_o}^* \mathcal{M} = i_{s_o}^* \mathcal{M}$ and, for \mathcal{M} underlying an object of $\text{MTM}^{\text{reg}}(X)$, $i_{s_o}^* \mathcal{M}$ is regular holonomic. The same property applies to \mathcal{N} . On the other hand, by Proposition 3.25, we can replace $Li_{s_o}^* \text{RH}^S(\text{pSol } \mathcal{N})$ with $T\mathcal{H}om(Li_{s_o}^* \text{Sol } \mathcal{N}, \mathcal{O}_X) = T\mathcal{H}om(\text{Sol } i_{s_o}^* \mathcal{N}, \mathcal{O}_X)$. In such a way, $Li_{s_o}^*(4.4)$ is an isomorphism, according to [6, Cor. 8.6], and this ends the proof of $(b)_n$. q.e.d.

Proof of $(c)_n \Rightarrow (a)_n$. This is an immediate consequence of Proposition 4.6 below. q.e.d.

Proposition 4.6. *Let $\mathcal{M}, \mathcal{N} \in \text{Mod}_{\text{hol}}(\mathcal{D}_{X \times S/S})$ and assume that both \mathcal{M} and \mathcal{N} underlie an object of $\text{MTM}(X)$. Then $R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{N})$ is S - \mathbb{C} -constructible.*

Proof. We shall apply the following relative versions of Lemmas 1.8, 1.9 and Proposition 4.7 of [5]. Since they are trivial adaptations of the original ones, we omit their proof.

Lemma 4.7. *Let $\mathcal{M} \in \text{D}^b(\mathcal{D}_{X \times S/S})^{\text{op}}$ and $\mathcal{N} \in \text{D}^b(\mathcal{D}_{X \times S/S})$. Then*

$$R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\Omega_{X \times S/S}^{\dim X}, \mathcal{M} \otimes_{\mathcal{O}_{X \times S}}^L \mathcal{N}) \simeq \mathcal{M} \otimes_{\mathcal{D}_{X \times S/S}}^L \mathcal{N}[-d_X].$$

Lemma 4.8. *Let $\mathcal{M} \in \text{D}_{\text{coh}}^b(\mathcal{D}_{X \times S/S})$ and let $\mathcal{N} \in \text{D}^b(\mathcal{D}_{X \times S/S})$. Then we have an isomorphism*

$$R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{N}) \simeq R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\Omega_{X \times S/S}^n, \mathbf{D}' \mathcal{M} \otimes_{\mathcal{O}_{X \times S}}^L \mathcal{N})[d_X].$$

Lemma 4.9. *Let \mathcal{M} and \mathcal{N} be two strict $\mathcal{D}_{X \times S/S}$ -modules. Then*

$$\mathcal{M} \otimes_{\mathcal{O}_{X \times S}}^L \mathcal{N} \simeq {}_{\mathbb{D}}i_X^*(\mathcal{M} \boxtimes \mathcal{N}),$$

where X is identified to the diagonal of $X \times X$ by the inclusion i_X and \boxtimes denotes the external product over $p^{-1}\mathcal{O}_S$.

Let us now return to the proof of Proposition 4.6. The assumption on \mathcal{M} and \mathcal{N} entail that $\mathcal{M} \boxtimes \mathcal{N}$ also underlies an object of MTM according to [14, Prop. 11.4.6], so the same condition is satisfied by the cohomology groups of ${}_{\mathbb{D}}i_X^*(\mathcal{M} \boxtimes \mathcal{N})$, in particular they are holonomic. In view of Lemmas 4.7, 4.8 and 4.9, the result follows from [16, Th 3.7] which entails that the de Rham complex of ${}_{\mathbb{D}}i_X^*(\mathcal{M} \boxtimes \mathcal{N})$ is S - \mathbb{C} -constructible. q.e.d.

4.3. End of the proof of Theorem 5

Proof of (c)_n by induction on n. We assume any of (a)_{n-1}, (b)_{n-1} and (c)_{n-1} is true, according to Lemma 4.5.

Reduction to the localized case. The main reason for restricting to the category of $\mathcal{D}_{X \times S/S}$ -modules underlying a regular mixed twistor \mathcal{D} -module is that, if Y is an hypersurface in X , the localization morphism $\mathcal{M} \rightarrow \mathcal{M}(*Y)$ underlies a morphism in MTM^{reg} , as proved in [14, §11.2.2], where the localization functor for mixed twistor \mathcal{D} -modules is denoted by the symbol $[*Y]$. The localization enables us to argue by induction on the dimension of the support of \mathcal{M} .

Since the constructibility is a local property (cf. Section 1.2), the question is local on X . Let $Z \subset X$ denote the support of \mathcal{M} . Locally, there exists an hypersurface Y in X such that $Z^* := Z \setminus Z \cap Y$ is a nonempty smooth submanifold of Z and $\mathcal{M}|_{Z \setminus Y}$ underlies the pushforward of a smooth object of $\text{MTM}^{\text{reg}}(Z^*)$ (i.e., an admissible variation of mixed twistor structure). Since the morphism $\mathcal{M} \rightarrow \mathcal{M}(*Y)$ underlies a morphism in MTM^{reg} , its kernel \mathcal{M}' , image \mathcal{I} and cokernel \mathcal{M}'' underlie objects of MTM^{reg} and \mathcal{M}' , \mathcal{M}'' have support of dimension $< d_Z = n$.

Together with the inductive assumption, we also assume that (c)_n holds true for $\mathcal{M}(*Y)$, and it holds for the kernel and cokernel of $\mathcal{M} \rightarrow \mathcal{M}(*Y)$ by induction.

From the distinguished triangle

$$\text{RH}^S \text{pSol } \mathcal{I} \longrightarrow \text{RH}^S \text{pSol } \mathcal{M}(*Y) \longrightarrow \text{RH}^S \text{pSol } \mathcal{M}'' \xrightarrow{+1}$$

and according to Proposition 4.6 applied to the last two terms, we find that $R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \text{RH}^S(\text{pSol } \mathcal{I}))$ is S - \mathbb{C} -constructible. Then, from the distinguished triangle

$$\text{RH}^S \text{pSol } \mathcal{M} \longrightarrow \text{RH}^S \text{pSol } \mathcal{I} \longrightarrow \text{RH}^S \text{pSol } \mathcal{M}'[1] \xrightarrow{+1}$$

and Proposition 4.6 applied similarly, we conclude that

$$R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \text{RH}^S(\text{pSol } \mathcal{M}))$$

is S - \mathbb{C} -constructible, that is, (a)_n for $\mathcal{N} = \mathcal{M}$. Then (b)_n for $\mathcal{N} = \mathcal{M}$ also holds, and then (c)_n for \mathcal{M} too. q.e.d.

Proof of (c)_n in the localized case. Recall that we work locally on X . We assume now that \mathcal{M} is as above and satisfies $\mathcal{M} = \mathcal{M}(*Y)$. We can then choose a projective morphism $\pi : X' \rightarrow X$ such that X' is a complex manifold, $\pi^{-1}(Y) = D$ is a normal crossing divisor in X' , and π induces an isomorphism $X' \setminus D \xrightarrow{\sim} Z \setminus Z \cap Y$ (hence $d_{X'} = d_Z$). We set $\mathcal{M}' = ({}_{\mathrm{D}}\pi^* \mathcal{M})(*D)[d_Z - d_X]$. By [14, §11.2.2], \mathcal{M}' underlies an object of $\mathrm{MTM}^{\mathrm{reg}}(X')$, in particular it is in degree zero, and is therefore of D-type. Moreover, by Lemma 4.2, (c) holds for \mathcal{M}' .

On the other hand, the adjunction isomorphism of Corollary 1.8 induces an isomorphism

$$\mathrm{Hom}_{\mathcal{D}_{X \times S/S}}(\mathcal{M}', \mathcal{M}') \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{D}_{Y \times S/S}}({}_{\mathrm{D}}\pi_* \mathcal{M}', \mathcal{M}),$$

and $\mathrm{Id}_{\mathcal{M}'}$ provides the adjunction morphism ${}_{\mathrm{D}}\pi_* \mathcal{M}' \rightarrow \mathcal{M}$. Note that ${}_{\mathrm{D}}\pi_* \mathcal{M}' = ({}_{\mathrm{D}}\pi_* \mathcal{M}')(*Y)$, so finally ${}_{\mathrm{D}}\pi_* \mathcal{M}' \simeq \mathcal{M}$. We end the proof that (c)_n holds for \mathcal{M} by considering the isomorphisms

$$\mathcal{M} \simeq {}_{\mathrm{D}}\pi_* \mathcal{M}' \stackrel{4.2}{\simeq} {}_{\mathrm{D}}\pi_* \mathrm{RH}^S(\mathrm{PSol} \mathcal{M}') \stackrel{3.24}{\simeq} \mathrm{RH}^S(R\pi_* \mathrm{PSol} \mathcal{M}') \stackrel{1.17(c)}{\simeq} \mathrm{RH}^S(\mathrm{PSol} \mathcal{M}).$$

q.e.d.

Proof of the functoriality in Theorem 5. Let $\varphi : \mathcal{M} \rightarrow \mathcal{N}$ be a morphism in the category $\mathrm{Mod}_{\mathrm{rhol}}(\mathcal{D}_{X \times S/S})$ and assume that \mathcal{M}, \mathcal{N} underlie objects of $\mathrm{MTM}^{\mathrm{reg}}(X)$, so that (4.4) is an isomorphism, according to the proof above. By the bi-functoriality of (4.4), we have a commutative diagram

$$\begin{array}{ccc} \mathrm{Hom}_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathrm{RH}^S(\mathrm{PSol} \mathcal{M})) & \xrightarrow{\sim} & \mathrm{Hom}_{p^{-1}\mathcal{O}_S}(\mathrm{Sol} \mathcal{M}, \mathrm{Sol} \mathcal{M}) \\ \downarrow (\mathrm{RH}^S \mathrm{PSol} \varphi) \circ & & \downarrow \circ (\mathrm{Sol} \varphi) \\ \mathrm{Hom}_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathrm{RH}^S(\mathrm{PSol} \mathcal{N})) & \xrightarrow{\sim} & \mathrm{Hom}_{p^{-1}\mathcal{O}_S}(\mathrm{Sol} \mathcal{N}, \mathrm{Sol} \mathcal{M}) \\ \circ \varphi \uparrow & & \uparrow (\mathrm{Sol} \varphi) \circ \\ \mathrm{Hom}_{\mathcal{D}_{X \times S/S}}(\mathcal{N}, \mathrm{RH}^S(\mathrm{PSol} \mathcal{N})) & \xrightarrow{\sim} & \mathrm{Hom}_{p^{-1}\mathcal{O}_S}(\mathrm{Sol} \mathcal{N}, \mathrm{Sol} \mathcal{N}) \end{array}$$

If $\eta_{\mathcal{M}} : \mathcal{M} \xrightarrow{\sim} \mathrm{RH}^S(\mathrm{PSol} \mathcal{M})$ corresponds to $\mathrm{Id}_{\mathrm{Sol} \mathcal{M}}$ via the horizontal isomorphism, and similarly for $\eta_{\mathcal{N}}$, then the diagram shows that $(\mathrm{RH}^S \mathrm{PSol} \varphi) \circ \eta_{\mathcal{M}} = \eta_{\mathcal{N}} \circ \varphi$, since both correspond to $\mathrm{Sol} \varphi$ via the middle horizontal isomorphism. q.e.d.

Appendix. Locally constant sheaves of $p^{-1}\mathcal{O}_S$ -modules

In this appendix, S denotes a complex analytic space which is not necessarily reduced, S_{red} denotes the associated reduced space and \mathcal{O}_S denotes its structure sheaf (a sheaf of rings on S_{red}). When there is no risk of confusion, we will use the notation S instead of S_{red} as the underlying space. We state the results we need without proofs, which are straightforward.

An S -constant sheaf of $p^{-1}\mathcal{O}_S$ -modules on $X \times S$ is a sheaf of the form $p^{-1}G$ for some sheaf G of \mathcal{O}_S -modules. We say that F is $p^{-1}\mathcal{O}_S$ -coherent if G is \mathcal{O}_S -coherent. Similarly, there is the notion of S -constant sheaf of \mathbb{C} -vector spaces.

Proposition A.1 (S -constant sheaves). *Let X be a topological space. An S -constant sheaf F of $p^{-1}\mathcal{O}_S$ -modules on $X \times S$ satisfies the following properties.*

- (1) *If $f : Y \rightarrow X$ is a continuous map, then $f^{-1}F$ is S -constant on Y .*

(2) If $U \subset X$ is a connected open set in X and $x \in U$, the natural morphism $p_{U,*}F \rightarrow i_x^{-1}F$ is an isomorphism of \mathcal{O}_S -modules. Conversely, if X is a connected topological space and F is a sheaf of $p^{-1}\mathcal{O}_S$ -modules on $X \times S$ such that the natural morphism $p_*F \rightarrow i_x^{-1}F$ is an isomorphism of \mathcal{O}_S -modules for each $x \in X$, then F is S -constant. In particular, if X is a connected topological space and G is a sheaf of \mathcal{O}_S -modules, the sheaf $p_*p^{-1}G$ is naturally identified to G .

(3) If G, G' are \mathcal{O}_S -modules, there are canonical isomorphisms:

$$p^{-1}(G \oplus G') \simeq p^{-1}G \oplus p^{-1}G', \quad p^{-1}(G \otimes_{\mathcal{O}_S} G') \simeq p^{-1}G \otimes_{p^{-1}\mathcal{O}_S} p^{-1}G',$$

and if moreover G is \mathcal{O}_S -coherent or X is locally connected

$$p^{-1}\mathcal{H}om_{\mathcal{O}_S}(G, G') \simeq \mathcal{H}om_{p^{-1}\mathcal{O}_S}(p^{-1}G, p^{-1}G').$$

(4) The functor p^{-1} , from the category of \mathcal{O}_S -modules to that of $p^{-1}\mathcal{O}_S$ -modules is exact. Moreover, if X is connected, this functor is fully faithful.

(5) If X is a connected topological space, the kernel, the image and the cokernel of a morphism between S -constant sheaves of $p^{-1}\mathcal{O}_S$ -modules are S -constant sheaves of $p^{-1}\mathcal{O}_S$ -modules.

We say that a sheaf F on $X \times S$ of $p^{-1}\mathcal{O}_S$ -modules is S -locally constant if each point $(x, s) \in X \times S$ has a neighbourhood on which F is S -constant. We then say that F is $p^{-1}\mathcal{O}_S$ -coherent if it is locally (on $X \times S$) isomorphic to the pull-back by p of a \mathcal{O}_S -coherent sheaf.

Proposition A.2. *If X is connected and locally connected, and if F is S -locally constant on $X \times S$, then there exists a sheaf G of \mathcal{O}_S -modules such that, locally on $X \times S$, we have $F \simeq p^{-1}G$. We can choose for G any of the sheaves $i_x^{-1}F$ for $x \in X$. Moreover, F is $p^{-1}\mathcal{O}_S$ -coherent if and only if G is \mathcal{O}_S -coherent.*

In other words, the isomorphisms are locally defined, but the sheaf G exists globally on S . However, this sheaf is not unique.

Proposition A.3. *Assume that X is locally connected. Let F be a sheaf of $p^{-1}\mathcal{O}_S$ -modules on $X \times S$. Then F is an S -locally constant sheaf of $p^{-1}\mathcal{O}_S$ -modules if and only if it is S -locally constant as a sheaf of \mathbb{C} -vector spaces.*

Proposition A.4. *If F, F' are S -locally constant on $X \times S$, and $\varphi : F \rightarrow F'$ is $p^{-1}\mathcal{O}_S$ -linear, then $F \oplus F'$, $F \otimes_{p^{-1}\mathcal{O}_S} F'$, $\mathcal{H}om_{p^{-1}\mathcal{O}_S}(F, F')$, $\ker \varphi$, $\operatorname{im} \varphi$ and $\operatorname{coker} \varphi$ are also S -locally constant. If F and F' are moreover $p^{-1}\mathcal{O}_S$ -coherent, so are these sheaves.*

Corollary A.5. *The category of S -locally constant sheaves of $p^{-1}\mathcal{O}_S$ -modules (resp. and $p^{-1}\mathcal{O}_S$ -coherent) is a full abelian subcategory of the category of sheaves of $p^{-1}\mathcal{O}_S$ -modules.*

Corollary A.6. *Let $0 \rightarrow F' \rightarrow F \rightarrow F'' \rightarrow 0$ be an exact sequence of sheaves of $p^{-1}\mathcal{O}_S$ -modules. If F, F' (resp. F, F'') are S -locally constant (resp. and coherent), then so are F' (resp. F'').*

Proposition A.7. *Set $I = [0, 1]$. Let F be an S -locally constant sheaf of $p^{-1}\mathcal{O}_S$ -modules on $X \times S$, with $X = I$ or $X = I \times I$. Then F is S -constant.*

Let $\gamma : I \rightarrow X$ be a continuous map, with $\gamma(0) = x_o$, $\gamma(1) = x_1$. If F is S -locally constant on $X \times S$, then so is $\gamma^{-1}F$ on $I \times S$, hence it is S -constant, and it defines an isomorphism $T_\gamma : i_{x_o}^{-1}F \xrightarrow{\sim} i_{x_1}^{-1}F$ of \mathcal{O}_S -modules.

Proposition A.8. *If γ and γ' are homotopic with fixed endpoints, then $T_\gamma = T_{\gamma'}$. If γ and γ' can be composed, we have $T_{\gamma \cdot \gamma'} = T_\gamma \circ T_{\gamma'}$.*

Let us now assume that X is connected and locally path-connected, and let us fix a base point $x_o \in X$. We consider the category $\text{Rep}_{\mathcal{O}_S}(\pi_1(X, x_o))$ whose objects are representations $\rho : \pi_1(X, x_o) \rightarrow \text{Aut}_{\mathcal{O}_S}(G)$ for some sheaf G of \mathcal{O}_S -modules and the morphisms $\rho \rightarrow \rho'$ are \mathcal{O}_S -linear morphisms $\varphi : G \rightarrow G'$ which satisfy $\rho'(\gamma) \circ \varphi = \varphi \circ \rho(\gamma)$ for any $\gamma \in \pi_1(X, x_o)$.

Given an S -locally constant sheaf F on $X \times S$, Proposition A.8 shows that $\gamma \mapsto T_\gamma$ defines a representation $\rho : \pi_1(X, x_o) \rightarrow \text{Aut}_{\mathcal{O}_S}(i_{x_o}^{-1}F)$, called the monodromy representation attached to F . A morphism of S -locally constant sheaves obviously gives rise to a morphism of their monodromy representation. We thus get a functor from the category of S -constant local systems of $p^{-1}\mathcal{O}_S$ -modules to $\text{Rep}_{\mathcal{O}_S}(\pi_1(X, x_o))$.

Proposition A.9. *The monodromy representation functor is an equivalence of categories.*

Remark A.10 (Riemann-Hilbert). By the Riemann-Hilbert correspondence for coherent S -local systems proved in [2, Th. 2.23 p. 14], the functor $F \mapsto \mathcal{O}_{X \times S} \otimes_{p^{-1}\mathcal{O}_S} F$ induces an equivalence between the category of coherent S -locally constant sheaves of $p^{-1}\mathcal{O}_S$ -modules and the category of coherent $\mathcal{O}_{X \times S}$ -modules F equipped with an integrable relative connection $\nabla : F \rightarrow \Omega_{X \times S/S}^1 \otimes_{\mathcal{O}_{X \times S}} F$.

Proposition A.11. *Let F be a coherent S -locally constant local system on $X \times S$. Then the following properties are equivalent:*

- (1) *there exists an \mathcal{O}_S -locally free sheaf of finite rank G such that locally $F \simeq p^{-1}G$;*
- (2) *any coherent \mathcal{O}_S -module G such that locally $F \simeq p^{-1}G$ is \mathcal{O}_S -locally free of finite rank.*

If S is a complex manifold with its reduced structure, (1) and (2) are also equivalent to

- (3) *the dual $\mathbf{D}F := \mathbf{R}\mathcal{H}om_{p^{-1}\mathcal{O}_S}(F, p^{-1}\mathcal{O}_S)$ is an S -locally constant sheaf.*

If X is connected and locally path-connected, and if $\pi_1(X, x_o)$ has finite presentation, so that $\text{Hom}(\pi_1(X, x_o), \text{GL}_r(\mathbb{C}))$ is naturally an affine complex algebraic variety, then (1) and (2) are also equivalent to

- (4) *for any open subset V of S on which some G as in (2) is free or rank r , giving $F|_{X \times V}$ is equivalent to giving a holomorphic map $V \rightarrow \text{Hom}(\pi_1(X, x_o), \text{GL}_r(\mathbb{C}))$.*

Proposition A.12. *Let Y be a contractible topological space and let F be an S -locally constant sheaf on $Y \times S$. Then $R^k p_{Y,*}F = 0$ for each $k \geq 1$ and F is constant.*

References

- [1] A. D’AGNOLO, S. GUILLERMOU & P. SCHAPIRA – “Regular holonomic $\mathcal{D}[[\hbar]]$ -modules”, *Publ. RIMS, Kyoto Univ.* **47** (2011), no. 1, p. 221–255.
- [2] P. DELIGNE – *Équations différentielles à points singuliers réguliers*, Lect. Notes in Math., vol. 163, Springer-Verlag, 1970.
- [3] M.J. EDMUNDO & L. PRELLI – “Sheaves on \mathcal{T} -topologies”, *J. Math. Soc. Japan* **68** (2016), no. 1, p. 347–381.
- [4] R. Hotta, K. TAKEUCHI & T. TANISAKI – *D-Modules, perverse sheaves, and representation theory*, Progress in Math., vol. 236, Birkhäuser, Boston, Basel, Berlin, 2008, in Japanese: 1995.
- [5] M. KASHIWARA – “On the holonomic systems of differential equations II”, *Invent. Math.* **49** (1978), p. 121–135.
- [6] ———, “The Riemann-Hilbert problem for holonomic systems”, *Publ. RIMS, Kyoto Univ.* **20** (1984), p. 319–365.
- [7] ———, *D-modules and microlocal calculus*, Translations of Mathematical Monographs, vol. 217, American Mathematical Society, Providence, R.I., 2003.
- [8] M. KASHIWARA & T. KAWAI – “On the holonomic systems of differential equations (systems with regular singularities) III”, *Publ. RIMS, Kyoto Univ.* **17** (1981), p. 813–979.
- [9] M. KASHIWARA & P. SCHAPIRA – *Sheaves on Manifolds*, Grundlehren Math. Wiss., vol. 292, Springer-Verlag, 1990.
- [10] ———, *Moderate and formal cohomology associated with constructible sheaves*, Mém. Soc. Math. France (N.S.), vol. 64, Société Mathématique de France, Paris, 1996.
- [11] ———, *Ind-sheaves*, Astérisque, vol. 271, Société Mathématique de France, Paris, 2001.
- [12] ———, *Categories and sheaves*, Grundlehren Math. Wiss., vol. 332, Springer-Verlag, 2006.
- [13] T. MOCHIZUKI – *Asymptotic behaviour of tame harmonic bundles and an application to pure twistor D-modules*, vol. 185, Mem. Amer. Math. Soc., no. 869-870, American Mathematical Society, Providence, R.I., 2007.
- [14] ———, *Mixed twistor D-Modules*, Lect. Notes in Math., vol. 2125, Springer-Verlag, 2015.
- [15] T. MONTEIRO FERNANDES & L. PRELLI – “Relative subanalytic sheaves”, *Fund. Math.* **226** (2014), no. 1, p. 79–100.
- [16] T. MONTEIRO FERNANDES & C. SABBABH – “On the de Rham complex of mixed twistor \mathcal{D} -modules”, *Internat. Math. Res. Notices* (2013), no. 21, p. 4961–4984.
- [17] L. PRELLI – “Sheaves on subanalytic sites”, *Rend. Sem. Mat. Univ. Padova* **120** (2008), p. 167–216.
- [18] C. SABBABH – “Proximité évanescence, I. La structure polaire d’un \mathcal{D} -module, Appendice en collaboration avec F. Castro”, *Compositio Math.* **62** (1987), p. 283–328.
- [19] ———, *Polarizable twistor \mathcal{D} -modules*, Astérisque, vol. 300, Société Mathématique de France, Paris, 2005.
- [20] P. SCHAPIRA & J.-P. SCHNEIDERS – *Index theorem for elliptic pairs*, Astérisque, vol. 224, Société Mathématique de France, Paris, 1994.
- [21] L. WANG – “The constructibility theorem for differential modules”, Ph.D. Thesis, University of Illinois at Chicago, 2008, <http://indigo.uic.edu/handle/10027/13547>.

T. MONTEIRO FERNANDES, Centro de Matemática e Aplicações Fundamentais – Centro de investigação Operacional e Departamento de Matemática da FCUL, Edifício C 6, Piso 2, Campo Grande, 1700, Lisboa, Portugal • *E-mail* : mtfernandes@fc.ul.pt

C. SABBABH, CMLS, École polytechnique, CNRS, Université Paris-Saclay, F-91128 Palaiseau cedex, France • *E-mail* : Claude.Sabbah@polytechnique.edu
Url : <http://www.math.polytechnique.fr/perso/sabbah>