

DYNAMICS OF VORTICES FOR THE COMPLEX GINZBURG-LANDAU EQUATION

EVELYNE MIOT

ABSTRACT. We study a complex Ginzburg-Landau equation in the plane, which has the form of a Gross-Pitaevskii equation with some dissipation added. We focus on the regime corresponding to well-prepared unitary vortices and derive their asymptotic motion law.

2000 Mathematics Subject Classification: 35B20,35B40,35Q40,82D55.
Keywords: Complex Ginzburg-Landau equation, Vortex dynamics.

1. INTRODUCTION

In this paper, we study the dynamics of vortices for a complex Ginzburg-Landau equation on the plane, namely

$$\frac{\delta}{|\log \varepsilon|} \partial_t u_\varepsilon + \alpha i \partial_t u_\varepsilon = \Delta u_\varepsilon + \frac{1}{\varepsilon^2} u_\varepsilon (1 - |u_\varepsilon|^2) \quad (\text{CGL})_\varepsilon$$

where $u_\varepsilon : \mathbb{R}_+ \times \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a complex valued map. Here δ , α and ε denote positive real parameters, and we will mainly focus on the asymptotics as ε tends to zero while δ and α are kept fixed. Up to a change of scale, we may further assume that $\alpha = 1$, and we set $k_\varepsilon = \frac{\delta}{|\log \varepsilon|}$. The complex Ginzburg-Landau equation $(\text{CGL})_\varepsilon$ reduces to the Gross-Pitaevskii equation when $\delta = 0$ and to the parabolic Ginzburg-Landau equation when $\alpha = 0$. Both the Gross-Pitaevskii and the Ginzburg-Landau equations have been widely investigated in the regime which we will consider (see e.g. [8, 18, 15, 4] for the Gross-Pitaevskii equation and [12, 19, 6] and references therein for the parabolic Ginzburg-Landau equation). Typical functions u_ε in this regime are given explicitly by

$$u_\varepsilon^*(a_i, d_i)(z) := \prod_{i=1}^l u_{\varepsilon, d_i}(z - a_i) = \prod_{i=1}^l f_{1, d_i} \left(\frac{|z - a_i|}{\varepsilon} \right) \left(\frac{z - a_i}{|z - a_i|} \right)^{d_i},$$

where the points $a_i \in \mathbb{R}^2$, $d_i = \pm 1$, and the functions $f_{1, d_i} : \mathbb{R}^+ \mapsto [0, 1]$ which satisfy $f_{1, d_i}(0) = 0$, $f_{1, d_i}(+\infty) = 1$ are in some sense optimal profiles. The points a_i are called the vortices of the fields u_ε^* and the d_i their degrees. This class of functions u_ε^* is of course not invariant by any of the flows corresponding to these equations, but not far from it¹, and it is in particular possible to define notions of point vortices for solutions of $(\text{CGL})_\varepsilon$, at least in an asymptotic way as $\varepsilon \rightarrow 0$, and to study their dynamics. This dynamics is eventually governed by a system of ordinary differential equations, at least before collisions.

Date: September 29, 2008.

¹see the notion of well-preparedness in Section 1.2.

Two relevant quantities in the study of vortex dynamics are the Ginzburg-Landau energy

$$E_\varepsilon(u) = \int_{\mathbb{R}^2} e_\varepsilon(u) dx = \int_{\mathbb{R}^2} \frac{|\nabla u|^2}{2} + \frac{(1 - |u|^2)^2}{4\varepsilon^2} dx,$$

through its energy density $e_\varepsilon(u)$, and the Jacobian

$$Ju = \frac{1}{2} \operatorname{curl}(u \times \nabla u)$$

through its primitive $j(u) = u \times \nabla u$. In the regime which we will consider, one has

$$\frac{e_\varepsilon(u_\varepsilon)}{|\log \varepsilon|} dx \rightarrow \pi \sum_{i=1}^l \delta_{a_i} \quad \text{and} \quad Ju_\varepsilon dx \rightarrow \pi \sum_{i=1}^l d_i \delta_{a_i}$$

as $\varepsilon \rightarrow 0$, which describes asymptotically the positions and the degrees of the vortices. The quantity $e_\varepsilon(u_\varepsilon)$ was especially used in the study of the parabolic Ginzburg-Landau equation while $j(u_\varepsilon)$ was used in the study of the Gross-Pitaevskii equation. Here, we will rely on both of them.

In the case of the domain being the entire plane \mathbb{R}^2 , which we consider here, the reference fields $u_\varepsilon^*(a_i, d_i)$ have infinite Ginzburg-Landau energy E_ε whenever $d = \sum d_i \neq 0$. In [7], a notion of renormalized energy² for such data was introduced in order to solve the Cauchy problem for the Gross-Pitaevskii equation. This notion was later used in [4] in order to study the dynamics of vortices for the Gross-Pitaevskii equation in the plane. Our definition of well-prepared data below and part of the subsequent analysis is borrowed from [4].

The complex Ginzburg-Landau equation $(\text{CGL})_\varepsilon$, either in the plane or in the real line, has been vastly considered in the literature, especially as a model for amplitude oscillation in weakly nonlinear systems undergoing a Hopf bifurcation (see e.g. [2] for a survey paper). The mathematical analysis of vortices for $(\text{CGL})_\varepsilon$ has been initiated in [18], where it was presented as an alternative approach (a regularized version) for the study of the Gross-Pitaevskii equation. We believe however that the conclusion regarding the dynamics of vortices for $(\text{CGL})_\varepsilon$ in [18] is erroneous, and that Theorem 2 yields the corrected version.

After the completion of this work we were informed that Kurzke, Melcher, Moser and Spirn [16] independently obtained similar results concerning the dynamics of vortices for $(\text{CGL})_\varepsilon$ in bounded and simply connected domains.

1.1. Renormalized energy and Cauchy Problem. As already mentioned, for $d = \sum d_i \neq 0$ the Ginzburg-Landau energy of $u_\varepsilon^*(a_i, d_i)$ is infinite. It can actually be computed that

$$\int_{\mathbb{R}^2} \frac{|\nabla |u_\varepsilon^*(a_i, d_i)||^2}{2} + \frac{(1 - |u_\varepsilon^*(a_i, d_i)|^2)^2}{4\varepsilon^2} dz < +\infty,$$

whereas as $|z| \rightarrow +\infty$,

$$|\nabla |u_\varepsilon^*(a_i, d_i)||^2(z) \sim \frac{d^2}{|z|^2},$$

²not to be merged with the notion in [3].

so that

$$\int_{\mathbb{R}^2} \frac{|\nabla u_\varepsilon^*(a_i, d_i)|^2}{2} = +\infty.$$

The renormalized energy introduced in [7] is obtained by subtracting the diverging part of the gradient at infinity. More precisely, given a smooth map U_d such that

$$U_d = \left(\frac{z}{|z|} \right)^d \quad \text{on } \mathbb{R}^2 \setminus B(0, 1),$$

we have as $|z| \rightarrow +\infty$

$$|\nabla u_\varepsilon^*(a_i, d_i)|^2 \sim |\nabla U_d|^2$$

and one may define

$$\mathcal{E}_{\varepsilon, U_d}(u_\varepsilon^*(a_i, d_i)) := \lim_{R \rightarrow +\infty} \int_{B(R)} \left[e_\varepsilon(u_\varepsilon^*(a_i, d_i)) - \frac{|\nabla U_d|^2}{2} \right] < +\infty. \quad (1)$$

This definition extends to a larger class of functions, and is a useful ingredient in solving the Cauchy problem. Following [7], we define

$$\mathcal{V} = \{U \in L^\infty(\mathbb{R}^2, \mathbb{C}), \nabla^k U \in L^2, \forall k \geq 2, (1 - |U|^2) \in L^2, \nabla|U| \in L^2\}.$$

In particular, the space \mathcal{V} contains all the maps u_ε^* as well as the reference maps U_d . Our first result, which we prove in the Appendix, establishes global well-posedness in the class³ $\mathcal{V} + H^1(\mathbb{R}^2)$.

Theorem 1. *Let $u_0 = U + w_0$ be in $\mathcal{V} + H^1(\mathbb{R}^2)$. Then there exists a unique global solution u to $(\text{CGL})_\varepsilon$ such that $u \in C^0(\{U\} + H^1(\mathbb{R}^2))$. If we write $u(t) = U + w(t)$, then w is the unique solution in $C^0(\mathbb{R}_+, H^1(\mathbb{R}^2))$ to*

$$\begin{cases} (k_\varepsilon + i)\partial_t w = \Delta w + f_U(w) \\ w(0) = w_0, \end{cases} \quad (2)$$

where

$$f_U(w) = \Delta U + \frac{1}{\varepsilon^2}(U + w)(1 - |U + w|^2).$$

In addition, w satisfies

$$w \in L_{\text{loc}}^1(\mathbb{R}_+, H^2(\mathbb{R}^2)) \cap L_{\text{loc}}^\infty(\mathbb{R}_+, L^\infty(\mathbb{R}^2)), \partial_t w \in L_{\text{loc}}^1(\mathbb{R}_+, L^2(\mathbb{R}^2))$$

and

$$w \in C^\infty(\mathbb{R}_+, C^\infty(\mathbb{R}^2)).$$

Finally, the functional $E_{\varepsilon, U}(u) := E_{\varepsilon, U}(w)$ defined by

$$E_{\varepsilon, U}(u) = \int_{\mathbb{R}^2} \frac{|\nabla w|^2}{2} - \int_{\mathbb{R}^2} \Delta U \cdot w + \int_{\mathbb{R}^2} \frac{(1 - |U + w|^2)^2}{4\varepsilon^2}$$

satisfies

$$\frac{d}{dt} E_{\varepsilon, U}(u) = -k_\varepsilon \int_{\mathbb{R}^2} |\partial_t w|^2 dx, \quad \forall t \geq 0.$$

³In [10], the Cauchy problem in local spaces is investigated for a more general class of complex Ginzburg-Landau equations.

As a matter of fact, it follows from an integration by part that if $u \in \{U\} + H^1(\mathbb{R}^2)$ is as in Theorem 1 and if U satisfies in addition $|\nabla U(x)| \leq \frac{C}{\sqrt{|x|}}$, then

$$E_{\varepsilon,U}(u(t)) \equiv \mathcal{E}_{\varepsilon,U}(u(t)) = \lim_{R \rightarrow +\infty} \int_{B(R)} \left[e_{\varepsilon}(u(t)) - \frac{|\nabla U|^2}{2} \right] dx.$$

The functions $u_{\varepsilon}^*(a_i, d_i)$ are not H^1 perturbations one of the other, even for fixed $d = \sum d_i$, unless some algebraic relations involving the a_i 's and d_i 's hold. In order to handle a class of functions containing them all, it is useful to introduce the following equivalence relation on the set \mathcal{V} :

$$\begin{aligned} \forall U, U' \in \mathcal{V}, \quad U \sim U' \text{ iff} \\ \deg_{\infty}(U) = \deg_{\infty}(U') \text{ and } |\nabla U|^2 - |\nabla U'|^2 \in L^1(\mathbb{R}^2). \end{aligned}$$

Denoting by $[U]$ the corresponding equivalence class of U , we observe that for any configuration (a_i, d_i) such that $\sum d_i = d$, we have $u_{\varepsilon}^*(a_i, d_i) \in [U_d]$. Therefore the space $[U_d] + H^1(\mathbb{R}^2)$ contains in particular all H^1 perturbations of all reference maps u_{ε}^* of degree d at infinity.

For a map u in $[U_d] + H^1(\mathbb{R}^2)$, we may now define

$$\mathcal{E}_{\varepsilon,[U_d]}(u) := \lim_{R \rightarrow +\infty} \int_{B(R)} \left[e_{\varepsilon}(u) - \frac{|\nabla U_d|^2}{2} \right],$$

which is a finite quantity. Moreover, for any solution $u \in C^0(\{U\} + H^1(\mathbb{R}^2))$ with $U \in [U_d]$, we infer from Theorem 1 that

$$\frac{d}{dt} \mathcal{E}_{\varepsilon,[U_d]}(u) = \frac{d}{dt} \mathcal{E}_{\varepsilon,U}(u) = -k_{\varepsilon} \int_{\mathbb{R}^2} |\partial_t u|^2,$$

which means that the renormalized energy has the same dissipation rate as the Ginzburg-Landau energy in the finite energy case $d = 0$.

1.2. Statement of the result. In the sequel, A_n denotes the annulus $B(2^{n+1}) \setminus B(2^n)$ for $n \in \mathbb{N}$, so that $\mathbb{R}^2 = B(2^{n_0}) \cup (\cup_{n \geq n_0} A_n)$.

Definition 1. Let a_1, \dots, a_l be l distinct points in \mathbb{R}^2 , $d_i \in \{-1, +1\}$ for $i = 1, \dots, l$ and set $d = \sum d_i$. Let $(u_{\varepsilon})_{0 < \varepsilon < 1}$ be a family of maps in $[U_d] + H^1(\mathbb{R}^2)$. We say that $(u_{\varepsilon})_{0 < \varepsilon < 1}$ is well-prepared with respect to the configuration (a_i, d_i) if there exist $R = 2^{n_0} > \max |a_i|$ and a constant $K_0 > 0$ such that⁴

$$\lim_{\varepsilon \rightarrow 0} \|Ju_{\varepsilon} - \pi \sum_{i=1}^l d_i \delta_{a_i}\|_{W_0^{1,\infty}(B(R))^*} = 0, \quad (\text{WP}_1)$$

$$\sup_{0 < \varepsilon < 1} E_{\varepsilon}(u_{\varepsilon}, A_n) \leq K_0 \quad \forall n \geq n_0, \quad (\text{WP}_2)$$

and

$$\lim_{\varepsilon \rightarrow 0} \left(\mathcal{E}_{\varepsilon,[U_d]}(u_{\varepsilon}) - \mathcal{E}_{\varepsilon,[U_d]}(u_{\varepsilon}^*(a_i, d_i)) \right) = 0. \quad (\text{WP}_3)$$

We can now state our main theorem as follows

⁴Here, $E_{\varepsilon}(u, B) \equiv \int_B e_{\varepsilon}(u)$.

Theorem 2. Let $(u_\varepsilon^0)_{0 < \varepsilon < 1}$ in $[U_d] + H^1(\mathbb{R}^2)$ be a family of well-prepared initial data with respect to the configuration (a_i^0, d_i) with $d_i = \pm 1$, and let $(u_\varepsilon(t))_{0 < \varepsilon < 1}$ in $C(\mathbb{R}_+, [U_d] + H^1(\mathbb{R}^2))$ be the corresponding solution of $(\text{CGL})_\varepsilon$. Let $\{a_i(t)\}_{\{i=1, \dots, l\}}$ denote the solution of the ordinary differential equation

$$\begin{cases} \pi \dot{a}_i(t) = C_i(\delta d_i \mathbb{I}_2 - \mathbb{J}_2) \nabla_{a_i} W, & C_i = \frac{-d_i}{1 + \delta^2} \\ a_i(0) = a_i, & i = 1, \dots, l \end{cases} \quad (3)$$

where

$$\mathbb{I}_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \mathbb{J}_2 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

and W is the Kirchhoff-Onsager functional defined by

$$W(a_i, d_i) = -\pi \sum_{i \neq j} d_i d_j \log |a_i - a_j|.$$

We denote by $[0, T^*)$ its maximal interval of existence. Then, for every $t \in [0, T^*)$, the family $(u_\varepsilon(t))_{0 < \varepsilon < 1}$ is well-prepared with respect to the configuration $(a_i(t), d_i)$.

2. EVOLUTION FORMULA FOR u_ε

In this section, we recall or derive a number of evolution formulae involving quantities related to u_ε which we introduce now.

2.1. Notations. Given $x = (x_1, x_2) \in \mathbb{R}^2$, we set $x^\perp = \mathbb{J}x = (-x_2, x_1)$, which with the standard identification of \mathbb{R}^2 and \mathbb{C} reads $x^\perp = ix$. For z and $z' \in \mathbb{C}$, $z \cdot z' = \text{Re}(z\bar{z}')$ denotes the scalar product and $z \times z' = z^\perp \cdot z' = -\text{Im}(z\bar{z}')$ the exterior product of z and z' in \mathbb{R}^2 . For $\vec{a} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$, we define $\text{curl}(\vec{a}) = \partial_1 a_2 - \partial_2 a_1$. If $u : \mathbb{R}^2 \rightarrow \mathbb{C}$, we denote by

$$j(u) = u \times \nabla u = iu \cdot \nabla u = u^\perp \cdot \nabla u$$

the linear momentum and

$$J(u) = \partial_1 u \times \partial_2 u = \det(\nabla u)$$

the Jacobian of u . For $u \in H_{\text{loc}}^1(\mathbb{R}^2)$, it can be checked that $J(u) = \frac{1}{2} \text{curl} j(u)$ in the distribution sense. On the set where u does not vanish, we have for $k = 1, 2$

$$\partial_k u = \partial_k u \cdot \frac{u}{|u|} \frac{u}{|u|} + \partial_k u \cdot \frac{i u}{|u|} \frac{i u}{|u|}.$$

This yields

$$\partial_k u = \partial_k |u| \frac{u}{|u|} + \frac{j_k(u)}{|u|} \frac{u^\perp}{|u|}, \quad (2.1)$$

hence we have

$$\partial_k u \cdot \partial_l u = \partial_k |u| \partial_l |u| + \frac{j_k(u) j_l(u)}{|u|^2} \quad (2.2)$$

and it follows that

$$|\nabla u|^2 = |\nabla|u||^2 + \frac{|j(u)|^2}{|u|^2}. \quad (2.3)$$

The Hopf differential of u is defined as

$$\omega(u) = |\partial_1 u|^2 - |\partial_2 u|^2 - 2i\partial_1 u \cdot \partial_2 u = 4\partial_z u \overline{\partial_{\bar{z}} u}.$$

It follows from (2.2) that $\omega(u)$ may be rewritten in terms of the components of $\nabla|u|$ and $j(u)$ as

$$\begin{aligned} \omega(u) &= \partial_1|u|^2 - \partial_2|u|^2 - 2i\partial_1|u|\partial_2|u| \\ &\quad + \frac{1}{|u|^2}(j_1^2(u) - j_2^2(u) - 2ij_1(u)j_2(u)). \end{aligned} \quad (2.4)$$

We recall that the Ginzburg-Landau energy density is defined by

$$e_\varepsilon(u) = \frac{|\nabla u|^2}{2} + \frac{(1 - |u|^2)^2}{4\varepsilon^2} = \frac{|\nabla u|^2}{2} + V(u),$$

and we set

$$\mu_\varepsilon(u) = \frac{e_\varepsilon(u)}{|\log \varepsilon|}.$$

In view of (2.3), we then have

$$e_\varepsilon(u) = e_\varepsilon(|u|) + \frac{|j(u)|^2}{|u|^2}. \quad (2.5)$$

Finally, we write the right-hand side in $(\text{CGL})_\varepsilon$ as

$$\nabla E(u) = \nabla E_\varepsilon(u) = \Delta u + \frac{1}{\varepsilon^2}u(1 - |u|^2).$$

2.2. Evolution formulae involving the Jacobian and the energy density. For a smooth map u in space-time, direct computations by integration by part yield for the energy

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}^2} e_\varepsilon(u) \varphi \, dx &= - \int_{\mathbb{R}^2} \partial_t u \cdot \nabla E(u) \varphi \, dx \\ &\quad - \int_{\mathbb{R}^2} \nabla \varphi \cdot (\partial_t u \cdot \nabla u) \, dx \end{aligned} \quad (2.6)$$

and for the Jacobian

$$\frac{d}{dt} \int_{\mathbb{R}^2} J(u) \chi \, dx = - \int_{\mathbb{R}^2} \nabla^\perp \chi \cdot (\partial_t u^\perp \cdot \nabla u) \, dx, \quad (2.7)$$

where $\chi, \varphi \in \mathcal{D}(\mathbb{R}^2)$.

On the other hand, the Pohozaev identity (see e.g. [5]) yields for any vector field $\vec{X} \in \mathcal{D}(\mathbb{R}^2, \mathbb{C})$

$$\int_{\mathbb{R}^2} \vec{X} \cdot (\nabla E(u) \cdot \nabla u) \, dx = - \int_{\mathbb{R}^2} \text{Re} \left(\omega(u) \frac{\partial \vec{X}}{\partial \bar{z}} \right) \, dz + \int_{\mathbb{R}^2} V(u) \nabla \cdot \vec{X} \, dx.$$

In particular, the choice of $\vec{X} = \nabla\varphi$, for which $\partial_{\bar{z}}X = 2\frac{\partial^2\varphi}{\partial\bar{z}^2}$, or $\vec{X} = \nabla^\perp\chi$, for which $\partial_{\bar{z}}X = 2i\frac{\partial^2\chi}{\partial\bar{z}^2}$, leads to

$$\int_{\mathbb{R}^2} \nabla\varphi \cdot (\nabla E(u) \cdot \nabla u) dx = -2 \int_{\mathbb{R}^2} \operatorname{Re}\left(\omega(u) \frac{\partial^2\varphi}{\partial\bar{z}^2}\right) dz + \int_{\mathbb{R}^2} V(u) \Delta\varphi dx$$

and

$$\int_{\mathbb{R}^2} \nabla^\perp\chi \cdot (\nabla E(u) \cdot \nabla u) dx = 2 \int_{\mathbb{R}^2} \operatorname{Im}\left(\omega(u) \frac{\partial^2\chi}{\partial\bar{z}^2}\right) dz. \quad (2.8)$$

We next consider a solution u of $(\text{CGL})_\varepsilon$, which is smooth in view of Theorem 1. In this case, $\nabla E(u)$ and $\partial_t u$ are related by

$$\partial_t u = \frac{1}{\alpha_\varepsilon} \nabla E(u) = \beta_\varepsilon \nabla E(u), \quad (2.9)$$

where $\alpha_\varepsilon = \frac{\delta}{|\log \varepsilon|} + i = k_\varepsilon + i$. Using (2.9) in (2.6) and (2.7), we obtain

$$\frac{d}{dt} \int_{\mathbb{R}^2} e_\varepsilon(u) \varphi dx = -k_\varepsilon \int_{\mathbb{R}^2} |\partial_t u|^2 \varphi dx - \int_{\mathbb{R}^2} \nabla\varphi \cdot (\beta_\varepsilon \nabla E(u) \cdot \nabla u) dx$$

and

$$\frac{d}{dt} \int_{\mathbb{R}^2} J(u) \chi dx = - \int_{\mathbb{R}^2} \nabla^\perp\chi \cdot (i\beta_\varepsilon \nabla E(u) \cdot \nabla u) dx.$$

In order to get rid of the terms of the form $\int_{\mathbb{R}^2} \vec{X} \cdot (i\nabla E(u) \cdot \nabla u)$, we compute

$$\frac{d}{dt} \int_{\mathbb{R}^2} [bJ(u)\chi - ae_\varepsilon(u)\varphi]$$

where $\beta_\varepsilon = a + ib$. This yields

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}^2} [bJ(u)\chi - ae_\varepsilon(u)\varphi] &= (b^2 + a^2) \int_{\mathbb{R}^2} \nabla^\perp\chi \cdot (\nabla E \cdot \nabla u) + ak_\varepsilon \int_{\mathbb{R}^2} |\partial_t u|^2 dx \\ &\quad + \int_{\mathbb{R}^2} (\nabla\varphi - \nabla^\perp\chi) \cdot (a(a + ib)\nabla E \cdot \nabla u). \end{aligned}$$

Since $a = \frac{k_\varepsilon}{k_\varepsilon^2 + 1}$ and $b = \frac{-1}{k_\varepsilon^2 + 1}$, we finally infer from the above relation and (2.8) the following

Proposition 1. *Let u solve $(\text{CGL})_\varepsilon$. Then for all $\varphi, \chi \in \mathcal{D}(\mathbb{R}^2)$,*

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}^2} [J(u)\chi + k_\varepsilon e_\varepsilon(u)\varphi] &= -k_\varepsilon^2 \int_{\mathbb{R}^2} |\partial_t u|^2 \varphi - 2 \int_{\mathbb{R}^2} \operatorname{Im}\left(\omega(u) \frac{\partial^2\chi}{\partial\bar{z}^2}\right) \\ &\quad + R_\varepsilon(t, \varphi, \chi, u), \end{aligned}$$

where the remainder R_ε is defined by

$$R_\varepsilon(t, \varphi, \chi, u) = -k_\varepsilon \int_{\mathbb{R}^2} (\nabla\varphi - \nabla^\perp\chi) \cdot (\beta_\varepsilon \nabla E(u) \cdot \nabla u)$$

or equivalently

$$R_\varepsilon(t, \varphi, \chi, u) = -k_\varepsilon \int_{\mathbb{R}^2} (\nabla\varphi - \nabla^\perp\chi) \cdot (\partial_t u \cdot \nabla u).$$

Proposition 1 allows to derive formally the motion law for the vortices. Indeed, assume that we have

$$Ju_\varepsilon(t) \rightarrow \pi \sum_{i=1}^l d_i \delta_{a_i(t)},$$

$$\mu_\varepsilon(u_\varepsilon)(t) \rightarrow \pi \sum_{i=1}^l \delta_{a_i(t)}$$

and $u_\varepsilon(t)$ is close in some sense to $u_\varepsilon^*(a_i(t), d_i)$, and therefore to $u^*(a_i(t), d_i)$, where

$$u^*(a_i, d_i) = \prod_{i=1}^l \left(\frac{z - a_i}{|z - a_i|} \right)^{d_i}.$$

We use Proposition 1 with u formally replaced by $u^*(a_i(t), d_i)$ and with choices of test functions φ and χ which are localized and affine near each point $a_i(t)$ and satisfy $\nabla\varphi = \nabla^\perp\chi$ there, so that both terms $k_\varepsilon^2 \int_{\mathbb{R}^2} |\partial_t u|^2 \varphi$ and $R_\varepsilon(t, \varphi, \chi, u_\varepsilon)$ vanish in the limit $\varepsilon \rightarrow 0$. Using the formula (see (7.2) in [5])

$$2 \int_{\mathbb{R}^2} \operatorname{Im} \left(\omega(u^*(a_i, d_i)) \frac{\partial^2 \chi}{\partial \bar{z}^2} \right) = 2\pi \sum_{j \neq i} d_i d_j \frac{(a_i - a_j)^\perp}{|a_i - a_j|^2} \cdot \nabla \chi(a_i),$$

we then obtain that for each i

$$\pi d_i \dot{a}_i(t) \cdot \nabla \chi(a_i) + \delta \pi \dot{a}_i(t) \cdot \nabla \varphi(a_i) = -2\pi \sum_{j \neq i} d_i d_j \frac{(a_i - a_j)^\perp}{|a_i - a_j|^2} \cdot \nabla \chi(a_i).$$

Taking into account the fact that $\nabla \varphi(a_i) = \nabla^\perp \chi(a_i)$, we infer that

$$\pi (d_i \dot{a}_i(t) - \delta \dot{a}_i^\perp(t)) \cdot \nabla \chi(a_i) = -2\pi \sum_{j \neq i} d_i d_j \frac{(a_i - a_j)^\perp}{|a_i - a_j|^2} \cdot \nabla \chi(a_i),$$

which yields the ODE (3).

In Sections 4 and 5, in order to give a rigorous meaning to the previous computations, we will prove the convergence of the Jacobians and of the energy densities to the weighted sums of dirac masses mentioned above, and then show that both the energy dissipation $k_\varepsilon^2 \int_{\mathbb{R}^2} |\partial_t u_\varepsilon|^2$ and the remainder $R_\varepsilon(t, \varphi, \chi, u_\varepsilon)$ vanish when ε tends to zero. In Section 6, we will establish some asymptotic control of $\omega(u_\varepsilon) - \omega(u^*(a_i), d_i)$ away from the vortices in terms of the excess energy $\mathcal{E}_{\varepsilon, [U_d]}(u_\varepsilon) - \mathcal{E}_{\varepsilon, [U_d]}(u_\varepsilon^*(a_i))$, and finally prove that this excess energy converges to zero by mean of a Gronwall inequality.

3. SOME RESULTS ON THE RENORMALIZED ENERGY

3.1. Degree and energy at infinity. In this paragraph, we collect some results from [4] related to the energy at infinity, which require the notion of degree at infinity.

Let A be the annulus $B(2) \setminus B(1)$. We define

$$T_d = \{u \in H^1(A) \text{ s.t. } \exists B \subset B(u), |B| \geq \frac{3}{4}, \forall r \in B, \deg(u, \partial B(r)) = d\},$$

where $B(u)$ is the set of radii $r \in [1, 2]$ such that the restriction $u|_{\partial B(r)}$ is continuous and does not vanish, and we define the sublevel sets

$$E_\varepsilon^\Lambda = \{u \in H^1(A) \text{ s.t. } E_\varepsilon(u, A) < \Lambda\}.$$

The topological sector of degree d is then defined as

$$S_{d,\varepsilon}^\Lambda = E_\varepsilon^\Lambda \cap T_d.$$

The following Theorem is proved in [1].

Theorem 3 ([1]). *For all $\Lambda > 0$, there exists $\varepsilon_\Lambda > 0$ such that for every $0 < \varepsilon < \varepsilon_\Lambda$, we have*

$$E_\varepsilon^\Lambda = \bigcup_{d \in \mathbb{Z}} S_{d,\varepsilon}^\Lambda.$$

The map $\text{deg} : E_\varepsilon^\Lambda \rightarrow \mathbb{Z}$, $u \in S_{d,\varepsilon}^\Lambda \mapsto d$ is continuous.

In the sequel of this section, we fix $\Lambda > \Lambda_d = 2\pi d^2 \log 2$ and we set

$$S_d \equiv S_{d,\varepsilon_\Lambda}^\Lambda,$$

so that in particular the map U_d belongs to S_d , since $|U_d| \equiv 1$ on A and $\int_A \frac{|\nabla U_d|^2}{2} = \pi d^2 \log 2$.

One may easily infer from Theorem 3 that if $u \in [U_d] + H^1(\mathbb{R}^2)$, then for any sufficiently large k the map $u(2^k \cdot)$ belongs to some $S_{d(k)}$. In fact, one can find a radius from which $d(k) \equiv d$, i.e., u has well defined and constant degree d at infinity.

Proposition 2 ([4]). *Let $d \in \mathbb{Z}$, $\Lambda > \Lambda_d$ and $u \in [U_d] + H^1(\mathbb{R}^2)$. There exists an integer $n \in \mathbb{N}^*$ such that for any $k \geq n$, the map $u_k : z \in A \mapsto u(2^k z)$ belongs to the topological sector S_d . We denote by $n(u)$ the smallest integer for which this property holds.*

For maps $u \in [U_d] + H^1(\mathbb{R}^2)$ satisfying in addition a uniform energy bound on large annuli one can characterize $n(u)$ as follows (see e.g. the proof of Lemma 7.1 in [4]).

Lemma 1. *Let $\Lambda > \Lambda_d$ be given and $0 < \varepsilon < \varepsilon_\Lambda$. Let $u \in [U_d] + H^1(\mathbb{R}^2)$ and assume that there exists some $n_0 \in \mathbb{N}^*$ such that for all $n \geq n_0$,*

$$E_\varepsilon(u, A_n) < \Lambda.$$

Then we have $n(u) \leq n_0$.

The next Lemma provides a lower bound for the energy on large annuli.

Lemma 2 ([4]). *Let $d \in \mathbb{Z}$ and $u \in [U_d] + H^1(\mathbb{R}^2)$. Then, for any $k \geq n(u)$, we have for $0 < \varepsilon < \varepsilon_\Lambda$*

$$\int_{A_k} [e_\varepsilon(u) - \frac{|\nabla U_d|^2}{2}] \geq -C2^{-2k} \varepsilon^2.$$

One can then derive from Lemma 2 an upper bound for $E_\varepsilon(u, B) - E_\varepsilon(u_\varepsilon^*, B)$ on large balls B in terms of the excess energy $\mathcal{E}_{\varepsilon, [U_d]}(u) - \mathcal{E}_{\varepsilon, [U_d]}(u_\varepsilon^*)$. We will therefore be able to rely on properties of the Ginzburg-Landau energy on bounded domains in the course of the proof of Theorem 2.

Lemma 3 ([4]). *Let $d \in \mathbb{Z}$, $u \in [U_d] + H^1(\mathbb{R}^2)$, $a_1, \dots, a_l \in \mathbb{R}^2$ and $d_1, \dots, d_l \in \mathbb{Z}^*$ such that $d = \sum d_i$. Let $k \geq 1 + \max\{\log_2 |a_1|, \dots, \log_2 |a_l|, n(u)\}$ and $R = 2^k$. Then, we have*

$$\int_{B(R)} [e_\varepsilon(u) - e_\varepsilon(u_\varepsilon^*(a_i, d_i))] \leq \mathcal{E}_{\varepsilon, [U_d]}(u) - \mathcal{E}_{\varepsilon, [U_d]}(u_\varepsilon^*(a_i, d_i)) + \frac{C}{R},$$

where C depends only on l and d .

3.2. Explicit identities for the reference map u_ε^* . We present here an account of some classical identities for the energy of u_ε^* , which are borrowed from [4].

In the sequel, we consider a configuration (a_i, d_i) with $d_i \in \mathbb{Z}^*$ and we set $d = \sum d_i$. We begin with an explicit expansion near each vortex a_j .

Lemma 4. *For $j \in \{1, \dots, l\}$ and $0 < \varepsilon < 1$,*

$$\int_{B(a_j, r)} e_\varepsilon(u_\varepsilon^*(a_i, d_i)) = \pi d_j^2 \log\left(\frac{r}{\varepsilon}\right) + \gamma(|d_j|) + O\left(\frac{r}{r_a}\right)^2 + O\left(\frac{\varepsilon}{r}\right)^2,$$

where $\gamma(|d_j|)$ is some universal constant.

On the other hand, $u_\varepsilon^*(a_i, d_i)$ behaves as $u^*(a_i, d_i)$ away from the vortices, so its energy on $\Omega_{R,r} = B(R) \setminus \cup_{j=1}^l B(a_j, r)$ is close to the energy of $u^*(a_i, d_i)$ on $\Omega_{R,r}$ which we can compute explicitly (see [3]). Combining the previous expansions, we obtain

Proposition 3. *Let*

$$r_a = \frac{1}{8} \min_{i \neq j} \{|a_i - a_j|\}, \quad R_a = \max\{|a_i|\}.$$

Then for $R > R_a + 1$, we have as $\varepsilon \rightarrow 0$

$$\begin{aligned} \int_{B(R)} e_\varepsilon(u_\varepsilon^*(a_i, d_i)) &= \pi \sum_{i=1}^l d_i^2 |\log \varepsilon| + W(a_i, d_i) + \sum_{i=1}^l \gamma(|d_i|) \\ &\quad + \pi d^2 \log R + O\left(\frac{R_a}{R}\right) + o_\varepsilon(1). \end{aligned}$$

We observe that $\pi d^2 \log R = \int_{B(R) \setminus B(1)} \frac{|\nabla U_d|^2}{2}$. This yields the following expansion for the renormalized energy

Corollary 1. *When $\varepsilon \rightarrow 0$, the following holds*

$$\begin{aligned} \mathcal{E}_{\varepsilon, [U_d]}(u_\varepsilon^*(a_i, d_i)) &= \pi \sum_{i=1}^l d_i^2 |\log \varepsilon| + W(a_i, d_i) \\ &\quad + \sum_{i=1}^l \gamma(|d_i|) - \int_{B(1)} \frac{|\nabla U_d|^2}{2} + o_\varepsilon(1). \end{aligned}$$

Concerning the energy on annuli, we finally quote the following

Lemma 5. *For $R > R_a$, we have*

$$\int_{B(2R) \setminus B(R)} e_\varepsilon(u_\varepsilon^*(a_i, d_i)) = \pi d^2 \log 2 + O\left(\frac{R_a}{R}\right)$$

or, in view of the properties of U_d at infinity,

$$\int_{B(2R) \setminus B(R)} e_\varepsilon(u_\varepsilon^*(a_i, d_i)) = \int_{B(2R) \setminus B(R)} \frac{|\nabla U_d|^2}{2} + O\left(\frac{R_a}{R}\right).$$

4. COERCIVITY FOR THE RENORMALIZED ENERGY

In this section, we supplement some results from [4] and [14] with estimates which we will later need. These results establish precise estimates in various norms for maps u being close to $u_\varepsilon^*(a_i, d_i)$ in terms of the excess energy with respect to the configuration (a_i, d_i) . For a map $u \in [U_d] + H^1(\mathbb{R}^2)$ and a given configuration (a_i, d_i) with $d_i = \pm 1$, we define this excess energy Σ_ε as

$$\Sigma_\varepsilon = \Sigma_\varepsilon(a_i, d_i) = \mathcal{E}_{\varepsilon, [U_d]}(u) - \mathcal{E}_{\varepsilon, [U_d]}(u_\varepsilon^*(a_i, d_i)).$$

We also set

$$r_a = \frac{1}{8} \min_{i \neq j} \{|a_i - a_j|\}, \quad R_a = \max_{i=1, \dots, l} \{|a_i|\}.$$

Theorem 4. *Let $r \leq r_a$ and $2^{n_0} = R_0 > R_a$ such that $\cup_{i=1}^l B(a_i, r) \subset B(R_0)$. Then there exist ε_0 and η_0 depending only on l, r, r_a, R_a, R_0 satisfying the following property. For all $u \in [U_d] + H^1(\mathbb{R}^2)$ such that*

$$\eta = \|Ju - \pi \sum_{i=1}^l d_i \delta_{a_i}\|_{W_0^{1, \infty}(B(R_0))^*} \leq \eta_0 \quad (4.1)$$

and

$$2^{n(u)} \leq R_0, \quad (4.2)$$

then for $\varepsilon \leq \varepsilon_0$ we have

$$\int_{B(R_0) \setminus \cup B(a_i, r)} e_\varepsilon(|u|) + \frac{1}{8} \left| \frac{j(u)}{|u|} - j(u^*(a_i, d_i)) \right|^2 \leq \Sigma_\varepsilon + C(\eta, \varepsilon, \frac{1}{R_0}), \quad (4.3)$$

where C is a continuous function on \mathbb{R}^3 vanishing at the origin. Furthermore, there exist points $b_i \in B(a_i, r/2)$ such that

$$\|Ju - \pi \sum_{i=1}^l d_i \delta_{b_i}\|_{W_0^{1, \infty}(B(R_0))^*} \leq f(R_0, \Sigma_\varepsilon) \varepsilon |\log \varepsilon| \quad (4.4)$$

and

$$\|\mu_\varepsilon(u) - \pi \sum_{i=1}^l \delta_{b_i}\|_{W_0^{1, \infty}(B(R_0))^*} \leq \frac{g(R_0, r, r_a, \Sigma_\varepsilon)}{|\log \varepsilon|}, \quad (4.5)$$

where f and g are continuous functions on \mathbb{R}^2 and \mathbb{R}^4 .

Proof. Except for the energy concentration (4.5), each of the other statements is already proved in Theorem 6.1 of [4]. We first infer from (4.1) that for all i $\|Ju - \pi d_i \delta_{a_i}\|_{W_0^{1, \infty}(B(a_i, r))^*} \leq \eta_0$. If η_0 is sufficiently small with respect to r this gives in view of Theorem 3 in [14] $K_0^i \geq C(r)$, where K_0^i is the local excess energy near the vortex i defined by $K_0^i = \int_{B(a_i, r)} e_\varepsilon(u) - \pi \log(\frac{r}{\varepsilon})$. It follows that

$$\int_{B(a_i, r)} e_\varepsilon(u) \leq \int_{B(R_0)} e_\varepsilon(u) - \pi(l-1)|\log \varepsilon| - C(r).$$

On the other hand, since $n(u) \leq n_0$, we have according to Lemma 3 and Proposition 3

$$\int_{B(R_0)} e_\varepsilon(u) \leq \int_{B(R_0)} e_\varepsilon(u_\varepsilon^*(a_i, d_i)) + \Sigma_\varepsilon + \frac{C}{R_0} \leq \pi l |\log \varepsilon| + \Sigma_\varepsilon + C.$$

This first implies that $K_0^i \leq C + \Sigma_\varepsilon$. Also, replacing r by $3r/4$ we see that $\int_{B(R_0) \setminus \cup B(a_i, 3r/4)} \mu_\varepsilon(u) \leq (C + \Sigma_\varepsilon) |\log \varepsilon|^{-1}$, where C only depends on R_0, r, r_a, R_a .

Now, according to Theorem 2' in [14], the energy density $\mu_\varepsilon(u)$ on $B(a_i, r)$ concentrates at the point $b_i \in B(a_i, r/2)$ where $J(u)$ concentrates. From Theorem 3.2.1 in [9] and the estimate for K_0^i it follows that

$$\|\mu_\varepsilon(u) - \pi \delta_{b_i}\|_{W_0^{1,\infty}(B(a_i, r))^*} \leq \frac{f(\Sigma_\varepsilon, C)}{|\log \varepsilon|}.$$

Combining the above and the upper bound for the energy density outside the vortex balls finally yields (4.5). \square

5. CONVERGENCE TO LIPSCHITZ VORTEX PATHS

In this section, we establish compactness for the Jacobians and the energy densities in a more general situation, replacing assumption (WP₃) in Theorem 2 by a uniform bound on the initial excess energy.

Theorem 5. *Let (a_i^0, d_i) with $d_i = \pm 1$ be a configuration of vortices. Let $R = 2^{n_0}$ and $(u_\varepsilon^0)_{0 < \varepsilon < 1}$ in $[U_d] + H^1(\mathbb{R}^2)$ such that*

$$\lim_{\varepsilon \rightarrow 0} \|Ju_\varepsilon^0 - \pi \sum_{i=1}^l d_i \delta_{a_i^0}\|_{W_0^{1,\infty}(B(R))^*} = 0, \quad (\text{WP}_1)$$

$$\sup_{0 < \varepsilon < 1} E_\varepsilon(u_\varepsilon^0, A_n) \leq K_0, \quad \forall n \geq n_0, \quad (\text{WP}_2)$$

and

$$\sup_{0 < \varepsilon < 1} \left(\mathcal{E}_{\varepsilon, [U_d]}(u_\varepsilon^0) - \mathcal{E}_{\varepsilon, [U_d]}(u_\varepsilon^*(a_i^0, d_i)) \right) \leq K_1. \quad (\text{WP}_{3'})$$

Then there exist $R' = 2^{n_1}$ and $T > 0$ depending only on K_1, R, r_{a^0} and R_{a^0} , a sequence $\varepsilon_k \rightarrow 0$ and l Lipschitz paths $b_i : [0, T] \rightarrow \mathbb{R}^2$ starting from a_i^0 such that

$$\sup_{t \in [0, T]} \|Ju_{\varepsilon_k}(t) - \pi \sum_{i=1}^l d_i \delta_{b_i(t)}\|_{W_0^{1,\infty}(B(R'))^*} \rightarrow 0, \quad k \rightarrow +\infty \quad (5.1)$$

and

$$\sup_{t \in [0, T]} \|\mu_{\varepsilon_k}(u_{\varepsilon_k})(t) - \pi \sum_{i=1}^l \delta_{b_i(t)}\|_{W^{1,\infty}(B(R'))^*} \rightarrow 0, \quad k \rightarrow +\infty. \quad (5.2)$$

Moreover, there exist a constant $C_0 > 0$ depending only on r_{a^0}, R, K_1 and K_0 and a constant $C_1 > 0$ depending on r_{a^0}, R and K_1 such that for all $t \in [0, T]$ and for $k \in \mathbb{N}$,

$$E_{\varepsilon_k}(u_{\varepsilon_k}(t), A_n) \leq C_0, \quad \forall n \geq n_1 \quad (5.3)$$

and

$$\mathcal{E}_{\varepsilon_k, [U_d]}(u_{\varepsilon_k}(t)) - \mathcal{E}_{\varepsilon_k, [U_d]}(u_{\varepsilon_k}^*(b_i(t), d_i)) \leq C_1. \quad (5.4)$$

Proof. The proof is very similar to that of Theorem 4 in [4]. In the sequel, C will stand for a constant depending only on r_{a^0}, R, R_{a^0} and K_1 . In order to simplify the notations we will further set $r_a = r_{a^0}$ and $R_a = R_{a^0}$.

We first consider $\Lambda > \max(K_0, \Lambda_d)$. Thanks to Lemma 1 and (WP₂), there exists $\varepsilon_\Lambda > 0$ such that for all $0 < \varepsilon < \varepsilon_\Lambda$, we have $n(u_\varepsilon^0) \leq n_0$. We fix such a Λ and from now on only consider $0 < \varepsilon < \varepsilon_\Lambda$.

We next introduce the smallest integer $n_1 \geq n_0$ such that $2^{n_1} \geq \max(R, R_a + r_a)$ and define $R' = 2^{n_1}$. In the remainder of the proof, we will write $\|\cdot\|$ instead of $\|\cdot\|_{W_0^{1,\infty}(B(R'))^*}$. Our aim is to apply Theorem 4 to each $u_\varepsilon(t)$ for the choice $r = r_a$ and $R_0 = R'$. Let η_0 and ε_0 be the constants provided by Theorem 4 for this choice. First, thanks to (WP₂) and (WP_{3'}) it turns out that the convergence in (WP₁) still holds on the larger ball $B(R')$ (see the proof of Lemma 7.3 in [4]). Therefore, since $t \mapsto Ju_\varepsilon(t) \in L^1(B(R'))$ is continuous for each ε , there exists a time $T_\varepsilon > 0$ such that

$$\|Ju_\varepsilon(s) - \pi \sum_{i=1}^l d_i \delta_{a_i^0}\| < \eta_0, \quad \forall s \in [0, T_\varepsilon].$$

We take T_ε to be the maximum time smaller than T^* having this property, where T^* is defined in Theorem 2.

On the other hand, since $t \mapsto E_\varepsilon(u_\varepsilon(t), A_n)$ is continuous uniformly with respect to n and $\Lambda > K_0$, we infer from (WP₂) that there exists $T'_\varepsilon > 0$ such that for $s \in [0, T'_\varepsilon]$

$$E_\varepsilon(u_\varepsilon(s), A_n) < \Lambda, \quad \forall n \geq n_1,$$

so according to Lemma 1 we have $n(u_\varepsilon(s)) \leq n_1$ for $s \in [0, T'_\varepsilon]$. We take $T'_\varepsilon \leq T^*$ maximal with this property.

We claim that there exists a constant D depending on K_1, r_a, R and K_0 such that for all $s \in [0, \min(T_\varepsilon, T'_\varepsilon)]$,

$$E_\varepsilon(u_\varepsilon(s), A_n) \leq D, \quad \forall n \geq n_1. \quad (5.5)$$

Consequently, if we assume from the beginning that

$$\Lambda > \max(K_0, \Lambda_d, D),$$

then $T'_\varepsilon \geq T_\varepsilon$, and it follows from Lemma 1 that $n(u_\varepsilon(s)) \leq n_1$ on $[0, T_\varepsilon]$.

Proof of (5.5). As in [4], we decompose for each $n \geq n_1$ $E_\varepsilon(u_\varepsilon(t), A_n) - E_\varepsilon(u_\varepsilon^*(a_i^0, d_i), A_n)$ as

$$\begin{aligned} & \sum_{\substack{k=n_1 \\ k \neq n}}^{+\infty} (E_\varepsilon(u_\varepsilon^*(a_i^0, d_i), A_k) - E_\varepsilon(u_\varepsilon(t), A_k)) \\ & + E_\varepsilon(u_\varepsilon^*(a_i^0, d_i), B(R')) - E_\varepsilon(u_\varepsilon(s), B(R')) \\ & + \mathcal{E}_{\varepsilon, [U_d]}(u_\varepsilon(s)) - \mathcal{E}_{\varepsilon, [U_d]}(u_\varepsilon^*(a_i^0, d_i)). \end{aligned}$$

We first handle each term of the sum in the right-hand side. In view of Lemmas 2 and 5, we have for $k \geq n_1$

$$\begin{aligned} E_\varepsilon(u_\varepsilon(t), A_k) &\geq -C\varepsilon^2 2^{-2k} + \int_{A_k} \frac{|\nabla U_d|^2}{2} \\ &\geq E_\varepsilon(u_\varepsilon^*(a_i^0, d_i), A_k) - C(R_a)2^{-k} - C\varepsilon^2 2^{-2k}, \end{aligned}$$

so we deduce that

$$\sum_{\substack{k=n_1 \\ k \neq n}}^{+\infty} (E_\varepsilon(u_\varepsilon^*(a_i^0, d_i), A_k) - E_\varepsilon(u_\varepsilon(t), A_k)) \leq C.$$

Next, we infer from the definition of T_ε and Theorem 3 in [14] that $\int_{B(a_i^0, r_a)} e_\varepsilon(u_\varepsilon(s)) \geq \pi |\log \varepsilon| - C$. Observe that R' is chosen so that $\cup B(a_i^0, r_a) \subset B(R')$, so this leads to

$$E_\varepsilon(u_\varepsilon(s), B(R')) \geq \pi l |\log \varepsilon| - C.$$

Using Proposition 3, we thus find

$$E_\varepsilon(u_\varepsilon^*(a_i^0, d_i), B(R')) - E_\varepsilon(u_\varepsilon(s), B(R')) \leq C. \quad (5.6)$$

Finally, we define $\Sigma_\varepsilon^0(s) := \mathcal{E}_{\varepsilon, [U_d]}(u_\varepsilon(s)) - \mathcal{E}_{\varepsilon, [U_d]}(u_\varepsilon^*(a_i^0, d_i))$. Since $\mathcal{E}_{\varepsilon, [U_d]}(u_\varepsilon(t))$ is non-increasing, we obtain in view of (WP_{3'})

$$\Sigma_\varepsilon^0(s) \leq \mathcal{E}_{\varepsilon, [U_d]}(u_\varepsilon^0) - \mathcal{E}_{\varepsilon, [U_d]}(u_\varepsilon^*(a_i^0, d_i)) \leq K_1,$$

and (5.5) follows.

We may now apply Theorem 4 to each $u_\varepsilon(t)$ on $[0, T_\varepsilon]$. This provides points $b_i^\varepsilon(s) \in B(a_i^0, \frac{r_a}{2})$ for $0 \leq s \leq T_\varepsilon$. Since $\Sigma_\varepsilon^0(s) \leq K_1$, estimate (4.3) turns into

$$\int_{\Omega_{R', r_a}} e_\varepsilon(|u_\varepsilon(s)|) + \frac{1}{8} \left| \frac{j(u_\varepsilon(s))}{|u_\varepsilon(s)|} - j(u_\varepsilon^*(a_i^0, d_i)) \right|^2 \leq C,$$

where $\Omega_{R', r_a} = B(R') \setminus \cup B(a_i^0, r_a)$. Also, we have by (2.4) and (2.5)

$$\int_{\Omega_{R', r_a}} e_\varepsilon(u_\varepsilon(s)) \leq C \quad (5.7)$$

and

$$\|\omega(u_\varepsilon(s))\|_{L^1(\Omega_{R', r_a})} \leq C, \quad (5.8)$$

where $C = C(R, r_a, K_1)$. For notation convenience, we may now write μ_ε instead of $\mu_\varepsilon(u_\varepsilon)$.

In the sequel, given any configuration (a_i, d_i) , we denote by $\mathcal{H}(a_i)$ the set of functions $\chi, \varphi \in \mathcal{D}(\mathbb{R}^2)$ such that

$$\chi = \sum_{i=1}^l \chi_i, \quad \varphi = \sum_{i=1}^l \varphi_i,$$

where for all i

$$\chi_i, \varphi_i \in \mathcal{D}\left(B\left(a_i, \frac{3r_a}{2}\right)\right), \quad \nabla \varphi_i = \nabla^\perp \chi_i \text{ on } B(a_i, r_a)$$

and χ_i (hence φ_i) is affine on $B(a_i, r_a)$ with $|\nabla\chi_i(a_i)| = |\nabla\varphi_i(a_i)| \leq 1$.

By definition of r_a such functions χ and φ always exist, and we can moreover estimate their L^∞ norms by

$$\|D\varphi\|_\infty, \|D\chi\|_\infty \leq \frac{C}{r_a}, \quad \|D^2\varphi\|_\infty, \|D^2\chi\|_\infty \leq \frac{C}{r_a^2}.$$

We next establish a control of the remainder terms appearing in Proposition 1.

Lemma 6. *There exists a constant $C = C(r_a, R, K_1, T^*)$ such that*

$$\int_0^{T_\varepsilon} \int_{\mathbb{R}^2} \frac{|\partial_t u_\varepsilon|^2}{|\log \varepsilon|^2} ds \leq \frac{C}{|\log \varepsilon|}$$

and for all $\chi, \varphi \in \mathcal{H}(a_i^0)$

$$\left| \int_0^{T_\varepsilon} \int_{\mathbb{R}^2} (\nabla^\perp \chi - \nabla \varphi) \cdot \frac{\partial_t u_\varepsilon \cdot \nabla u_\varepsilon}{|\log \varepsilon|} ds \right| \leq \frac{C}{|\log \varepsilon|^{\frac{1}{2}}}.$$

Proof. In order to prove the first inequality, we use Theorem 1 and obtain

$$\begin{aligned} \frac{\delta}{|\log \varepsilon|} \int_0^{T_\varepsilon} \int_{\mathbb{R}^2} |\partial_t u_\varepsilon|^2 &= \mathcal{E}_{\varepsilon, [U_d]}(u_\varepsilon^0) - \mathcal{E}_{\varepsilon, [U_d]}(u_\varepsilon(T_\varepsilon)) \\ &\leq K_1 + \mathcal{E}_{\varepsilon, [U_d]}(u_\varepsilon^*(a_i^0, d_i)) - \mathcal{E}_{\varepsilon, [U_d]}(u_\varepsilon(T_\varepsilon)). \end{aligned}$$

Since $n(u_\varepsilon(T_\varepsilon)) \leq n_1$ we have by Lemma 3

$$\begin{aligned} \mathcal{E}_{\varepsilon, [U_d]}(u_\varepsilon^*(a_i^0, d_i)) - \mathcal{E}_{\varepsilon, [U_d]}(u_\varepsilon(T_\varepsilon)) \\ \leq \int_{B(R')} e_\varepsilon(u_\varepsilon^*(a_i^0, d_i)) - \int_{B(R')} e_\varepsilon(u_\varepsilon(T_\varepsilon)) + \frac{C}{R'} \end{aligned}$$

which is bounded in view of (5.6). It then suffices to divide all terms by $|\log \varepsilon|$.

For the second assertion, we set $\xi = \nabla^\perp \chi - \nabla \varphi$ which has compact support in $A = \cup A_i$, where $A_i = B(a_i^0, \frac{3r_a}{2}) \setminus B(a_i^0, r_a)$, and we apply Cauchy-Schwarz inequality. We obtain

$$\begin{aligned} \left(\int_0^{T_\varepsilon} \int_{\mathbb{R}^2} (\nabla^\perp \chi - \nabla \varphi) \cdot \frac{\partial_t u_\varepsilon \cdot \nabla u_\varepsilon}{|\log \varepsilon|} \right)^2 \\ \leq \left(\int_0^{T_\varepsilon} \int_{\mathbb{R}^2} \frac{|\partial_t u_\varepsilon|^2}{|\log \varepsilon|^2} \right) \cdot \left(\int_0^{T_\varepsilon} \int_A |\nabla u_\varepsilon|^2 |\xi|^2 \right). \end{aligned}$$

Since $A \subset \Omega_{R', r_a}$ and $\sup_{0 < \varepsilon < 1} T_\varepsilon \leq T^*$, we infer from (5.7)

$$\int_0^{T_\varepsilon} \int_A |\nabla u_\varepsilon|^2 |\xi|^2 \leq \|\xi\|_\infty^2 \int_0^{T_\varepsilon} \int_A |\nabla u_\varepsilon|^2 \leq CT^* \|\xi\|_\infty^2,$$

and the conclusion finally follows from the first part of the proof. \square

We may now establish the following

Lemma 7. *There exists $T = T(r_a, R_a, R, K_1) > 0$ such that*

$$\liminf_{\varepsilon \rightarrow 0} T_\varepsilon \geq T.$$

Proof. The first step consists in showing that for $(\chi, \varphi) \in \mathcal{H}(a_i^0)$, for $s, t \in [0, T_\varepsilon]$ and $i = 1, \dots, l$ we have

$$\begin{aligned} & |\langle \chi_i, Ju_\varepsilon(t) - Ju_\varepsilon(s) \rangle + \delta \langle \varphi_i, \mu_\varepsilon(t) - \mu_\varepsilon(s) \rangle| \\ & \leq C|t - s| + \frac{C}{|\log \varepsilon|^{\frac{1}{2}}}. \end{aligned} \quad (5.9)$$

Indeed, we fix i and we invoke Proposition 1 for $u \equiv u_\varepsilon$ and the choice of test functions (χ_i, φ_i) . Integrating the formula given by Proposition 1 on $[s, t]$ yields

$$\begin{aligned} & |\langle \chi_i, Ju_\varepsilon(t) - Ju_\varepsilon(s) \rangle + \delta \langle \varphi_i, \mu_\varepsilon(t) - \mu_\varepsilon(s) \rangle| \leq 2 \int_s^t \int \left| \operatorname{Im} \left(\omega(u_\varepsilon) \frac{\partial^2 \chi_i}{\partial \bar{z}^2} \right) \right| \\ & \quad + \int_s^t \int \left| \frac{|\partial_t u_\varepsilon|^2}{|\log \varepsilon|^2} \varphi_i + (\nabla^\perp \chi_i - \nabla \varphi_i) \cdot \frac{\partial_t u_\varepsilon \cdot \nabla u_\varepsilon}{|\log \varepsilon|} \right|, \end{aligned}$$

where $\frac{\partial^2 \chi_i}{\partial \bar{z}^2}$ has support in $C_i \subset \Omega_{R', r_a}$, and it finally suffices to use (5.8) and Lemma 6.

In a second step, we take advantage of the equality

$$\|Ju_\varepsilon(T_\varepsilon) - \pi \sum_{i=1}^l d_i \delta_{a_i^0}\| \equiv \eta_0.$$

We set

$$\nu_{i,\varepsilon} = d_i \frac{b_i^\varepsilon(T_\varepsilon) - a_i^0}{|b_i^\varepsilon(T_\varepsilon) - a_i^0|}, \quad i = 1, \dots, l$$

and we define $\chi_{i,\varepsilon}, \varphi_{i,\varepsilon}$ so that for $x \in B(a_i^0, r_a)$,

$$\chi_{i,\varepsilon}(x) = \nu_{i,\varepsilon} \cdot x, \quad \varphi_{i,\varepsilon}(x) = \nu_{i,\varepsilon}^\perp \cdot x,$$

and we require additionally that $\chi = \sum \chi_{i,\varepsilon}$ and $\varphi = \sum \varphi_{i,\varepsilon}$ belong to $\mathcal{H}(a_i^0)$; we can moreover choose $\varphi_{i,\varepsilon}$ and $\chi_{i,\varepsilon}$ so that their norms in $C^2(B(R))$ remain bounded uniformly in ε . As $b_i^\varepsilon(T_\varepsilon) \in B(a_i^0, r_a/2)$, we have

$$|d_i ||b_i^\varepsilon(T_\varepsilon) - a_i^0| = d_i \chi(b_i^\varepsilon(T_\varepsilon) - a_i^0) + \delta \varphi(b_i^\varepsilon(T_\varepsilon) - a_i^0),$$

so that

$$\|\pi \sum_{i=1}^l d_i (\delta_{b_i^\varepsilon(T_\varepsilon)} - \delta_{a_i^0})\| = \langle \pi \sum_{i=1}^l d_i (\delta_{b_i^\varepsilon(T_\varepsilon)} - \delta_{a_i^0}), \chi \rangle + \delta \langle \pi \sum_{i=1}^l d_i (\delta_{b_i^\varepsilon(T_\varepsilon)} - \delta_{a_i^0}), \varphi \rangle.$$

On the other hand, we have

$$\|Ju_\varepsilon(T_\varepsilon) - \pi \sum_{i=1}^l d_i \delta_{a_i^0}\| \leq \|Ju_\varepsilon(T_\varepsilon) - \pi \sum_{i=1}^l d_i \delta_{b_i^\varepsilon(T_\varepsilon)}\| + \|\pi \sum_{i=1}^l d_i (\delta_{b_i^\varepsilon(T_\varepsilon)} - \delta_{a_i^0})\|.$$

The second term in the right-hand side may be rewritten as

$$\langle \pi \sum_{i=1}^l d_i (\delta_{b_i^\varepsilon(T_\varepsilon)} - \delta_{a_i^0}), \chi \rangle + \delta \langle \pi \sum_{i=1}^l d_i (\delta_{b_i^\varepsilon(T_\varepsilon)} - \delta_{a_i^0}), \varphi \rangle = A + B + C,$$

where

$$\begin{aligned} A &= \left\langle \pi \sum_{i=1}^l d_i \delta_{b_i^\varepsilon(T_\varepsilon)} - Ju_\varepsilon(T_\varepsilon), \chi \right\rangle + \delta \left\langle \pi \sum_{i=1}^l \delta_{b_i^\varepsilon(T_\varepsilon)} - \mu_\varepsilon(T_\varepsilon), \varphi \right\rangle \\ &\leq C \left(\|Ju_\varepsilon(T_\varepsilon) - \sum_{i=1}^l d_i \delta_{b_i^\varepsilon(T_\varepsilon)}\| + \delta \|\mu_\varepsilon(T_\varepsilon) - \sum_{i=1}^l \delta_{b_i^\varepsilon(T_\varepsilon)}\| \right), \end{aligned}$$

B is given by

$$B = \langle Ju_\varepsilon(T_\varepsilon) - Ju_\varepsilon(0), \chi \rangle + \delta \langle \mu_\varepsilon(T_\varepsilon) - \mu_\varepsilon(0), \varphi \rangle$$

and finally

$$\begin{aligned} C &= \left\langle Ju_\varepsilon^0 - \pi \sum_{i=1}^l d_i \delta_{a_i^0}, \chi \right\rangle + \delta \left\langle \mu_\varepsilon(u_\varepsilon^0) - \pi \sum_{i=1}^l \delta_{a_i^0}, \varphi \right\rangle \\ &\leq C \left(\|Ju_\varepsilon^0 - \sum_{i=1}^l d_i \delta_{a_i^0}\| + \delta \|\mu_\varepsilon(u_\varepsilon^0) - \sum_{i=1}^l \delta_{a_i^0}\| \right). \end{aligned}$$

In view of the bound provided by (5.9) for B , estimates (4.4)-(4.5) and the fact that $\Sigma_\varepsilon^0(s) \leq K_1$ for $0 \leq s \leq T_\varepsilon$, this implies

$$\eta_0 = \|Ju_\varepsilon(T_\varepsilon) - \pi \sum_{i=1}^l d_i \delta_{a_i^0}\| \leq C(\varepsilon |\log \varepsilon| + |\log \varepsilon|^{-1} + |\log \varepsilon|^{-\frac{1}{2}}) + CT_\varepsilon,$$

and letting $\varepsilon \rightarrow 0$ yields the conclusion. Lemma 7 is proved. \square

Proof of Theorem 5 completed.

We consider $t, s \in [0, T]$. Arguing as in the proof of Lemma 7 (with T_ε and 0 replaced by t and s), we find that for all χ, φ belonging to $\mathcal{H}(a_i^0)$

$$\begin{aligned} &\left| \sum_{i=1}^l d_i [\chi(b_i^\varepsilon(t)) - \chi(b_i^\varepsilon(s))] + \delta [\varphi(b_i^\varepsilon(t)) - \varphi(b_i^\varepsilon(s))] \right| \\ &\leq C \sup_{\tau \in [0, T]} \left(\|Ju_\varepsilon(\tau) - \sum_{i=1}^l d_i \delta_{b_i^\varepsilon(\tau)}\| + \delta \|\mu_\varepsilon(\tau) - \sum_{i=1}^l \delta_{b_i^\varepsilon(\tau)}\| \right) \\ &\quad + \left| \langle Ju_\varepsilon(t) - Ju_\varepsilon(s), \chi \rangle + \delta \langle \mu_\varepsilon(t) - \mu_\varepsilon(s), \varphi \rangle \right|, \end{aligned}$$

which is bounded by $o_\varepsilon(1) + c|t - s|$ by (4.4)-(4.5) and (5.9). Considering successively $\chi(x) = e_1 \cdot x$ and $\chi(x) = e_2 \cdot x$ on each $B(a_i^0, r_a)$, we obtain

$$|b_i^\varepsilon(t) - b_i^\varepsilon(s)| \leq c|t - s| + o_\varepsilon(1). \quad (5.10)$$

Next, using that $b_i^\varepsilon \in B(a_i^0, r_a)$ and a standard diagonal argument, we may construct a sequence $(\varepsilon_k) \rightarrow 0$ and paths $b_i(t)$ such that $b_i^{\varepsilon_k}(t)$ converges to $b_i(t)$ for all $t \in \mathbb{Q} \cap [0, T]$. We infer then from (4.4)-(4.5) that the convergence statements (5.1)-(5.2) in Theorem 5 hold for these times. Moreover, in view of (5.10) these paths are Lipschitz on $[0, T] \cap \mathbb{Q}$, so that they can be extended in a unique way to Lipschitz paths (still denoted by $b_i(t)$) on the whole of $[0, T]$. We can finally establish that the convergence (5.1)-(5.2) holds uniformly with respect to $t \in [0, T]$ by using again (5.10) and (4.4)-(4.5).

Finally, we already know from (5.5) that estimate (5.3) holds for the full family $(u_\varepsilon)_{\varepsilon < \varepsilon_\Lambda}$. In order to show (5.4), we recall first the uniform bound $\mathcal{E}_{\varepsilon, [U_d]}(u_\varepsilon(t)) - \mathcal{E}_{\varepsilon, [U_d]}(u_\varepsilon^*(a_i^0, d_i)) \leq K_1$. On the other hand, Corollary 1 gives $\mathcal{E}_{\varepsilon, [U_d]}(u_\varepsilon^*(a_i^0, d_i)) - \mathcal{E}_{\varepsilon, [U_d]}(u_\varepsilon^*(b_i(t), d_i)) = W(a_i^0, d_i) - W(b_i(t), d_i) \leq C$, since the b_i 's are continuous and remain separated on $[0, T]$. This yields the bound (5.4) and concludes the proof of Theorem 5. \square

As mentioned in the beginning of the proof of Theorem 5, the convergence of the initial data in (WP_1) actually holds on every large ball $B(L)$, $L = 2^n \geq R$, so that we find the same conclusions when replacing R by L .

Lemma 8 ([4], Lemma 7.3). *There exists a subsequence, still denoted by ε_k , such that for all $L \geq 2^{n_1}$,*

$$\eta_k := \sup_{[0, T]} \|Ju_{\varepsilon_k}(t) - \pi \sum_{i=1}^l d_i \delta_{b_i(t)}\|_{W_0^{1, \infty}(B(L))^*} \rightarrow 0, \quad k \rightarrow +\infty.$$

For $t \in [0, T]$ and sufficiently large $k \in \mathbb{N}$, we may therefore apply Theorem 4 to $u_{\varepsilon_k}(t)$ with respect to the configuration $(b_i(t), d_i)$ and with the choice $R_0 = L = 2^n$ for each $n \geq n_1$. We are led to introduce the excess energy at time t with respect to the configuration $(b_i(t), d_i)$ by

$$\Sigma_{\varepsilon_k}(t) = \mathcal{E}_{\varepsilon_k, U_d}(u_{\varepsilon_k}(t)) - \mathcal{E}_{\varepsilon_k, U_d}(u_{\varepsilon_k}^*(b_i(t), d_i)),$$

which is uniformly bounded on $[0, T]$ in view of (5.4). Letting first k , then n tend to $+\infty$, we can get rid of the dependance on R in (4.3).

Lemma 9. *For all $r \leq r_a/2$ and $K \geq 2^{n_1}$, we have for sufficiently large k and $t, t_1, t_2 \in [0, T]$*

$$\begin{aligned} \int_{B(K) \setminus \cup B(b_i(t), r)} e_{\varepsilon_k}(|u_{\varepsilon_k}(t)|) + \frac{1}{8} \left| \frac{j(u_{\varepsilon_k}(t))}{|u_{\varepsilon_k}(t)|} - j(u^*(b_i(t), d_i)) \right|^2 \\ \leq \Sigma_{\varepsilon_k}(t) + C(\varepsilon_k, \eta_k, \frac{1}{K}). \end{aligned}$$

Therefore, we have as $k \rightarrow +\infty$

$$\begin{aligned} \limsup_{k \rightarrow +\infty} \int_{t_1}^{t_2} \int_{B(K) \setminus \cup B(b_i(t), r)} e_{\varepsilon_k}(|u_{\varepsilon_k}(t)|) + \frac{1}{8} \left| \frac{j(u_{\varepsilon_k}(t))}{|u_{\varepsilon_k}(t)|} - j(u^*(b_i(t), d_i)) \right|^2 \\ \leq \limsup_{k \rightarrow +\infty} \int_{t_1}^{t_2} \Sigma_{\varepsilon_k}(t). \end{aligned}$$

Consequently, it appears that the distance between $u_{\varepsilon_k}(t)$ and $u^*(b_i(t), d_i)$ may be asymptotically entirely controlled by $\limsup \Sigma_{\varepsilon_k}(t)$.

We now define the trajectory set

$$\mathcal{T} = \{(t, b_i(t)), t \in [0, T], i = 1, \dots, l\}$$

and

$$\mathcal{G} = [0, T] \times \mathbb{R}^2 \setminus \mathcal{T}.$$

Thanks to the uniform bounds in $L_{\text{loc}}^2(\mathcal{G})$ provided by Lemma 9, we establish the following

Proposition 4. *There exists a subsequence, still denoted ε_k , such that*

$$\frac{j(u_{\varepsilon_k})}{|u_{\varepsilon_k}|} \rightharpoonup j(u^*(b_i(\cdot), d_i))$$

weakly in $L^2_{\text{loc}}(\mathcal{G})$ as $k \rightarrow +\infty$.

Proof. Let B be any bounded subset of \mathbb{R}^2 . First, we observe that according to Lemma 8

$$\text{curl}(j(u_{\varepsilon_k})) = 2Ju_{\varepsilon_k} \rightarrow 2\pi \sum_{i=1}^l d_i \delta_{b_i(\cdot)} = \text{curl}(j(u^*(b_i(\cdot), d_i))) \quad (5.11)$$

in $\mathcal{D}'([0, T] \times B)$.

On the other hand, we have

$$\text{div}(j(u_{\varepsilon_k})) \rightarrow 0 = \text{div}(j(u^*(b_i(\cdot), d_i))) \quad (5.12)$$

in $\mathcal{D}'([0, T] \times B)$.

Indeed, since u_{ε_k} solves $(\text{CGL})_{\varepsilon}$ we obtain by considering the exterior product

$$k_{\varepsilon_k} u_{\varepsilon_k} \times \partial_t u_{\varepsilon_k} + u_{\varepsilon_k} \cdot \partial_t u_{\varepsilon_k} = u_{\varepsilon_k} \times \Delta u_{\varepsilon_k} = \text{div}(j(u_{\varepsilon_k})),$$

so we are led to

$$\text{div}(j(u_{\varepsilon_k})) = k_{\varepsilon_k} u_{\varepsilon_k} \times \partial_t u_{\varepsilon_k} + \frac{1}{2} \varepsilon_k \frac{d}{dt} \left(\frac{|u_{\varepsilon_k}|^2 - 1}{\varepsilon_k} \right). \quad (5.13)$$

Now, applying Lemma 3 to u_{ε_k} , we find

$$\sup_{[0, T]} E_{\varepsilon_k}(u_{\varepsilon_k}(t), B) \leq \pi l |\log \varepsilon| + \Sigma_{\varepsilon_k}(t) + C \leq \pi l |\log \varepsilon_k| + C, \quad (5.14)$$

where the second inequality is itself a consequence of (5.4). This implies first that $|u_{\varepsilon_k}| \rightarrow 1$ in $L^2([0, T] \times B)$. Moreover, we infer that the second term in the r.h.s of (5.13) converges to zero in the distribution sense on $[0, T] \times B$. For the first one, it suffices to use Cauchy-Schwarz inequality combined with the L^2 bound provided by Lemma 6 and the already mentioned uniform bounds of $|u_{\varepsilon_k}|$ in L^2_{loc} .

We then infer from Lemma 8 and (5.14) that $j(u_{\varepsilon_k})$ is uniformly bounded in $L^p_{\text{loc}}([0, T] \times \mathbb{R}^2)$ for all $p < 2$. This is e.g. a consequence of Theorem 3.2.1 in [9] and the remarks that follow. We deduce from (5.11) and (5.12) that up to a subsequence, we have

$$j(u_{\varepsilon_k}) \rightharpoonup j_1 = j(u^*(b_i(\cdot), d_i)) + H \quad (5.15)$$

weakly in $L^p_{\text{loc}}([0, T] \times \mathbb{R}^2)$, where H is harmonic in x on $[0, T] \times \mathbb{R}^2$.

On the other hand, it follows from the first part of Lemma 9 that there exists j_2 such that, taking subsequences if necessary, $j(u_{\varepsilon_k})/|u_{\varepsilon_k}| \rightharpoonup j_2$ weakly in $L^2_{\text{loc}}(\mathcal{G})$.

Taking into account the strong convergence $|u_{\varepsilon_k}| \rightarrow 1$ in $L^2_{\text{loc}}([0, T] \times \mathbb{R}^2)$, we obtain $j_1 = j_2 \in L^2_{\text{loc}}(\mathcal{G})$. The second part of Lemma 9 combined with (5.15) then yields

$$\|H\|_{L^2_{\text{loc}}(\mathcal{G})} \leq \liminf_{k \rightarrow +\infty} \left\| \frac{j(u_{\varepsilon_k})}{|u_{\varepsilon_k}|} - j(u^*(b_i, d_i)) \right\|_{L^2_{\text{loc}}(\mathcal{G})} \leq CT,$$

where C depends only on K_1 , R and r_a , so finally $\|H\|_{L^2([0,T] \times \mathbb{R}^2)} \leq CT$. Since H is harmonic in x , we find that $H(t, \cdot)$ is bounded on \mathbb{R}^2 for almost every t and therefore is identically zero. We end up with $j_1 = j_2 = j(u^*(b_i(\cdot), d_i))$ in \mathcal{G} , and the conclusion follows. \square

6. PROOF OF THEOREM 2

In this section, we present the proof of Theorem 2. We let $\{b_i(t)\}$ be the l Lipschitz paths on $[0, T]$ provided by Theorem 5 and $\{a_i(t)\}$ be the unique maximal solution defined on $I = [0, T^*)$ to (3) with initial conditions a_i^0 . Our aim is to show that $a_i(t) \equiv b_i(t)$ on I . We will first prove that this holds on $[0, T]$. By Rademacher's Theorem, the time derivatives $\dot{b}_i(t)$ exist and are bounded almost everywhere on $[0, T]$. Without loss of generality, we may assume $T < T^*$, so that

$$|\dot{a}_i(t)| \leq C, \quad |\dot{b}_i(t)| \leq C, \quad \text{a.e. on } [0, T]. \quad (6.1)$$

Moreover, we may assume, decreasing possibly T , that $|a_i(t) - b_i(t)| \leq r_a/2$ for all i . Hence, the trajectories $a_i(t)$ remain in $B(a_i^0, r_a)$ on $[0, T]$. We introduce

$$h(t) = \sum_{i=1}^l \int_0^t |\dot{a}_i(s) - \dot{b}_i(s)| ds, \quad \sigma(t) = \sum_{i=1}^l |a_i(t) - b_i(t)|,$$

then h is Lipschitz on $[0, T]$ and for almost every $t \in [0, T]$ we have $h'(t) = \sum_{i=1}^l |\dot{a}_i(t) - \dot{b}_i(t)|$. Note that since σ is absolutely continuous and $\sigma(0) = 0$, we have for all $t \in [0, T]$

$$\sigma(t) = \int_0^t \sigma'(s) ds \leq h(t)$$

therefore it suffices to show that h is identically zero on $[0, T]$. This will be done by mean of Gronwall's Lemma.

Lemma 10. *For all $t_1, t_2, t \in [0, T]$, we have*

$$\limsup_{k \rightarrow +\infty} \Sigma_{\varepsilon_k}(t) \leq Ch(t)$$

and

$$\limsup_{k \rightarrow +\infty} \int_{t_1}^{t_2} \Sigma_{\varepsilon_k}(s) ds \leq C \int_{t_1}^{t_2} h(s) ds,$$

where C only depends on r_a, K_0, R_a .

Proof. For $t \in [0, T]$, we decompose $\Sigma_{\varepsilon_k}(t)$ as

$$\begin{aligned} \Sigma_{\varepsilon_k}(t) &= \mathcal{E}_{\varepsilon_k, [U_d]}(u_{\varepsilon_k}(t)) - \mathcal{E}_{\varepsilon_k, [U_d]}(u_{\varepsilon_k}^0) + \Sigma_{\varepsilon_k}(0) \\ &\quad + \mathcal{E}_{\varepsilon_k, [U_d]}(u_{\varepsilon_k}^*(a_i^0, d_i)) - \mathcal{E}_{\varepsilon_k, [U_d]}(u_{\varepsilon_k}^*(b_i(t), d_i)). \end{aligned}$$

Appealing to Corollary 1 and Theorem 1, we obtain

$$\Sigma_{\varepsilon_k}(t) = -\delta \int_0^t \int_{\mathbb{R}^2} \frac{|\partial_t u_{\varepsilon_k}|^2}{|\log \varepsilon_k|} + \Sigma_{\varepsilon_k}(0) + W(a_i^0, d_i) - W(b_i(t), d_i) + o_{\varepsilon_k}(1).$$

Using that W is Lipschitz away from zero, we estimate the last term as follows

$$\begin{aligned} W(a_i^0, d_i) - W(b_i(t), d_i) &= W(a_i^0, d_i) - W(a_i(t), d_i) + W(a_i(t), d_i) - W(b_i(t), d_i) \\ &\leq - \int_0^t \sum_{i=1}^l \dot{a}_i(s) \cdot \nabla_{a_i} W(s) ds + C\sigma(t). \end{aligned}$$

Since the a_i solve the Cauchy problem (3), an explicit computation gives

$$\dot{a}_i(s) \cdot \nabla_{a_i} W(s) = \frac{\delta}{\pi} C_i d_i |\nabla_{a_i} W|^2 = -\delta\pi |\dot{a}_i(s)|^2,$$

so that

$$\Sigma_{\varepsilon_k}(t) \leq \Sigma_{\varepsilon_k}(0) + \delta\pi \int_0^t \sum_{i=1}^l |\dot{a}_i(s)|^2 ds - \delta \int_0^t \int_{\mathbb{R}^2} \frac{|\partial_t u_{\varepsilon_k}|^2}{|\log \varepsilon_k|} + C\sigma(t) + o_{\varepsilon_k}(1).$$

For the energy dissipation in the right-hand side, we need a lower bound as ε_k tends to zero. In view of the convergence of the Jacobians (5.1) and the upper bound for the energy $\sup_{t \in [0, T]} E_{\varepsilon_k}(u_{\varepsilon_k}(t), B(R')) \leq \pi l |\log \varepsilon_k| + C$ stated in (5.14), Proposition 3 in [11] (see also Corollary 7 in [20]) provides the lower mobility bound

$$\liminf_{k \rightarrow +\infty} \int_0^t \int_{\mathbb{R}^2} \frac{|\partial_t u_{\varepsilon_k}|^2}{|\log \varepsilon_k|} \geq \pi \sum_{i=1}^l \int_0^t |\dot{b}_i(t)|^2 ds. \quad (6.2)$$

Now, we have thanks to (6.1)

$$\sum_{i=1}^l \int_0^t (|\dot{a}_i(s)|^2 - |\dot{b}_i(s)|^2) \leq C \sum_{i=1}^l \int_0^t |\dot{a}_i(s) - \dot{b}_i(s)| ds = Ch(t),$$

whereas $\Sigma_{\varepsilon_k}(0) \rightarrow 0$ by assumption, hence we get

$$\limsup_{k \rightarrow +\infty} \Sigma_{\varepsilon_k}(t) \leq C(\sigma(t) + h(t)).$$

Applying Fatou's Lemma in (6.2) finally also yields the corresponding integral version, and lastly, it suffices to use that $\sigma \leq h$. \square

As suggested in the introduction, the map $u^*(a_i(t), d_i)$ solves the evolution formula provided by Proposition 1 in the asymptotics $\varepsilon \rightarrow 0$.

Lemma 11. *We have for $t \in [0, T]$ and $\chi, \varphi \in \mathcal{H}(a_i^0)$*

$$\pi \frac{d}{dt} \sum_{i=1}^l d_i [\chi(a_i(t)) + \delta\varphi(a_i(t))] = -2 \int_{\mathbb{R}^2} \operatorname{Im} \left(\omega(u^*(a_i(t), d_i)) \frac{\partial^2 \chi}{\partial \bar{z}^2} \right).$$

Proof. We use the following formula proved in [5], valid for any configuration (a_i, d_i) and any test function χ which is affine near the point vortices:

$$-2 \int_{\mathbb{R}^2} \operatorname{Im} \left(\omega(u^*(a_i(t), d_i)) \frac{\partial^2 \chi}{\partial \bar{z}^2} \right) = -2\pi \sum_{i \neq j} d_i d_j \frac{(a_i(t) - a_j(t))^\perp}{|a_i(t) - a_j(t)|^2} \cdot \nabla \chi(a_i(t)).$$

On the other hand, we compute

$$\begin{aligned} \frac{d}{dt} \sum_{i=1}^l [d_i \chi(a_i) + \delta \varphi(a_i)] &= \sum_{i=1}^l [d_i \nabla \chi(a_i^0) \cdot \dot{a}_i(t) + \delta \nabla \varphi(a_i^0) \cdot \dot{a}_i(t)] \\ &= \sum_{i=1}^l d_i \nabla \chi(a_i^0) \cdot (\dot{a}_i(t) - \delta d_i \dot{a}_i^\perp(t)), \end{aligned}$$

where the second equality follows from the relation $\nabla \varphi(a_i^0) = \nabla^\perp \chi(a_i^0)$. Next, we deduce from (3)

$$\pi \left(\dot{a}_i(t) - \delta d_i \dot{a}_i^\perp(t) \right) = -C_i (1 + \delta^2 d_i^2) \nabla_{a_i}^\perp W = d_i \nabla_{a_i}^\perp W,$$

and we obtain

$$\begin{aligned} \pi \frac{d}{dt} \sum_{i=1}^l [d_i \chi(a_i) + \delta \varphi(a_i)] &= \sum_{i=1}^l \nabla \chi(a_i) \cdot \nabla_{a_i}^\perp W \\ &= -2\pi \sum_{i \neq j} d_i d_j \frac{(a_i - a_j)^\perp}{|a_i - a_j|^2} \cdot \nabla \chi(a_i), \end{aligned}$$

which yields the conclusion. \square

Lemma 12. *Set $A = \cup_{i=1}^l B(a_i^0, 2r_a) \setminus B(a_i^0, r_a)$ and let $t_1, t_2 \in [0, T]$. Then for all $\varphi \in \mathcal{D}(A)$, we have*

$$\limsup_{k \rightarrow +\infty} \left| \int_{t_1}^{t_2} \int_A [\omega(u_{\varepsilon_k}(s)) - \omega(u^*(b_i(s), d_i))] \varphi \right| \leq C \|\varphi\|_\infty \int_{t_1}^{t_2} h(s) ds.$$

Proof. We apply the pointwise equality (2.4) to $u \equiv u_{\varepsilon_k}(t)$ and $u^* \equiv u^*(b_i(t), d_i)$ for all t . Since $|u^*(b_i(t), d_i)| = 1$, this gives

$$\omega(u) - \omega(u^*) = \sum_{k,l=1}^2 \left(a_{k,l} \partial_l |u| \partial_k |u| + b_{k,l} \left[\frac{j_k(u)}{|u|} \frac{j_l(u)}{|u|} - j_k(u^*) j_l(u^*) \right] \right),$$

where $a_{k,l}, b_{k,l} \in \mathbb{C}$. We rewrite the terms involving the components of j as

$$\begin{aligned} \frac{j_k(u)}{|u|} \frac{j_l(u)}{|u|} - j_k(u^*) j_l(u^*) &= \left(\frac{j_k(u)}{|u|} - j_k(u^*) \right) \left(\frac{j_l(u)}{|u|} - j_l(u^*) \right) \\ &\quad + j_k(u^*) \left(\frac{j_l(u)}{|u|} - j_l(u^*) \right) + j_l(u^*) \left(\frac{j_k(u)}{|u|} - j_k(u^*) \right). \end{aligned}$$

We multiply the previous equality by φ , integrate on $[t_1, t_2] \times A$ and let k go to $+\infty$. Using the weak convergence in L^2 of $j(u_{\varepsilon_k})/|u_{\varepsilon_k}|$ to $j(u^*(b_i(\cdot), d_i))$ on $[0, T] \times A \subset \mathcal{G}$, we deduce that

$$\begin{aligned} \limsup_{k \rightarrow +\infty} \left| \int_{t_1}^{t_2} \int_A [\omega(u_{\varepsilon_k}(s)) - \omega(u^*(b_i(s), d_i))] \varphi \right| \\ \leq C \|\varphi\|_\infty \limsup_{k \rightarrow +\infty} \int_{t_1}^{t_2} \int_A [|\nabla |u_{\varepsilon_k}||^2 + \left| \frac{j(u_{\varepsilon_k})}{|u_{\varepsilon_k}|} - j(u^*(b_i, d_i)) \right|^2]. \end{aligned}$$

The conclusion finally follows from Lemmas 9 and 10. \square

We are now in position to complete the proof of Theorem 2. We consider arbitrary χ, φ belonging to $\mathcal{H}(a_i^0)$, we fix $0 \leq s \leq t \leq T$ and we integrate the evolution formula given by Proposition 1 on $[s, t]$. We obtain

$$\int_s^t \frac{d}{d\tau} \int_{\mathbb{R}^2} J u_{\varepsilon_k}(\tau) \chi + \delta \int_{\mathbb{R}^2} \mu_{\varepsilon_k}(\tau) \varphi = \int_s^t g_k^1(\tau) + \int_s^t g_k^2(\tau),$$

where

$$g_k^1(\tau) = -\delta \int_{\mathbb{R}^2} \frac{|\partial_t u_{\varepsilon_k}|^2}{|\log \varepsilon_k|^2} + R_{\varepsilon_k}(\tau, \chi, \varphi, u_{\varepsilon_k})$$

and

$$g_k^2(\tau) = -2 \int_{\mathbb{R}^2} \operatorname{Im} \left(\omega(u_{\varepsilon_k}(\tau)) \frac{\partial^2 \chi}{\partial \bar{z}^2} \right),$$

which we decompose as

$$\begin{aligned} g_k^2 &= -2 \int_{\mathbb{R}^2} \operatorname{Im} \left([\omega(u_{\varepsilon_k}) - \omega(u^*(b_i, d_i))] \frac{\partial^2 \chi}{\partial \bar{z}^2} \right) \\ &\quad - 2 \int_{\mathbb{R}^2} \operatorname{Im} \left([\omega(u^*(b_i, d_i)) - \omega(u^*(a_i, d_i))] \frac{\partial^2 \chi}{\partial \bar{z}^2} \right) \\ &\quad - 2 \int_{\mathbb{R}^2} \operatorname{Im} \left(\omega(u^*(a_i, d_i)) \frac{\partial^2 \chi}{\partial \bar{z}^2} \right) = A_k(\tau) + B_k(\tau) + C_k(\tau). \end{aligned}$$

We next substitute the formula given by Lemma 11 for C_k in the previous equalities. Setting

$$f_{k, \chi, \varphi}(\tau) = \int_{\mathbb{R}^2} J u_{\varepsilon_k}(\tau) \chi + \delta \int_{\mathbb{R}^2} \mu_{\varepsilon_k}(\tau) \varphi - \pi \sum_{i=1}^l [d_i \chi(a_i(\tau)) + \delta \varphi(a_i(\tau))],$$

we obtain

$$f_{k, \chi, \varphi}(t) - f_{k, \chi, \varphi}(s) = \int_s^t g_k^1 + \int_s^t A_k + \int_s^t B_k.$$

Lemma 6 with $T_\varepsilon = T$ first gives $|\int_s^t g_k^1(\tau) d\tau| \leq C |\log \varepsilon_k|^{-\frac{1}{2}}$ for all k . Moreover, it follows from Lemma 12 and the fact that $\operatorname{supp} \frac{\partial^2 \chi}{\partial \bar{z}^2} \subset A$ that

$$\limsup_{k \rightarrow +\infty} \left| \int_s^t A_k(\tau) d\tau \right| \leq C \int_s^t h(\tau) d\tau.$$

Finally, we infer from the regularity of $\omega(u^*)$ away from the vortices that

$$\int_s^t |B_k(\tau)| d\tau \leq C \int_s^t \sigma(\tau) d\tau \leq C \int_s^t h(\tau) d\tau.$$

Letting k go to $+\infty$, we finally deduce from the convergence statements in Theorem 5 that for $0 \leq s \leq t \leq T$,

$$|f_{\chi, \varphi}(t) - f_{\chi, \varphi}(s)| \leq C \int_s^t h(\tau) d\tau, \quad (6.3)$$

where $f_{\chi, \varphi}$ is defined by

$$f_{\chi, \varphi} = \pi \sum_{i=1}^l \left[d_i (\chi(b_i) - \chi(a_i)) + \delta (\varphi(b_i) - \varphi(a_i)) \right].$$

Here the constant C depends only on χ, φ and the initial conditions.

We now fix a time $t \in [0, T]$ at which all the vortices b_i have a time derivative. Since the a_i are C^1 , it follows that $f_{\chi, \varphi}$ is differentiable at t with time derivative given by

$$f'_{\chi, \varphi}(t) = \pi \sum_{i=1}^l \left(d_i \nabla \chi(a_i^0) + \delta \nabla^\perp \chi(a_i^0) \right) \cdot (\dot{b}_i(t) - \dot{a}_i(t)).$$

Dividing by $t - s$ in (6.3) and letting $s \rightarrow t$ gives then

$$\left| \pi \sum_{i=1}^l \left(d_i \nabla \chi(a_i^0) + \delta \nabla^\perp \chi(a_i^0) \right) \cdot (\dot{b}_i(t) - \dot{a}_i(t)) \right| \leq C h(t).$$

So, considering in particular $\chi, \varphi \in \mathcal{H}(a_i^0)$ such that χ and φ vanish near each point a_i^0 except for one, we obtain for all $i = 1, \dots, l$

$$\left| \pi \left(d_i \nabla \chi(a_i^0) + \delta \nabla^\perp \chi(a_i^0) \right) \cdot (\dot{b}_i(t) - \dot{a}_i(t)) \right| \leq C h(t).$$

Choosing then successively $\chi(x) = x_1$ and $\chi(x) = x_2$ near a_i^0 we end up with $|\dot{b}_i(t) - \dot{a}_i(t)| \leq C h(t)$, and it follows by summation

$$h'(t) \leq C h(t) \quad \text{a.e. } t \in [0, T].$$

Since $h(0) = 0$, this implies that $h = 0$ on $[0, T]$, and hence $\sigma = 0$ on $[0, T]$. Applying Lemma 10, we infer that $\limsup_{k \rightarrow +\infty} \Sigma_{\varepsilon_k}(t) \leq 0$. Besides, Lemma 3 yields for all $L \geq 2^{n_1}$

$$\begin{aligned} \liminf_{k \rightarrow +\infty} \Sigma_{\varepsilon_k}(t) &\geq \liminf_{k \rightarrow +\infty} \int_{B(L)} \left[e_{\varepsilon_k}(u_{\varepsilon_k}(t)) - e_{\varepsilon_k}(u_{\varepsilon_k}^*(a_i(t), d_i)) \right] - \frac{C}{L} \\ &\geq -\frac{C}{L}, \end{aligned}$$

where the second inequality is a consequence of the convergence of Jacobians on $B(L)$ stated in Lemma 8 (see [14, 18]). Letting L tend to $+\infty$, we obtain $\liminf_{k \rightarrow +\infty} \Sigma_{\varepsilon_k}(t) \geq 0$, so we deduce from (5.3) that $(u_{\varepsilon_k}(t))_{k \in \mathbb{N}}$ is well-prepared with respect to the configuration $(a_i(t), d_i)$. By uniqueness of the limit, this finally holds for the full family $(u_\varepsilon(t))_{0 < \varepsilon < 1}$ on $[0, T]$.

In conclusion, we observe that in our definition T only depends on K_1 , r_a and $\max(R, R_a + r_a)$, so that we can extend our results to the whole of $[0, T^*)$ by repeating the previous arguments.

APPENDIX: CAUCHY PROBLEM FOR $(\text{CGL})_\varepsilon$

We present here the proof of Theorem 1. We omit the dependence on ε and rewrite (2) in the following form

$$\begin{cases} \partial_t w = (a + ib)(\Delta w + f_{U_0}(w)), \\ w(0) = w_0 \in H^1(\mathbb{R}^2), \end{cases} \quad (\text{CGL})$$

where

$$f_{U_0}(w) = \Delta U_0 + (U_0 + w)(1 - |U_0 + w|^2),$$

a is positive and $b \in \mathbb{R}$. We denote by $S = S(t, x)$ the semi-group operator associated to the corresponding homogeneous linear equation. Every

solution $w \in C^0([0, T], H^1(\mathbb{R}^2))$ to (CGL) satisfies the Duhamel formula

$$w(t, \cdot) = S(t, \cdot) * w_0 + \int_0^t S(t-s, \cdot) * g_{U_0}(w(s), \cdot) ds, \quad s \in [0, T],$$

where $g_{U_0} = (a + ib)f_{U_0}$. The kernel S is explicitly given by

$$S(t, x) = \frac{1}{4\pi(a + ib)t} \exp\left(\frac{-|x|^2}{4(a + ib)t}\right).$$

Since a is positive, S decays at infinity like the standard heat Kernel, therefore (CGL) enjoys the same smoothing properties as the parabolic Ginzburg-Landau equation. In particular, we have for all $1 \leq r \leq +\infty$ and for all $t > 0$

$$\|S(t, \cdot)\|_{L^r(\mathbb{R}^2)} \leq \frac{C_{a,b}}{t^{1-\frac{1}{r}}} \quad (\text{a})$$

and concerning the space derivatives of $S(t)$,

$$\|D^k S(t, \cdot)\|_{L^r(\mathbb{R}^2)} \leq \frac{C_{a,b}}{t^{\frac{|k|}{2}+1-\frac{1}{r}}}. \quad (\text{b})$$

We will often use Young's inequality, which gives for $f \in L^p(\mathbb{R}^2)$ and $g \in L^q(\mathbb{R}^2)$ $\|f * g\|_{L^r(\mathbb{R}^2)} \leq \|f\|_{L^p(\mathbb{R}^2)} \|g\|_{L^q(\mathbb{R}^2)}$, where $1 + \frac{1}{r} = \frac{1}{p} + \frac{1}{q}$.

We first state local well-posedness for (CGL).

Proposition A.1. *Let $w_0 \in H^1(\mathbb{R}^2)$. Then there exists a positive time T^* depending on $\|w_0\|_{H^1}$ and a unique solution $w \in C^0([0, T^*), H^1(\mathbb{R}^2))$ to (CGL).*

Proof. We intend to apply the fixed point theorem to the map $\psi : w \in H^1(\mathbb{R}^2) \mapsto \psi(w)$, where

$$\psi(w)(t) = S(t) * w_0 + \int_0^t S(t-s) * g_{U_0}(w(s)) ds.$$

To this aim, we introduce $R = \|w_0\|_{H^1(\mathbb{R}^2)}$ and for $T > 0$

$$B(T, R) = \{w \in L^\infty([0, T], H^1(\mathbb{R}^2)) \text{ s.t. } \|w\|_{L^\infty(H^1)} \leq (2C_{a,b} + 1)R\},$$

where $C_{a,b}$ is the constant appearing in (a)-(b). We next show that we can choose $T = T(R)$ so that ψ maps $B(T(R), R)$ into itself and is a contraction on this ball.

For $T > 0$, we let $w \in B(T, R)$ and expand $f_{U_0}(w)$. Using that $H^1(\mathbb{R}^2)$ is continuously embedded in $L^p(\mathbb{R}^2)$ for all $2 \leq p < +\infty$ and that U_0 belongs to \mathcal{V} , it can be shown that ⁵

$$\|f_{U_0}\|_{L^\infty([0, T], L^2)} \leq C(U_0, R), \quad (\text{c})$$

and for $w_1, w_2 \in B(T, R)$

$$\|f_{U_0}(w_1) - f_{U_0}(w_2)\|_{L^\infty([0, T], L^2)} \leq C(U_0, R) \|w_1 - w_2\|_{L^\infty([0, T], H^1)}. \quad (\text{d})$$

⁵see Lemma 1 in [7].

We next apply Young's inequality to obtain

$$\begin{aligned} \|\psi(w)(t)\|_{H^1} &\leq \|\psi(w)(t)\|_{L^2} + \|\nabla\psi(w)(t)\|_{L^2} \\ &\leq 2\|S(t)\|_{L^1}\|w_0\|_{H^1} + \int_0^t \|S(t-s) + \nabla S(t-s)\|_{L^1}\|g_{U_0}(s)\|_{L^2} ds \\ &\leq 2C_{a,b}\|w_0\|_{H^1} + C \int_0^t \left(1 + (t-s)^{-\frac{1}{2}}\right) \|g_{U_0}(w(s))\|_{L^2} ds, \end{aligned}$$

where the last inequality is a consequence of (a) and (b) with the choice $r = 1$. This yields according to (c) and (d)

$$\sup_{t \in [0, T]} \|\psi(w)(t)\|_{H^1} \leq 2C_{a,b}\|w_0\|_{H^1} + C(U_0, R)(T + \sqrt{T})$$

and similarly,

$$\sup_{t \in [0, T]} \|\psi(w_1)(t) - \psi(w_2)(t)\|_{H^1} \leq C'(U_0, R)(T + \sqrt{T}) \sup_{t \in [0, T]} \|w_1(t) - w_2(t)\|_{H^1}.$$

The conclusion follows by choosing $T = T(R)$ sufficiently small so that $C(U_0, R)(T + \sqrt{T}) \leq R$ and $C'(U_0, R)(T + \sqrt{T}) < 1$. \square

We next show additional regularity for a solution to (CGL).

Lemma A.1. *Let $w \in C^0([0, T], H^1(\mathbb{R}^2))$ be a solution to (CGL). Then w belongs to $L^1_{\text{loc}}([0, T], H^2(\mathbb{R}^2)) \cap C^0((0, T], H^2(\mathbb{R}^2))$ and therefore to $L^1_{\text{loc}}([0, T], L^\infty(\mathbb{R}^2))$.*

Proof. We first differentiate $f_{U_0}(w)$ and use Lemma 2 in [7] which states by means of various Sobolev embeddings, Hölder and Gagliardo-Nirenberg inequalities that

$$\partial_i f_{U_0}(w) = g_1(w) + g_2(w) \in L^\infty([0, T], L^2(\mathbb{R}^2)) + L^\infty([0, T], L^r(\mathbb{R}^2))$$

for all $1 < r < 2$. Moreover, we have

$$\sup_{s \in [0, T]} (\|g_1(w)(s)\|_{L^2(\mathbb{R}^2)} + \|g_2(w)(s)\|_{L^r(\mathbb{R}^2)}) \leq C(U_0, A(T), r),$$

where $A(T) = \sup_{s \in [0, T]} \|w(s)\|_{H^1(\mathbb{R}^2)}$. Next, differentiating twice Duhamel formula gives

$$\partial_{ij} w(t) = \partial_j S(t) * \partial_i w_0 + \int_0^t \partial_j S(t-s) * \partial_i f_{U_0}(s) ds,$$

so taking into account the decomposition $\partial_i f_{U_0} = g_1 + g_2$ we get

$$\begin{aligned} \|\partial_{ij} w(t)\|_{L^2} &\leq \|\nabla S(t)\|_{L^1} \|\nabla w_0\|_{L^2} + \int_0^t \|\nabla S(t-s)\|_{L^1} \|g_1(s)\|_{L^2} ds \\ &\quad + \int_0^t \|\nabla S(t-s)\|_{L^\alpha} \|g_2(s)\|_{L^r} ds, \end{aligned}$$

where α is chosen so that $1 + \frac{1}{2} = \frac{1}{\alpha} + \frac{1}{r}$. This finally yields in view of (b)

$$\begin{aligned} \|\partial_{ij} w(t)\|_{L^2} &\leq Ct^{-\frac{1}{2}} \|w_0\|_{H^1} \\ &\quad + C(U_0, A(T), r) \int_0^t \left((t-s)^{-\frac{1}{2}} + (t-s)^{-\frac{1}{2}-1+\frac{1}{\alpha}} \right) ds. \end{aligned}$$

Since $\frac{1}{2} + 1 - \frac{1}{\alpha} = \frac{1}{r} < 1$, we conclude that the right-hand side is finite, so that $\partial_{ij} w \in L^1_{\text{loc}}([0, T], L^2(\mathbb{R}^2))$. \square

Lemma A.1 enables to show that the renormalized energy is non-increasing and hence to control $\|w(t)\|_{H^1(\mathbb{R}^2)}$. For equation (CGL), this energy is given by

$$E_{U_0}(w)(t) = \int_{\mathbb{R}^2} \frac{|\nabla w|^2}{2} - \int_{\mathbb{R}^2} \Delta U_0 \cdot w + \int_{\mathbb{R}^2} \frac{(1 - |U_0 + w|^2)^2}{4}.$$

It is well-defined and continuous in time for $w \in C^0(H^1(\mathbb{R}^2))$.

Lemma A.2. *Let $w \in C^0([0, T], H^1(\mathbb{R}^2))$ be a solution to (CGL). Then for all $t \in (0, T)$ we have*

$$\frac{d}{dt} E_{U_0}(w)(t) \leq 0.$$

Moreover, there exists C_{U_0, w_0} depending only on U_0 and $\|w_0\|_{H^1}$ such that

$$\|w(t)\|_{H^1} \leq C_{U_0, w_0} \exp(C_{U_0, w_0} t), \quad \forall t \in [0, T]. \quad (\text{e})$$

Proof. We infer from equation (CGL) and Lemma A.1 that $\partial_t w$ belongs to $L_{\text{loc}}^\infty((0, T], L^2(\mathbb{R}^2))$, so that we may compute

$$\begin{aligned} \frac{d}{dt} E_{U_0}(w(t)) &= \int_{\mathbb{R}^2} \nabla w \cdot \nabla \partial_t w - \Delta U_0 \cdot \partial_t w - \partial_t w \cdot (U_0 + w)(1 - |U_0 + w|^2) \\ &= - \int_{\mathbb{R}^2} \partial_t w \cdot (\Delta w + f_{U_0}(w)) \\ &= - \int_{\mathbb{R}^2} \partial_t w \cdot \left(\frac{1}{a + ib} \partial_t w \right) \\ &= \frac{-a}{a^2 + b^2} \int_{\mathbb{R}^2} |\partial_t w|^2 \leq 0. \end{aligned}$$

We now turn to (e). We compute for $t \in (0, T)$

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|w(t)\|_{L^2(\mathbb{R}^2)}^2 &= \int_{\mathbb{R}^2} w \cdot \partial_t w \\ &= \int_{\mathbb{R}^2} w \cdot [(a + ib)\Delta w] + \int_{\mathbb{R}^2} w \cdot [(a + ib)f_{U_0}(w)] \\ &= -a \int_{\mathbb{R}^2} |\nabla w|^2 + \int_{\mathbb{R}^2} w \cdot (a + ib)\Delta U_0 \\ &\quad + \int_{\mathbb{R}^2} w \cdot [(a + ib)(U_0 + w)(1 - |U_0 + w|^2)]. \end{aligned}$$

We then split the last term in the previous equality as

$$\begin{aligned} &\int_{\mathbb{R}^2} w \cdot [(a + ib)(U_0 + w)(1 - |U_0 + w|^2)] \\ &= \int_{\mathbb{R}^2} w \cdot [(a + ib)U_0(1 - |U_0 + w|^2)] + a \int_{\mathbb{R}^2} |w|^2(1 - |U_0 + w|^2). \end{aligned}$$

The second term in the r.h.s. is clearly bounded by $a\|w(t)\|_{L^2(\mathbb{R}^2)}^2$. Using Cauchy-Schwarz inequality for the first one, we obtain

$$\int_{\mathbb{R}^2} w \cdot [(a + ib)(U_0 + w)(1 - |U_0 + w|^2)] \leq C(U_0)\|w(t)\|_{L^2} V(t)^{\frac{1}{2}} + a\|w(t)\|_{L^2}^2,$$

where $V(t) = \int_{\mathbb{R}^2} (1 - |U_0 + w(t)|^2)^2$. We are led to

$$\frac{d}{dt} \|w(t)\|_{L^2(\mathbb{R}^2)}^2 \leq C(U_0)(\|w(t)\|_{L^2}^2 + 1 + V(t)). \quad (\text{f})$$

On the other hand, Cauchy-Schwarz inequality gives

$$E_{U_0}(w)(t) \geq \int_{\mathbb{R}^2} \frac{|\nabla w|^2}{2} dx - C(U_0)\|w(t)\|_{L^2} + \frac{V(t)}{4},$$

which yields, since $E_{U_0}(w)$ is non-increasing,

$$\frac{V(t)}{4} + \int_{\mathbb{R}^2} \frac{|\nabla w|^2}{2} \leq E_{U_0}(w_0) + C(U_0)\|w(t)\|_{L^2}. \quad (\text{g})$$

We infer from (f) and (g)

$$\|w(t)\|_{L^2} \leq (1 + \|w_0\|_{H^1}) \exp(Ct)$$

and finally deduce (e) by using (g) once more. \square

Lemma A.2 provides global well-posedness for (CGL).

Proposition A.2. *Let $w_0 \in H^1(\mathbb{R}^2)$. Then there exists a unique and global solution $w \in C^0(\mathbb{R}_+, H^1(\mathbb{R}^2))$ to (CGL).*

Proof. Let $w \in C^0([0, T^*), H^1(\mathbb{R}^2))$ be the unique maximal solution with initial condition w_0 . If T^* is finite, we have according to (e)

$$\limsup_{t \rightarrow T^*} \|w(t)\|_{H^1(\mathbb{R}^2)} \leq C(U_0, T^*, w_0) < +\infty,$$

so that we can extend w to a solution \bar{w} on $[0, T^* + \delta]$ for some positive δ . This yields a contradiction. \square

We conclude this section with the following

Proposition A.3. *Let $w \in C^0(\mathbb{R}_+, H^1(\mathbb{R}^2))$ be the solution to (CGL). Then we have $w \in C^\infty(\mathbb{R}_+, C^\infty(\mathbb{R}^2))$.*

Proof. We proceed in several steps.

Step 1 Let $p \geq 2$ and $v \in H^p(\mathbb{R}^2)$. Then $D^k f_{U_0}(v) \in L^2(\mathbb{R}^2) + L^{\frac{4}{3}}(\mathbb{R}^2)$ for all $|k| \leq p$.

Proof of Step 1. We may assume in view of the proof of Lemma A.1 that $|k| \geq 2$. We decompose $f_{U_0}(v)$ as $f_{U_0}(v) = \Delta U_0 + h_{U_0}(v)$, where

$$h_{U_0}(v) = (U_0 + v)(1 - |U_0 + v|^2).$$

Since $U_0 \in \mathcal{V}$, it suffices to show that $D^k h_{U_0}(v) \in L^2(\mathbb{R}^2) + L^{\frac{4}{3}}(\mathbb{R}^2)$. Applying Leibniz's formula to $h_{U_0}(v)$, we obtain

$$\begin{aligned} D^k h_{U_0}(v) &= \sum_{m \leq k} \binom{k}{m} D^{k-m}(U_0 + v) D^m(1 - |U_0 + v|^2) \\ &= D^k(U_0 + v) \\ &\quad - \sum_{\substack{m \leq k \\ n \leq m}} \binom{k}{m} \binom{m}{n} D^{k-m}(U_0 + v) D^n(U_0 + v) \cdot D^{m-n}(U_0 + v). \end{aligned}$$

Since $2 \leq |k| \leq p$, $v \in H^p(\mathbb{R}^2)$ and $U_0 \in \mathcal{V}$, we clearly have $D^k(U_0 + v) \in L^2(\mathbb{R}^2)$.

For the second term in the right-hand side, we write each product inside the sum as

$$D^a(U_0 + v)D^b(U_0 + v) \cdot D^c(U_0 + v)$$

with $|a| + |b| + |c| = |k| \geq 2$, and we examine all cases. We observe that $D^a(v + U_0)$ belongs to $H^1(\mathbb{R}^2)$ whenever $1 \leq |a| \leq p-1$ and hence to $L^4(\mathbb{R}^2)$, whereas $D^a(v + U_0)$ belongs to $L^2(\mathbb{R}^2)$ for $2 \leq |a| \leq p$. Since on the other hand $U_0 + v \in L^\infty$, we finally obtain

$$D^a(U_0 + v)D^b(U_0 + v) \cdot D^c(U_0 + v) \in L^2(\mathbb{R}^2) + L^{\frac{4}{3}}(\mathbb{R}^2),$$

which yields the conclusion.

We now turn to the regularity in space for a solution to (CGL).

Step 2 Let $w \in C^0(\mathbb{R}_+, H^1(\mathbb{R}^2))$ be the solution to (CGL). Then for all $p \geq 1$ we have $w \in C^0(\mathbb{R}_+^*, H^p(\mathbb{R}^2))$.

Proof of Step 2. We proceed by induction on p . The case $p = 2$ has already been treated in Lemma A.1. Let us thus assume that $w \in C^0(\mathbb{R}_+^*, H^p(\mathbb{R}^2))$ for some $p \geq 2$. For $|k| \leq p + 1$, we differentiate $w(t)$ and we find

$$D^k w(t) = D^k(S(t) * w_0) + D^k \int_0^t S(t-s) * g_{U_0}(s) ds$$

which we rewrite as

$$\begin{aligned} D^k w(t) &= D^k S(t) * w_0 + \int_0^{t/2} (D^k S(t-s)) * g_{U_0}(s) ds \\ &\quad + \int_{t/2}^t D^m S(t-s) * D^{k-m} g_{U_0}(s) ds, \end{aligned}$$

where m is a multi-index so that $|m| = 1$.

First, it follows from (b) that $t \mapsto D^k S(t) * w_0 \in C^0(\mathbb{R}_+^*, L^2(\mathbb{R}^2))$. Next, arguing that $g_{U_0} \in C^0(\mathbb{R}_+, L^2(\mathbb{R}^2))$ and using (b) with $r = 1$, we find

$$\left\| \int_0^{t/2} (D^k S(t-s)) * g_{U_0}(s) ds \right\|_{L^2} \leq C \int_0^{t/2} \frac{ds}{(t-s)^{\frac{|k|}{2}}} \leq \frac{C}{t^{\frac{|k|}{2}-1}}.$$

On the other hand, since $|k-m| = |k| - 1 \leq p$ and since by assumption $w(s) \in H^p(\mathbb{R}^2)$, Step 1 provides the decomposition

$$D^{k-m} g_{U_0}(s) = d^1(s) + d^2(s)$$

where d^1 belongs to $C^0(\mathbb{R}_+^*, L^2(\mathbb{R}^2))$ and d^2 to $C^0(\mathbb{R}_+^*, L^{\frac{4}{3}}(\mathbb{R}^2))$. It follows from (b) that

$$\begin{aligned} &\left\| \int_{t/2}^t D^m S(t-s) * D^{k-m} g_{U_0}(s) ds \right\|_{L^2} \\ &\leq \int_{t/2}^t \left(\|\nabla S(t-s)\|_{L^1} \|d^1(s)\|_{L^2} + \|\nabla S(t-s)\|_{L^r} \|d^2(s)\|_{L^{\frac{4}{3}}} \right) ds \\ &\leq C(t) \int_{t/2}^t \left((t-s)^{-\frac{1}{2}} + (t-s)^{-\frac{1}{2}-1+\frac{1}{r}} \right) ds, \end{aligned}$$

where r satisfies $1 + \frac{1}{2} = \frac{1}{r} + \frac{3}{4}$. The last term is finite since $\frac{1}{2} + 1 - \frac{1}{r} = \frac{3}{4} < 1$, so we infer that $w \in C^0(\mathbb{R}_+^*, H^{p+1}(\mathbb{R}^2))$, as we wanted.

Step 3 Let $w \in C^0(\mathbb{R}_+, H^1(\mathbb{R}^2))$ be the solution to (CGL). Then we have $w \in C^k(\mathbb{R}_+^*, C^l(\mathbb{R}^2))$ for all $k, l \in \mathbb{N}$.

Proof of Step 3. For fixed $k, l \in \mathbb{N}$, we show by induction on $0 \leq j \leq k$ that $w \in C^j(\mathbb{R}_+^*, C^{l+2k-2j}(\mathbb{R}^2))$.

This holds for $j = 0$ according to Step 2 and to Sobolev embeddings. We assume next that $w \in C^j(\mathbb{R}_+^*, C^{l+2k-2j}(\mathbb{R}^2))$ for some $0 \leq j \leq k - 1$, and it follows that

$$\Delta w, f_{U_0}(w) \in C^j(\mathbb{R}_+^*, C^{l+2k-2j-2}(\mathbb{R}^2)).$$

So, going finally back to equation (CGL), we obtain

$$w \in C^{j+1}(\mathbb{R}_+^*, C^{l+2k-2j-2}(\mathbb{R}^2)).$$

This concludes the proof of Proposition A.3. □

Acknowledgements. I warmly thank Didier Smets for his constant support during the preparation of this work. I am also indebted to Thierry Gallay and to Sylvia Serfaty for very helpful discussions.

This work was partly supported by the grant JC05-51279 of the Agence Nationale de la Recherche.

REFERENCES

- [1] L. Almeida, *Threshold transition energies for Ginzburg-Landau functionals*, Nonlinearity **12** (1999), 1389-1414.
- [2] I. S. Aranson and L. Kramer, *The world of the complex Ginzburg-Landau equation*, Rev. Mod. Phys. **74** (2002), 99-143.
- [3] F. Bethuel, H. Brezis and F. Hélein, *Ginzburg-Landau vortices*, Birkhäuser, Boston, 1994.
- [4] F. Bethuel, R. L. Jerrard and D. Smets, *On the NLS dynamics for infinite energy vortex configurations on the plane*, Rev. Mat. Iberoamericana **24** (2008), 671-702.
- [5] F. Bethuel, G. Orlandi and D. Smets, *Collisions and phase-vortex interactions in dissipative Ginzburg-Landau dynamics*, Duke Math. J. **130** (2005), 523-614.
- [6] F. Bethuel, G. Orlandi and D. Smets, *Dynamics of multiple degree Ginzburg-Landau vortices*, Comm. Math. Phys. **272** (2007), 229-261.
- [7] F. Bethuel and D. Smets, *A remark on the Cauchy Problem for the 2D Gross-Pitaevskii equation with non zero degree at infinity*, Differential Integral Equations **20** (2007), 325-338.
- [8] J. E. Colliander and R. L. Jerrard, *Vortex dynamics for the Ginzburg-Landau-Schrödinger equation*, Internat. Math. Res. Notices **7** (1998), 333-358.
- [9] J. E. Colliander and R. L. Jerrard, *Ginzburg-Landau vortices: weak stability and Schrödinger equation dynamics*, J. Anal. Math. **77** (1999), 129-205.
- [10] J. Ginibre and G. Velo, *The Cauchy problem in local spaces for the complex Ginzburg-Landau equation. II. Contraction methods*, Comm. Math. Phys. **187** (1997), no.1, 45-79.
- [11] R. L. Jerrard, *Vortex dynamics for the Ginzburg-Landau wave equation*, Calc. Var. PDE **9** (1999), no.1, 1-30.
- [12] R. L. Jerrard and H. M. Soner, *Dynamics of Ginzburg-Landau vortices*, Arch. Ration. Mech. Anal. **142** (1998), 99-125.
- [13] R. L. Jerrard and H. M. Soner, *The Jacobian and the Ginzburg-Landau energy*, Calc. Var. PDE **14** (2002), 141-191.
- [14] R. L. Jerrard and D. Spirn, *Refined Jacobian estimates for Ginzburg-Landau functionals*, Indiana Univ. Math. Jour. **56** (2007), 135-186.
- [15] R. L. Jerrard and D. Spirn, *Refined Jacobian estimates and Gross-Pitaevsky vortex dynamics*, Arch. Ration. Mech. Anal. **190** (2008), no. 3, 425-475.

- [16] M. Kurzke, C. Melcher, R. Moser and D. Spirn, *Dynamics for Ginzburg-Landau vortices under a mixed flow*, Indiana Univ. Math. J. (2008), to appear.
- [17] F. H. Lin, *Some dynamical properties of Ginzburg-Landau vortices*, Comm. Pure Appl. Math. **49** (1996), 323-359.
- [18] F. H. Lin and J. X. Xin, *On the incompressible fluid limit and the vortex motion law of the nonlinear Schrödinger equation*, Comm. Math. Phys. **200** (1999), 249-274.
- [19] S. Serfaty, *Vortex collisions and energy-dissipation rates in the Ginzburg-Landau heat flow, part II: The dynamics*, Journal Eur. Math Society **9** (2007), no. 3, 383-426.
- [20] E. Sandier and S. Serfaty, *A product-estimate for Ginzburg-Landau and corollaries*, J. of Funct. An. **211** (2004), 219-244.

(E. Miot) LABORATOIRE J.-L. LIONS UMR 7598, UNIVERSITÉ PIERRE ET MARIE CURIE, 175 RUE DU CHEVALERET, 75013 PARIS, FRANCE
E-mail address: miot@ann.jussieu.fr