

Incompressible Euler and e-MHD as scaling limits of the Vlasov-Maxwell system

Yann BRENIER ^{*}
Norbert J. MAUSER [†]
Marjolaine PUEL [‡]

June 1, 2003

Abstract

We consider two different asymptotic limits of the Vlasov-Maxwell system describing a quasineutral plasma with a uniform ionic background. In the first case, as the magnetic field is preserved in the limiting process, we obtain the so-called electron magnetohydrodynamics equations. In the second case, we obtain the incompressible Euler equations with no more magnetic field left.

1 Introduction

Let us consider the Vlasov-Maxwell system

$$\partial_t f + \xi \cdot \nabla_x f - (E + \alpha(\xi \wedge B)) \cdot \nabla_\xi f = 0, \quad (1)$$

$$\alpha \partial_t B + \nabla \wedge E = 0 \quad , \quad \varepsilon \nabla \cdot E = 1 - \rho, \quad (2)$$

$$\varepsilon \alpha \partial_t E - \nabla \wedge B = \alpha J \quad , \quad \nabla \cdot B = 0, \quad (3)$$

This system (1)–(3) describes the evolution of the electron phase density $f(t, x, \xi) \geq 0$ at time $t > 0$ and point $x \in \mathbb{R}^3$, with velocity $\xi \in \mathbb{R}^3$, in a uniform background of non-moving ions with unit density. The fields E and B are respectively the electric field and the magnetic field. The density ρ and the current J are given by

$$\rho = \int f(t, x, \xi) d\xi \quad , \quad J = \int \xi f(t, x, \xi) d\xi. \quad (4)$$

^{*}Lab. J.A Dieudonné, Parc Valrose, F-06100 Nice, France.

[†]Wolfgang Pauli Inst. c/o Inst. f. Math., Univ Wien, Strudlhofg. 4, A-1090 Wien, Austria.

[‡]Laboratoire d'analyse numérique, université Pierre et Marie Curie, BC 187 F-75252 PARIS Cedex 05, France.

E and B are coupled to the electron density through the Maxwell equations and act on electrons via the Lorentz force. Notice that we do not consider here the *relativistic* Vlasov-Maxwell system for which the impulse variable ξ and the velocity variable are not proportional. (This means that the speed of light must be infinite at leading order in any reasonable scaling.) After non-dimensionalization, parameters α and ϵ can be chosen independently on each other, according to the desired scaling. Indeed, in physical units

$$\alpha = \sqrt{\frac{r_0}{\epsilon_0}}, \quad \epsilon = \frac{\epsilon_0}{r_0 c^2},$$

where r_0 is the classical electron radius ($r_0 = 2.82 \times 10^{-15} m$), ϵ_0 is the vacuum dielectric constant and c is the speed of light.

We will concentrate on the so-called quasi-neutral regime when ϵ is a small parameter.

Taking the first two moments of (1), we obtain the following system,

$$\partial_t \rho + \nabla \cdot J = 0 \tag{5}$$

$$\partial_t J + \nabla \cdot \left(\int \xi \otimes \xi f d\xi \right) + E\rho + \alpha J \wedge B = 0 \tag{6}$$

Remark 1.1 Here notation $\nabla \cdot \left(\int \xi \otimes \xi f d\xi \right)$ corresponds to

$$\sum_{j=1}^3 \partial_j \int \xi_i \xi_j f d\xi, \quad i = 1, 2, 3.$$

Starting from this system, we may consider two different quasi-neutral scalings.

Scaling 1: $\alpha \ll 1, \epsilon \ll 1$

This scaling corresponds to a limit, without any magnetic field, coupled with a quasi-neutral limit (the electron density instantaneously adjusts itself to the unit background density). Formally, it is easy to guess the limit of (5),(6), (2),(3) when $\epsilon, \alpha \rightarrow 0$, provided that the distribution function tends to be monokinetic, i.e.

$$f(t, x, \xi) \sim \rho(t, x) \delta\left(\xi - \frac{J(t, x)}{\rho(t, x)}\right),$$

which means that the electronic temperature is close to zero and implies

$$\int \xi \otimes \xi f d\xi \sim \frac{J \otimes J}{\rho}.$$

This limit is the Euler equation of incompressible fluids

$$\partial_t J + \nabla \cdot (J \otimes J) + E = 0, \tag{7}$$

$$\nabla \cdot J = 0 \quad , \quad \rho = 1, \tag{8}$$

$$\nabla \wedge E = 0 \quad , \quad B = 0. \tag{9}$$

Scaling 2: $\alpha = 1$, $\varepsilon \ll 1$

Now, the magnetic field does not disappear any longer in the limit equations, the so-called electron magnetohydrodynamics equations [KCY]:

$$\partial_t J + \nabla \cdot (J \otimes J) + E + J \wedge B = 0, \quad (10)$$

$$J = -\nabla \wedge B \quad , \quad \rho = 1, \quad (11)$$

$$\partial_t B + \nabla \wedge E = 0, \quad \nabla \cdot B = 0. \quad (12)$$

As above, provided that the distribution function tends to be monokinetic, the formal limit is obvious.

In the present paper, we provide a rigorous derivation of these formal limits by using the modulated energy method designed in [Br] for the quasi-neutral limit of the Vlasov-Poisson system. This method has been used in a quantum context in [Pu1], [Pu2] and based on the concept of dissipative solutions due to P.-L. Lions [Li]. This method can be seen as a variant of both the relative entropy method [Da], [Ya] and the hamiltonian energy method by E. Grenier [Gr].

The key idea is to estimate a modulation of the energy by the solution of the formal limit equation. Typically, we replace (twice) the kinetic energy $\int |\xi|^2 f d\xi$ by $\int |\xi - v|^2 f d\xi$ where v is a smooth solution to the limit equations.

The Vlasov-Maxwell system is known to admit global strong solutions at least for smooth initial data depending only on two space coordinates, with a perpendicular magnetic field (see [GIsc] for details and [DiLi] for global weak solutions). To keep the proofs as simple as possible, we state our results in the case of spatially periodic solutions on the unit cube. All spatial integrals will be implicitly performed on the unit cube.

2 Convergence to the electron-Magnetohydrodynamics (e-MHD)

2.1 Result

In this section, we consider the Vlasov-Maxwell system (1), (2), (3), (4), where $\alpha > 0$ is a fixed constant of order one and ε is a small parameter. (This means that the magnetic field will not vanish in the limiting process.) The system that we consider for the limit $\varepsilon \rightarrow 0$, is the so-called e-MHD system [KCY]

$$\partial_t v + (v \cdot \nabla)v + e + \alpha v \wedge b = 0 \quad (13)$$

$$\text{with } \alpha \partial_t b + \nabla \wedge e = 0, \quad \alpha v = \nabla \wedge b \text{ and } \nabla \cdot b = 0.$$

Introducing $\omega = \nabla \wedge (v - \alpha A)$ where A is the magnetic potential such that $\nabla \wedge A = b$ and $\nabla \cdot A = 0$, and using identity

$$(v \cdot \nabla)v = (\nabla \wedge v) \wedge v + \nabla \frac{|v|^2}{2},$$

we can write e-MHD in a different way :

$$\partial_t \omega + v \cdot \nabla \omega = \omega \cdot \nabla v, \text{ with } -\Delta v + \alpha^2 v = \nabla \wedge \omega.$$

In particular, as the initial data depend only on the two first coordinates x_1, x_2 , then ω is aligned with the third coordinate and can be seen as a scalar field satisfying

$$\partial_t \omega + v \cdot \nabla \omega = 0, \text{ with } -\Delta v + \alpha^2 v = {}^\perp \nabla \omega = (-\partial_1 \omega_2, \partial_2 \omega_1, 0),$$

Therefore, the existence results are the same as for the incompressible Euler equations (see [Li], for instance). In particular, we have global strong solutions for smooth initial conditions depending only on two space coordinates. Let us now state our first main result

Theorem 2.1 *Let us fix $T > 0$, $\alpha > 0$ and $C > 0$. Let (f, B, E) and (v, b, e) be two smooth solutions of respectively the VM system and the e-MHD system on the time interval $[0, T]$. Assume their initial values to satisfy*

$$\int f(0, x, \xi) dx d\xi = 1, \tag{14}$$

$$\int f(0, x, \xi) |\xi|^2 dx d\xi \leq C, \tag{15}$$

$$\varepsilon \int |E(0, x)|^2 dx \leq C\sqrt{\varepsilon}, \tag{16}$$

$$\int |B(0, x) - b(0, x)|^2 dx \leq C\sqrt{\varepsilon}, \tag{17}$$

$$\int |\xi - v(0, x)|^2 f(0, x, \xi) dx d\xi \leq C\sqrt{\varepsilon}. \tag{18}$$

Then

$$\sup_{0 \leq t \leq T} \left(\int |\xi - v(t, x)|^2 f(t, x, \xi) d\xi dx + \int |B(t, x) - b(t, x)|^2 dx \right) \leq C' \sqrt{\varepsilon},$$

where C' depends only on T, C and (v, b, e) .

2.2 Moment equations and modulated energy

We deduce from the Vlasov system the following equations for the moments ρ and J :

$$\partial_t \rho + \nabla \cdot J = 0 \tag{19}$$

and

$$\partial_t J + \nabla \cdot \left(\int \xi \otimes \xi f d\xi \right) + \rho E + \alpha J \wedge B = 0$$

and the conserved energy is

$$H = \frac{1}{2} \int |\xi|^2 f(t, x, \xi) d\xi dx + \int \frac{\varepsilon |E(t, x)|^2 + |B(t, x)|^2}{2} dx.$$

Following [Br], we define a 'modulated energy' $H_{v,b}(t)$ by

$$H_{v,b}(t) := \frac{1}{2} \int |\xi - v(t, x)|^2 f(t, x, \xi) d\xi dx + \int \frac{\varepsilon |E(t, x)|^2 + |B(t, x) - b(t, x)|^2}{2} dx$$

where v and b are arbitrarily chosen smooth divergence free vector fields depending on $t \geq 0$ and $x \in \mathbb{R}^3/\mathbb{Z}^3$.

Proposition 2.2 *Let (f, B, E) be a smooth solution to the VM system and v, b, e three smooth vector fields, b and v being divergence free, then the modulated energy $H_{v,b}$ satisfies*

$$\frac{d}{dt}(H_{v,b} - \theta) = \int \sum_{i,j=1}^3 \partial_j v_i [- \int (\xi - v)_i (\xi - v)_j f d\xi + (B - b)_i (B - b)_j + \varepsilon E_i E_j] dx + \eta + r \quad (20)$$

where

$$\theta = \varepsilon \int E \cdot (e + \alpha v \wedge (b - B)) dx$$

$$\eta = -\varepsilon \int E \cdot (\partial_t e + \nabla(e \cdot v) + \alpha \partial_t (v \wedge b) - \alpha \partial_t v \wedge B) dx,$$

and

$$\begin{aligned} r = & - \int (e - E - \alpha v \wedge B) \cdot \left(\frac{\nabla \wedge b}{\alpha} + v \right) dx \\ & - \int (B - b) \cdot \left(\partial_t b + \frac{\nabla \wedge e}{\alpha} \right) dx - \int ((\partial_t + v \cdot \nabla)v + e + \alpha v \wedge b) \cdot (J - \rho v) dx. \end{aligned}$$

Theorem 2.1 follows easily from Proposition 2.2. Indeed, we first assume (v, b, e) to be solution to the e-MHD equation. As a consequence, r vanishes in the right-hand side of (20). Next, it follows from the assumptions of Theorem 2.1 that $\eta(t)$ and $\theta(t)$ are uniformly bounded by $C_1 \sqrt{\varepsilon}$, for $t \in [0, T]$ where C_1 depends only on T, C and (v, b, e) . Thus, we get from (20),

$$\frac{d}{dt}(H_{v,b} - \theta) \leq 2\lambda H_{v,b} + \eta \leq 2\lambda(H_{v,b} - \theta) + 2\lambda\theta + \eta \leq 2\lambda(H_{v,b} - \theta + C_2 \sqrt{\varepsilon}),$$

where

$$\lambda = \sup_{t,x,\xi} \sum_{i,j=1}^3 \partial_j v_i(t, x) \frac{\xi_i \xi_j}{|\xi|^2}$$

and C_2 depends only on T, C and (v, b, e) . Thus

$$H_{v,b}(t) - \theta(t) + C_2 \sqrt{\varepsilon} \leq \exp(2\lambda t) (H_{v,b}(0) - \theta(0) + C_2 \sqrt{\varepsilon}).$$

From the assumptions of Theorem 2.1, we know that $H_{v,b}(0) \leq 1.5C \sqrt{\varepsilon}$ and conclude that

$$H_{v,b}(t) \leq C' \sqrt{\varepsilon}$$

where C' depends only on T, C and (v, b, e) and the result of Theorem 2.1 easily follows.

2.3 Proof of Proposition 2.2

N.B. To perform the calculations required by the proof, we use notations with indices $i, j, k \in \{1, 2, 3\}$, implicit summation on repeated indices and abridged notations with comma for partial derivatives. For instance, $v_{,t} + v_{i,j}v_j$ stands for $\partial_t v + (v \cdot \nabla)v$. Consistently, we use the signature symbol $\epsilon_{ijk} = +1, -1$ to denote the wedge product and the curl operator. With these notations, the equations satisfied by (ρ, J, E, B) are

$$\rho_{,t} = -J_{j,j}, \quad (21)$$

$$-J_{i,t} = \left(\int \xi_i \xi_j f d\xi \right)_{,j} + \alpha \epsilon_{ijk} J_j B_k + \rho E_i, \quad (22)$$

$$\alpha B_{i,t} = -\epsilon_{ijk} E_{k,j}, \quad B_{i,i} = 0, \quad (23)$$

$$\alpha J_i = \varepsilon \alpha E_{i,t} - \epsilon_{ijk} B_{k,j}, \quad \rho = 1 - \varepsilon E_{i,i}. \quad (24)$$

Let us notice that we can write equation (22) in a different way, using equations (23) and (24), namely :

$$\begin{aligned} -J_{i,t} &= \left(\int \xi_i \xi_j f d\xi \right)_{,j} - \epsilon_{ijk} \epsilon_{jpk} B_{q,p} B_k + \varepsilon \alpha \epsilon_{ijk} E_{j,t} B_k + E_i (1 - \varepsilon E_{j,j}) \\ &= \left(\int \xi_i \xi_j f d\xi \right)_{,j} + B_k B_{k,i} - B_k B_{i,k} + \varepsilon \alpha \epsilon_{ijk} (E_j B_k)_{,t} + \varepsilon \alpha \epsilon_{ijk} E_j B_{k,t} + E_i (1 - \varepsilon E_{j,j}) \\ &= \left(\int \xi_i \xi_j f d\xi \right)_{,j} + B_j (B_{j,i} - B_{i,j}) + \varepsilon \alpha \epsilon_{ijk} (E_j B_k)_{,t} + \varepsilon \epsilon_{ijk} \epsilon_{kpq} E_j E_{q,p} + E_i (1 - \varepsilon E_{j,j}) \end{aligned}$$

(using (23)). Thus,

$$-J_{i,t} = \left(\int \xi_i \xi_j f d\xi \right)_{,j} + B_j (B_{j,i} - B_{i,j}) + \varepsilon \alpha \epsilon_{ijk} (E_j B_k)_{,t} + \varepsilon E_j (E_{j,i} - E_{i,j}) + E_i (1 - \varepsilon E_{j,j}). \quad (25)$$

The modulated energy is defined by

$$H_{b,v} = \frac{1}{2} \left(\int \int |\xi - v|^2 f d\xi + \int |B - b|^2 + \int \varepsilon |E|^2 \right). \quad (26)$$

We have

$$H_{b,v} = H + \int (-J_i v_i + \rho v_i v_i / 2) + \int (-B_i b_i + b_i b_i / 2)$$

where the energy

$$H = H_{0,0} = \frac{1}{2} \left(\int \int |\xi|^2 f d\xi + \int |B|^2 + \int \varepsilon |E|^2 \right)$$

is conserved and therefore does not depend on t . Thus

$$\frac{dH_{b,v}}{dt} = Q_1 + Q_2 + Q_3 + L_1 + L_2 + L_3, \quad (27)$$

where

$$Q_1 = - \int v_i J_{i,t}, \quad Q_2 = \int \rho_{,t} v_i v_i / 2, \quad Q_3 = - \int B_{i,t} b_i, \quad (28)$$

$$L_1 = - \int J_i v_{i,t}, \quad L_2 = \int \rho v_i v_{i,t}, \quad L_3 = - \int (B - b)_i b_{i,t}. \quad (29)$$

From equation (25), we deduce

$$\begin{aligned} Q_1 &= \int v_i \left(\int \xi_i \xi_j f d\xi \right)_{,j} + \int v_i B_j (B_{j,i} - B_{i,j}) + \varepsilon \alpha \int \epsilon_{ijk} v_i (E_j B_k)_{,t} \\ &\quad + \varepsilon \int v_i E_j (E_{j,i} - E_{i,j}) + \int v_i E_i (1 - \varepsilon E_{j,j}) \\ &= \int v_i \left(\int \xi_i \xi_j f d\xi \right)_{,j} + \int v_i \left(\left(\frac{|B|^2}{2} \right)_{,i} - B_j B_{i,j} \right) + \varepsilon \alpha \int \epsilon_{ijk} v_i (E_j B_k)_{,t} \\ &\quad + \varepsilon \int v_i \left(\left(\frac{|E|^2}{2} \right)_{,i} - (E_i E_j)_{,j} \right) + \int v_i E_i. \end{aligned}$$

Thus, after integrating by part and using that $v_{j,j} = B_{j,j} = 0$,

$$Q_1 = \int v_{i,j} \left(- \int \xi_i \xi_j f d\xi + B_j B_i + \varepsilon E_j E_i \right) + \int v_i E_i + \eta_0 + \frac{d}{dt} \theta_0,$$

where

$$\theta_0 = \varepsilon \alpha \int \epsilon_{ijk} v_i E_j B_k, \quad \eta_0 = -\varepsilon \alpha \int \epsilon_{ijk} v_{i,t} E_j B_k.$$

Next, from equation (21),

$$Q_2 = \int \rho_{,t} v_i v_i / 2 = - \int J_{j,j} v_i v_i / 2 = \int J_j v_i v_{i,j}$$

and, from (23),

$$Q_3 = - \int B_{k,t} b_k = \int \epsilon_{kji} \frac{E_{i,j}}{\alpha} b_k = \int E_i \epsilon_{ijk} \frac{b_{k,j}}{\alpha}.$$

So

$$\begin{aligned} Q = Q_1 + Q_2 + Q_3 &= \int v_{i,j} \left(- \int \xi_i \xi_j f d\xi + B_i B_j + \varepsilon E_i E_j \right) + \int J_j v_i v_{i,j} + \eta_0 + \frac{d}{dt} \theta_0 \\ &\quad + r_0 \end{aligned}$$

where

$$r_0 = \int E_i \left(\epsilon_{ijk} \frac{b_{k,j}}{\alpha} + v_i \right).$$

(Notice that $r_0 = 0$ as (b, v, e) satisfies the e-MHD system.) We also have

$$L_3 = - \int (B - b)_i b_{i,t} = \int (B - b)_i \frac{\epsilon_{ijk} e_{k,j}}{\alpha} + r_1,$$

where

$$r_1 = - \int (B - b)_i (b_{i,t} + \frac{\epsilon_{ijk} e_{k,j}}{\alpha}).$$

(Notice that $r_1 = 0$ as (b, v, e) satisfies the e-MHD system.) Thus

$$\begin{aligned} L_3 &= - \int e_k \epsilon_{ijk} \frac{(B - b)_{i,j}}{\alpha} + r_1 \\ &= - \int e_k (J_k - \epsilon E_{k,t} - v_k) + r_1 + r_2, \end{aligned}$$

(because of equation (24)), where

$$r_2 = - \int e_k (v_k + \frac{\epsilon_{kji} b_{i,j}}{\alpha}).$$

(Notice that $r_2 = 0$ if (b, v, e) is solution to the e-MHD system.) So

$$\begin{aligned} L_3 &= - \int e_k (J_k - \rho v_k) - \int e_k (\rho - 1) v_k + \frac{d}{dt} \epsilon \int E_k e_k - \epsilon \int E_k e_{k,t} + r_1 + r_2 \\ &= - \int e_k (J_k - \rho v_k) + \eta_1 + \eta_2 + \frac{d}{dt} \theta_1 + r_1 + r_2 \end{aligned}$$

(using $\rho = 1 - \epsilon E_{i,i}$) where

$$\begin{aligned} \eta_1 &= -\epsilon \int (e_k v_k)_{,i} E_i, \\ \eta_2 &= -\epsilon \int E_k e_{k,t} \\ \theta_1 &= \epsilon \int E_k e_k. \end{aligned}$$

It follows, using definition (29), that

$$L = L_1 + L_2 + L_3 = - \int (v_{k,t} + e_k) (J_k - \rho v_k) + \eta_1 + \eta_2 + \frac{d}{dt} \theta_1 + r_1 + r_2$$

Thus

$$L = \int v_{i,j} (v_j J_i - \rho v_j v_i) + L_4 + \eta_1 + \eta_2 + \frac{d}{dt} \theta_1 + r_1 + r_2 + r_3$$

where

$$r_3 = - \int (v_{i,t} + v_j v_{i,j} + e_i + \alpha \epsilon_{ijk} v_j b_k) (J_i - \rho v_i)$$

(which vanishes if (b, v, e) is a solution to the e-MHD system), and

$$L_4 = \int \alpha \epsilon_{ijk} v_j b_k (J_i - \rho v_i)$$

$$= \int \alpha \epsilon_{ijk} v_j b_k J_i = \varepsilon \alpha \int \epsilon_{ijk} v_j b_k E_{i,t} - \int \epsilon_{ijk} v_j b_k \epsilon_{ipq} B_{q,p}$$

(because of (24))

$$= \eta_3 + \frac{d}{dt} \theta_2 + L_5$$

where

$$\eta_3 = -\varepsilon \alpha \int \epsilon_{ijk} (v_j b_k)_t E_i,$$

$$\theta_2 = \varepsilon \alpha \int \epsilon_{ijk} v_j b_k E_i,$$

$$L_5 = - \int v_j b_k (B_{k,j} - B_{j,k}).$$

We have

$$L_5 = \int v_j b_{k,j} B_k - \int v_{j,k} b_k B_j$$

(since both v and b are divergence free)

$$= - \int v_{j,k} (b_k B_j + B_k b_j) + \int v_j (b_{k,j} - b_{j,k}) B_k$$

$$= - \int v_{j,k} (b_k B_j + B_k b_j) + \int \epsilon_{ijk} \epsilon_{ipq} b_{q,p} v_j B_k$$

$$= - \int v_{j,k} (b_k B_j + B_k b_j) + r_4,$$

where

$$r_4 = \int \epsilon_{ijk} (\epsilon_{ipq} b_{q,p} + \alpha v_i) v_j B_k$$

(and vanishes if (b, v, e) is solution to the e-MHD system). So

$$L_4 = \eta_3 + \frac{d}{dt} \theta_2 - \int v_{j,k} (b_k B_j + B_k b_j) + r_4$$

and

$$L = \int v_{i,j} (v_j J_i - \rho v_j v_i - b_i B_j - B_i b_j) \\ + \eta_1 + \eta_2 + \eta_3 + \frac{d}{dt} (\theta_1 + \theta_2) + r_1 + r_2 + r_3 + r_4$$

It follows, using definition (28), that

$$L + Q = Q' + \eta_0 + \eta_1 + \eta_2 + \eta_3 + \frac{d}{dt} (\theta_0 + \theta_1 + \theta_2) + r_0 + r_1 + r_2 + r_3 + r_4,$$

where

$$Q' = \int v_{i,j}((B-b)_j(B-b)_i + \varepsilon E_j E_i) - \int (\xi_i - v_i)(\xi_j - v_j) f d\xi.$$

So, we have finally obtained

$$\frac{d}{dt}(H_{b,v} - \theta) = Q' + \eta + r,$$

where

$$\eta = \eta_0 + \eta_1 + \eta_2 + \eta_3 = -\varepsilon \int \epsilon_{ijk} \alpha v_{i,t} E_j B_k - \varepsilon \int E_i (e_{i,t} + (e_k v_k)_{,i} + \alpha \epsilon_{ijk} (v_j b_k)_{,t}),$$

$$\theta = \theta_0 + \theta_1 + \theta_2 = \varepsilon \alpha \int \epsilon_{ijk} v_i E_j B_k, + \varepsilon \int E_i (e_i + \alpha \epsilon_{ijk} v_j b_k)$$

and

$$\begin{aligned} r = r_0 + r_1 + r_2 + r_3 + r_4 &= \int E_i (\epsilon_{ijk} \frac{b_{k,j}}{\alpha} + v_i) - \int (B-b)_i (b_{i,t} + \frac{\epsilon_{ijk} e_{k,j}}{\alpha}) \\ &- \int e_k (v_k + \frac{\epsilon_{kji} b_{i,j}}{\alpha}) - \int (v_{i,t} + v_j v_{i,j} + e_i + \alpha \epsilon_{ijk} v_j b_k) (J_i - \rho v_i) \\ &+ \int \epsilon_{ijk} (\epsilon_{ipq} b_{q,p} + \alpha v_i) v_j B_k \\ &= - \int (e_i - E_i - \epsilon_{ipq} \alpha v_p B_q) (\epsilon_{ijk} \frac{b_{k,j}}{\alpha} + v_i) \\ &- \int (B-b)_i (b_{i,t} + \frac{\epsilon_{ijk} e_{k,j}}{\alpha}) - \int (v_{i,t} + v_j v_{i,j} + e_i + \alpha \epsilon_{ijk} v_j b_k) (J_i - \rho v_i). \end{aligned}$$

This completes the proof of Proposition 2.2.

3 Convergence to the Euler equation

3.1 Result

In this section, we consider the Vlasov-Maxwell system (1), (2), (3), (4), with a different scaling where both α and ε go to zero. The limit system is now the usual Euler equations for incompressible fluids

$$\partial_t v + (v \cdot \nabla) v + e = 0 \tag{30}$$

$$\text{with } \nabla \wedge e = 0, \text{ and } \nabla \cdot v = 0.$$

Let us recall that, by introducing $\omega = \nabla \wedge v$, we get the standard vorticity formulation of the Euler equations, namely :

$$\partial_t \omega + v \cdot \nabla \omega = \omega \cdot \nabla v, \text{ with } -\Delta v = \nabla \wedge \omega.$$

Let us now state our second main result

Theorem 3.1 *Let us fix $T > 0$ and $C > 0$. Let (f, B, E) and (v, e) be two smooth solutions of respectively the VM system and the Euler system on the time interval $[0, T]$. For each $t \in [0, T]$, assume v to have zero mean and introduce β such that*

$$\nabla \wedge \beta = v, \quad \nabla \cdot \beta = 0.$$

Assume the initial values to satisfy

$$\int f(0, x, \xi) dx d\xi = 1, \tag{31}$$

$$\int f(0, x, \xi) |\xi|^2 dx d\xi \leq C, \tag{32}$$

$$\varepsilon \int |E(0, x)|^2 dx \leq (\alpha + \sqrt{\varepsilon})C, \tag{33}$$

$$\int |B(0, x) - \alpha\beta(0, x)|^2 dx \leq (\alpha + \sqrt{\varepsilon})C, \tag{34}$$

$$\int |\xi - v(0, x)|^2 f(0, x, \xi) dx d\xi \leq (\alpha + \sqrt{\varepsilon})C. \tag{35}$$

Then

$$\sup_{0 \leq t \leq T} \left(\int |\xi - v(t, x)|^2 f(t, x, \xi) d\xi dx + \int |B(t, x) - \alpha\beta(t, x)|^2 dx \right) \leq (\alpha + \sqrt{\varepsilon})C',$$

where C' depends only on T, C and (v, e) .

3.2 Proof of Theorem 3.1

We use again Proposition 2.2, but with a different choice for (v, b, e) . We assume (v, e) to be the smooth solution of the Euler equation considered in the assumptions of Theorem 3.1, and define

$$b(t, x) = \alpha\beta(t, x), \quad \nabla \wedge \beta = v, \quad \nabla \cdot \beta = 0.$$

This implies that

$$\theta = \varepsilon \int E \cdot (e + \alpha v \wedge (\alpha\beta - B))$$

$$\eta = -\varepsilon \int E \cdot (\partial_t e + \nabla(e \cdot v) + \alpha^2 \partial_t (v \wedge \beta) - \alpha \partial_t v \wedge B),$$

and

$$r = - \int (B - \alpha\beta) \cdot \alpha \partial_t \beta - \int \alpha^2 v \wedge \beta \cdot (J - \rho v).$$

(using that $\nabla \wedge \beta = v$ and (v, e) is a solution to the Euler equations which in particular implies $\nabla \wedge e = 0$). Thus, η, θ and r are uniformly bounded by $(\alpha + \sqrt{\varepsilon})C''$ for some constant C'' depending only on $T, (b, v)$ and the total energy H . Then, the proof of Theorem 3.1 immediately follows as in the previous section for Theorem 2.1.

Acknowledgments

The authors acknowledge the support of the European network IHP network "HYKE" HPRN-CT-2002-00282, the LRC Cea-Cadarache/UNSA, the START project (FWF, TEC-Y-137) of N.J.M and the Austrian-French program "Amadeus". Y.B. thanks D. Holm for informing him about e-MHD equations [Ho].

References

- [Br] Y. Brenier, Convergence of the Vlasov-Poisson system to the incompressible Euler equations, *Comm. PDEs* (3-4), (2000), 737-754.
- [BrMaPu] Y. Brenier, N. J. Mauser and M. Puel, "Sur les limites quasineutres/non-relativistes: de Vlasov-Maxwell vers Euler incompressible", *C. R. Acad. Sci.*, t 334, Série I Math., 1-6, 2001.
- [Da] C. Dafermos, *Hyperbolic conservation laws in continuum physics, chapt. 5-2, Grundlehren der Mathematischen Wissenschaften, 325, Springer-Verlag, Berlin, 2000.*
- [DiLi] R.J. Di Perna and P.L. Lions, "Global Weak Solutions of Vlasov-Maxwell Systems", *Comm. Pure and Appl. Math* 42 (1989) 729-757.
- [GlSc] R. Glassey, J. Schaeffer, The relativistic Vlasov-Maxwell system in two space dimensions, *Arch. Rational Mech. Anal.* 141 (1998) 331-354.
- [Gr] E. Grenier, Boundary layers of 2D inviscid fluids from a Hamiltonian viewpoint, *Math. Res. Lett.* 6 (1999) 257-269.
- [Ho] D. D. Holm, "Notes on Electron MHD vs Euler-alpha", private communication to Y.B.
- [KCY] A. Kingsep, K. Chukbar, V. Yankov **Electron Magnetohydrodynamics**, *Review of Plasma Physics 16, ed. by B.B. Kadomtsev, New York, 1990.*
- [Li] P.L. Lions, **Mathematical topics in fluid mechanics, vol. 1. incompressible models**, *Oxford Lecture in Mathematics and its Applications*, Oxford Univ. Press (1996).
- [Pu1] M. Puel, Convergence of the Schrödinger Poisson system to the incompressible Euler equations, submitted (2001).
- [Pu2] M. Puel, Convergence of the Schrödinger Poisson system to the Euler equations under the influence of a strong magnetic field, submitted (2001).
- [Ya] H.T. Yau, Relative entropy and hydrodynamics of Ginzburg-Landau models, *Lett. Math. Phys.*, 22(1) (1991) 63-80.